



Publication Year	2015
Acceptance in OA	2020-04-03T17:25:35Z
Title	VizieR Online Data Catalog: Photometry and abundances of NGC1851 stars (Marino+, 2014)
Authors	Marino, A. F., Milone, A. P., Yong, D., Dotter, A., Da Costa, G., Asplund, M., Jerjen, H., Mackey, D., Norris, J., CASSISI, Santi, Sbordone, L., Stetson, P. B., Weiss, A., Aparicio, A., BEDIN, Luigi, Lind, K., Monelli, M., Piotto, G., Angeloni, R., Buonanno, R.
Handle	http://hdl.handle.net/20.500.12386/23822
Journal	VizieR Online Data Catalog

VizieR Online Data Catalog: Photometry and abundances of NGC1851 stars (Marino+, 2014)

Show affiliations

- [Marino, A. F.](#)
- [Milone, A. P.](#)
- [Yong, D.](#)
- [Dotter, A.;](#)
- [Da Costa, G.](#)
- [Asplund, M.;](#)
- [Jerjen, H.](#)
- [Mackey, D.;](#)
- [Norris, J.;](#)
- [Cassisi, S.;](#)
- [Sbordone, L.;](#)
- [Stetson, P. B.;](#)
- [Weiss, A.;](#)
- [Aparicio, A.;](#)
- [Bedin, L. R.;](#)
- [Lind, K.](#)
- [Monelli, M.;](#)
- [Piotto, G.;](#)
- [Angeloni, R.;](#)
- [Buonanno, R.](#)

Abstract

In this paper, we used four distinct photometric data sets. First, we used Stetson (2000PASP..112..925S) ground-based B, V, R and I photometry. This photometric catalogue has been established from about 550 images taken at different telescopes, i.e. the Max Planck 2.2m, the Cerro Tololo Inter-American Observatory (CTIO) 4, 1.5 and 0.9m telescopes, and the Dutch 0.9m telescope in La Silla. In the present work, we have complemented the Stetson catalogue with images collected with the Wide Field Imager (WFI) of the Max Planck 2.2m telescope at La Silla (WFI@2.2m) through the U filter under the SURvey of Multiple pOpulations (SUMO) campaign.

Secondly, to study stars in the halo of NGC 1851, we collected BVI images with WFI@2.2m of a field between ~ 10 and 35 arcmin to the south of the cluster centre. Photometry and astrometry for this data set have been obtained by using the program `img2xym_WFI` and the procedure described by Anderson et al. (2006A&A...454.1029A).

Third and finally, to investigate the most crowded central regions, we use Hubble Space Telescope (HST) F606W and F814W photometry obtained with the Wide Field Channel of the Advanced Camera for Survey (WFC/ACS) and F275W photometry

collected with the Ultraviolet and Visual Channel of the Wide Field Camera 3 (UVIS/WFC3).
(2 data files).

Publication:

VizieR On-line Data Catalog: J/MNRAS/442/3044. Originally published in:
2014MNRAS.442.3044M

Pub Date:

March 2015

Bibcode:

[2015yCat..74423044M](#)

Keywords:

- Clusters: globular
- Photometry
- Radial velocities
- Abundance

Comments:

table2.dat 191x106 Coordinates, basic photometric data and radial;
velocities for the NGC 1851-halo and ; cluster (inner field) stars ; table3.dat
104x96 Adopted atmospheric parameters and chemical; abundances for the
NGC 1851-halo and ; cluster (central field) stars