



Publication Year	2019
Acceptance in OA	2021-04-21T14:13:19Z
Title	The surface composition of asteroid 162173 Ryugu from Hayabusa2 near-infrared spectroscopy
Authors	Kitazato, K., Milliken, R. E., Iwata, T., Abe, M., Ohtake, M., Matsuura, S., Arai, T., Nakauchi, Y., Nakamura, T., Matsuoka, M., Senshu, H., Hirata, N., Hiroi, T., Pilorget, C., Brunetto, R., Poulet, F., Riu, L., Bibring, J. -P., Takir, D., Domingue, D. L., Vilas, F., Barucci, M. A., PERNA, Davide, PALOMBA, Ernesto, GALIANO, ANNA, Tsumura, K., Osawa, T., Komatsu, M., Nakato, A., Takato, N., Matsunaga, T., Takagi, Y., Matsumoto, K., Kouyama, T., Yokota, Y., Tatsumi, E., Sakatani, N., Yamamoto, Y., Okada, T., Sugita, S., Honda, R., Morota, T., Kameda, S., Sawada, H., Honda, C., Yamada, M., Suzuki, H., Yoshioka, K., Hayakawa, M., Ogawa, K., Cho, Y., Shirai, K., Shimaki, Y., Yamaguchi, A., Ogawa, N., Terui, F., Yamaguchi, T., Takei, Y., Saiki, T., Nakazawa, S., Tanaka, S., Yoshikawa, M., Watanabe, S., Tsuda, Y.
Publisher's version (DOI)	10.1126/science.aav7432
Handle	http://hdl.handle.net/20.500.12386/30832
Journal	SCIENCE
Volume	364

Title: The surface composition of asteroid 162173 Ryugu from Hayabusa2 near-infrared spectroscopy

Authors: K. Kitazato^{1*}, R. E. Milliken², T. Iwata^{3,4}, M. Abe^{3,4}, M. Ohtake^{3,4}, S. Matsuura⁵,
T. Arai⁶, Y. Nakauchi³, T. Nakamura⁷, M. Matsuoka³, H. Senshu⁸, N. Hirata¹, T. Hiroi²,
C. Pilorget⁹, R. Brunetto⁹, F. Poulet⁹, L. Riu³, J.-P. Bibring⁹, D. Takir¹⁰, D. L. Domingue¹¹,
F. Vilas¹¹, M. A. Barucci¹², D. Perna^{13,12}, E. Palomba¹⁴, A. Galiano¹⁴, K. Tsumura^{7,15},
T. Osawa¹⁶, M. Komatsu⁴, A. Nakato³, T. Arai⁸, N. Takato^{17,4}, T. Matsunaga¹⁸, Y. Takagi¹⁹,
K. Matsumoto^{17,4}, T. Kouyama²⁰, Y. Yokota^{3,21}, E. Tatsumi²², N. Sakatani³, Y. Yamamoto^{3,4},
T. Okada^{3,22}, S. Sugita²², R. Honda²¹, T. Morota²³, S. Kameda²⁴, H. Sawada³, C. Honda¹,
M. Yamada⁸, H. Suzuki²⁵, K. Yoshioka²², M. Hayakawa³, K. Ogawa²⁶, Y. Cho²², K. Shirai³,
Y. Shimaki³, N. Hirata²⁶, A. Yamaguchi^{27,4}, N. Ogawa³, F. Terui³, T. Yamaguchi²⁸, Y. Takei³,
T. Saiki³, S. Nakazawa³, S. Tanaka^{3,4}, M. Yoshikawa^{3,4}, S. Watanabe^{23,3}, Y. Tsuda^{3,4}

Affiliations:

¹The University of Aizu, Fukushima, Japan.

²Brown University, Providence, RI, USA.

³Institute of Space and Astronautical Science (ISAS), Japan Aerospace Exploration Agency (JAXA), Sagamihara, Japan.

⁴The Graduate University for Advanced Studies (SOKENDAI), Kanagawa, Japan.

⁵Kwansei Gakuin University, Hyogo, Japan.

⁶Ashikaga University, Tochigi, Japan.

⁷Tohoku University, Sendai, Japan.

⁸Chiba Institute of Technology, Chiba, Japan.

⁹Institut d'Astrophysique Spatial, Université Paris-Sud, Orsay, France.

¹⁰Jacobs, Astromaterials Research and Exploration Science, NASA Johnson Space Center, Houston, TX, USA.

¹¹Planetary Science Institute, Tucson, AZ, USA.

¹²Laboratoire d'Etudes Spatiales et d'Instrumentation en Astrophysique (LESIA), Observatoire de Paris, Meudon, France.

¹³Osservatorio Astronomico di Roma, Istituto Nazionale di Astrofisica (INAF), Monte Porzio Catone, Italy.

¹⁴Istituto di Astrofisica e Planetologia Spaziali, INAF, Roma, Italy.

¹⁵Tokyo City University, Tokyo, Japan.

¹⁶Japan Atomic Energy Agency, Ibaraki, Japan.

¹⁷National Astronomical Observatory of Japan, Tokyo, Japan.

¹⁸National Institute for Environmental Studies, Ibaraki, Japan.

¹⁹Aichi Toho University, Nagoya, Japan.

²⁰National Institute of Advanced Industrial Science and Technology, Tokyo, Japan.

²¹Kochi University, Kochi, Japan.

²²The University of Tokyo, Tokyo, Japan.

²³Nagoya University, Nagoya, Japan.

5 ²⁴Rikkyo University, Tokyo, Japan.

²⁵Meiji University, Tokyo, Japan.

²⁶Kobe University, Kobe, Japan.

²⁷National Institute of Polar Research, Tokyo, Japan.

²⁸Mitsubishi Electric Corporation, Kanagawa, Japan.

10 *Correspondence to: kitazato@u-aizu.ac.jp

Abstract: The near-Earth asteroid 162173 Ryugu, the target of Hayabusa2 sample return mission, is believed to be a primitive carbonaceous object. The Near Infrared Spectrometer (NIRS3) on Hayabusa2 acquired reflectance spectra of Ryugu's surface to provide direct measurements of the surface composition and geological context for the returned samples. A weak, narrow absorption feature centered at 2.72 μm was detected across the entire observed surface, indicating that hydroxyl (OH)-bearing minerals are ubiquitous there. The intensity of the OH feature and low albedo are similar to thermally- and/or shock-metamorphosed carbonaceous chondrite meteorites. There are few variations in the OH-band position, consistent with Ryugu being a compositionally homogeneous rubble-pile object generated from impact fragments of an undifferentiated aqueously altered parent body.

One Sentence Summary: Ryugu possesses a very dark, hydroxyl-bearing surface, spectrally similar to thermally- and/or shock-metamorphosed carbonaceous meteorites.

Main Text: Scientific knowledge of extraterrestrial materials collected by sample-return missions is enhanced if they can be analyzed in the broader context of their parent body. In June 2018, the Japanese Aerospace Exploration Agency (JAXA) spacecraft and sample return mission Hayabusa2 (1) arrived at the near-Earth target asteroid 162173 Ryugu. A mapping campaign with the onboard remote sensing instruments was performed to provide context for sample collection. Prior to arrival, ground- and space-based telescopic measurements indicated the surface of Ryugu was consistent with that of a low-albedo C-type (assumed to be carbon-bearing) asteroid (2, 3). Carbonaceous chondrite meteorites that exhibit evidence of chemical alteration by water, such as CM and CI type, have been proposed as possible compositional analogues for Ryugu (4-7). However, the lack of spectral data in the 3 μm region, where OH stretching and H₂O bending vibrational modes occur, has precluded definitive compositional identification. We report results of near-infrared (NIR) reflectance observations of Ryugu by the Near Infrared Spectrometer (NIRS3) on the Hayabusa2 spacecraft.

NIRS3 is a point spectrometer with a 0.1° field of view that takes continuous point-target spectra over the effective wavelength range 1.8-3.2 μm (8). On June 21 2018, NIRS3 began observing Ryugu from a distance of 70 km. On July 11 and 19, NIRS3 operated in a scanning mode, in which slews of the spacecraft were combined with the rotational motion of the asteroid, to acquire near-global coverage at a surface spatial resolution of 40 and 20 m, respectively. During a descent close to the surface on August 6-7, NIRS3 continuously acquired spectra down

to 1 km altitude, corresponding to a spatial resolution of 2 m. By the time of sampling site selection on 17 August, NIRS3 had acquired over 69,000 spectra of Ryugu's surface. These data can be used to constrain surface composition at sampling locations and across the asteroid as a whole (9).

All reflectance spectra of Ryugu exhibit thermally emitted radiation at wavelengths longer than 2.1-2.4 μm (Fig. 1A). Daytime surface temperatures of Ryugu were estimated from the intensity of the thermal component; they range between 330 to 370 K when the asteroid was located at a heliocentric distance of 1.01 astronomical units (au) (Fig. 2C-D). The temperature estimated from each spectrum was used to model and subtract the contribution due to thermal emission from the total observed radiance, thus isolating the surface reflected sunlight (9). After conversion from radiance to reflectance, the thermally-corrected spectra of Ryugu exhibit several common features. There is a very low reflectance factor (simply "reflectance" hereafter) value across nearly the entire body. The reflectance has been corrected to a standard viewing geometry (incidence angle 30° , emission angle 0° , phase angle 30°) using the radiative transfer theory of Hapke (9, 10). The globally-averaged reflectance at 2.0 μm is 0.017 ± 0.002 , consistent with values at visible wavelengths observed by the Hayabusa2 Optical Navigation Camera Telescope (ONC-T) (11) (Fig. S3). Reflectance values vary within 15% across the entire observed surface, excluding regions in shadow (Fig. 2A-B). Brighter surfaces are primarily observed along the equatorial ridge, crater rims and for individual boulders, again similar to visible wavelength images. The spectra commonly exhibit a weak positive spectral slope (0.2 to 0.6 % μm^{-1}) between 2.0 and 2.5 μm .

All spectra of Ryugu exhibit a weak, narrow absorption feature centered at 2.72 μm (Fig. 1B) and with absorption intensity values that range from 7 to 10% of the spectral continuum. The 2.72 μm feature is observable even in the radiance data of Ryugu, and its veracity was confirmed by searching for a similar feature in NIRS3 calibration observations of the Moon, where it was found to be lacking (9). The absolute band depth of this feature in processed reflectance data is dependent on the thermal correction. The intensity of the 2.72 μm feature exhibits a positive correlation with estimated surface temperatures (Fig. S5), which indicates that uncertainties in the radiometric calibration and/or thermal emission component could be responsible for the observed variations in the band depth of this feature. When normalized by the observed temperature trend, the intensity of the 2.72 μm feature exhibits no clear correlation with topographic/morphologic features (Fig. 2E-F).

Ryugu is darker than other objects visited by spacecraft, including the nucleus of comet 67P/Churyumov-Gerasimenko (11-13). It is also darker than many laboratory reflectance spectra of carbonaceous chondrite meteorites, which are believed to be derived from C-type or similar asteroids (14). Meteorite samples have reflectance values as dark as 0.03 to 0.10 at 2.0 μm wavelength (15-17), in part because they contain Fe-bearing phyllosilicates, opaque minerals (magnetite and Fe sulfides), and organic carbon, yet the surface of Ryugu is darker still. Such a low and relatively homogeneous albedo indicates the presence of a highly absorbing component dispersed throughout the surface materials of the asteroid, though we cannot uniquely determine its composition and origin. Possible explanations include high abundances of carbon, opaque minerals (e.g., magnetite), and/or products of shock-induced metamorphism (e.g., dark glassy components).

The detection of an absorption feature at 2.72 μm indicates the presence of OH attached to a cation, and the position of the reflectance minimum indicates is most likely Mg. The band

position is similar to Mg-OH features observed in Mg-rich phyllosilicates, such as serpentine and saponite, which are known to be present in aqueously altered CI and CM chondrites (18, 19). It is also similar to OH features observed in spectra of the dwarf planet Ceres and some C-complex main belt asteroids, which have been interpreted as evidence for Mg-rich phyllosilicates (20, 21). Our spectra indicate that the OH band position does not vary across the surface of Ryugu within the ~18 nm spectral sampling of the instrument (Fig. S7), suggesting a relatively homogeneous phyllosilicate cation composition.

No other absorption features are detected in the 3 μm region in addition to the 2.72 μm feature. Spectra of some C-complex main belt asteroids have been found to exhibit absorption features that may be attributable to H₂O frost (22, 23), H₂O-bearing minerals (24) or NH₄-bearing phyllosilicates (20, 21) in the NIRS3 wavelength range. A decrease in reflectance values at 3 μm can be produced in the NIRS3 data by overestimating the surface temperatures during thermal correction, and it is also possible to mask a weak 3 μm feature if the thermal contribution is underestimated. Nevertheless, a 3 μm feature is expected to be detectable in the radiance data if it were similar to that found on other asteroids.

There are no published meteorite samples whose reflectance spectra perfectly match that of Ryugu at visible to near-infrared wavelengths. However, spectra of thermally-metamorphosed CI chondrites and shocked CM chondrites are similar to Ryugu at NIR wavelengths in brightness and shape. Laboratory spectra of samples taken from the Ivuna meteorite (classified as CI1) heated to 500°C and from MET 01072 (classified as shocked CM2) are darker than unaltered carbonaceous chondrites, flat, and yet retain a weak 2.72 μm feature (Fig. 3). These meteorite data suggest that thermal alteration processes, such as partial dehydration and decomposition of hydrated minerals induced by static or shock heating, can act to darken hydrated carbonaceous chondrites. Such processes are consistent with current interpretations of Ryugu's formation history. The low bulk density (~1.2 g cm⁻³) of Ryugu suggests that it is a rubble-pile asteroid formed by a collisional event with the parent body (25), thus it is likely to have experienced shock and post-shock heating that could lead to darkening, dehydration, and dehydroxylation. However, it is also possible that the weak OH absorption is because the degree of aqueous alteration on Ryugu was never extensive to begin with, perhaps due to low water-to-rock ratios or slow/incomplete hydration reactions on the parent body.

Alternatively, it has been suggested that Ryugu's orbit might have had shorter perihelion distances in the past, a characteristic that would have increased radiative heating from the Sun (26) and altered the mineralogy of the uppermost surface. Similarly, the surface of Ryugu has experienced solar-wind irradiation and micrometeorite impacts (space weathering), which can alter surface composition and spectral properties. Laboratory experiments suggest space weathering can destroy OH bonds, causing the disappearance or reduction of a 2.72 μm feature (27), a process that may explain the uniformly weak OH signature observed in NIRS3 data. These processes – solar radiative heating and space weathering – represent near-surface phenomena that continue to operate at Ryugu today, whereas the other interpretations for the apparent low hydration state represent inherent chemical and mineralogical attributes of the asteroid as a result of its early geological history.

Reflectance spectra of Ryugu differ from the heated Ivuna and MET 01072 meteorite samples at visible wavelengths (Fig. 3). This may be due to effects of space weathering or physical properties that could preferentially affect shorter wavelengths. Laboratory experiments show that reflectance spectra of some ion irradiated CI and CM chondrites tend to increase in

brightness at visible wavelengths (27). Moreover, there are differences in particle size and likely porosity between the meteorite samples and Ryugu's surface materials. For CM chondrite-like materials, an increase in the average particle size of a powdered sample results in spectra with more positive slope and lower reflectance values, as well as an increase in band depth values of diagnostic absorption bands (28). With increasing porosity, overall reflectance decreases (28). Despite these differences, hydrated carbonaceous chondrites that have been metamorphosed by thermal and/or shock processes are the best available spectral matches to the NIRS3 data, so Ryugu may be analogous in composition.

Another possible spectral analogue is interplanetary dust particles (IDPs). IDPs are known to be darker than CV meteorite samples and have featureless reflectance spectra between 0.4 and 1.0 μm (29). However, no published reflectance spectra of IDPs span the NIRS3 wavelength range and therefore a direct comparison cannot be made. In this scenario, and if the spectral properties observed by NIRS3 are indicative of the bulk asteroid, then Ryugu may consist of carbon-rich, partially-hydrated, possibly comet-like material that is darker than common aqueously altered carbonaceous chondrites. If Ryugu is indeed such a carbon-rich body, then nonlinear masking effects induced by carbonaceous material (30) may result in an underestimation of the relative abundance of phyllosilicates and the degree of aqueous alteration based on the apparent weak 2.72 μm feature.

References and Notes:

1. Y. Tsuda, M. Yoshikawa, M. Abe, H. Minamino, S. Nakazawa, System design of the Hayabusa 2 – Asteroid sample return mission to 1999 JU3. *Acta Astronaut.* **91**, 356-362, (2013).
2. R. P. Binzel, A. W. Harris, S. J. Bus, T. H. Burbine, Spectral properties of near-Earth objects: Palomar and IRTF results for 48 objects including spacecraft targets (9969) Braille and (10302) 1989 ML, *Icarus* **151**, 139-149, (2001).
3. H. Campins *et al.*, Spitzer observations of spacecraft target 162173 (1999 JU3). *Astron. Astrophys.* **503**, L17-L20 (2009).
4. F. Vilas, Spectral characteristics of Hayabusa 2 near-Earth asteroid targets 162173 1999 JU3 and 2001 QC34. *Astron. J.* **135**, 1101-1105 (2008).
5. N. A. Moskovitz *et al.*, Rotational characterization of Hayabusa II target asteroid (162173) 1999 JU3. *Icarus* **224**, 24-31 (2013).
6. D. Perna *et al.*, Spectral and rotational properties of near-Earth asteroid (162173) Ryugu, target of the Hayabusa2 sample return mission. *Astron. Astrophys.* **599**, L1 (2017).
7. L. Le Corre *et al.*, Ground-based characterization of Hayabusa2 mission target asteroid 162173 Ryugu: constraining mineralogical composition in preparation for spacecraft operations. *Mon. Not. R. Astron. Soc.* **475**, 614-623 (2018).
8. T. Iwata *et al.*, NIRS3: the near-infrared spectrometer on Hayabusa2. *Space Sci. Rev.* **208**, 317-337 (2017).
9. Materials and methods are available as supplementary materials.

10. B. Hapke, Theory of Reflectance and Emittance Spectroscopy. (Cambridge University Press, 1993).
11. S. Sugita *et al.*, The geomorphology, color, and thermal properties of Ryugu: Implications for parent-body processes. *Science* (this issue).
- 5 12. S. Fornasier *et al.*, Spectrophotometric properties of the nucleus of comet 67P/Churyumov-Gerasimenko from the OSIRIS instrument onboard the ROSETTA spacecraft. *Astron. Astrophys.* **583**, A30 (2015).
13. M. Ciarniello *et al.*, Photometric properties of comet 67P/Churyumov-Gerasimenko from VIRTIS-M onboard Rosetta. *Astron. Astrophys.* **583**, A31 (2015).
- 10 14. T. H. Burbine, T. J. McCoy, A. Meibom, B. Gladman, K. Keil, “Meteoritic parent bodies: their number and identification”, in Asteroids III, W. F. Bottke Jr., A. Cellino, P. Paolicchi, R. P. Binzel, Eds. (Univ. of Arizona Press, Tucson, AZ, 2002), pp. 653-667.
15. E. A. Cloutis, T. Hiroi, M. J. Gaffey, C. M. O’D. Alexander, P. Mann, Spectral reflectance properties of carbonaceous chondrites: 1. CI chondrites. *Icarus* **212**, 180-209 (2011).
- 15 16. E. A. Cloutis, P. Hudon, T. Hiroi, M. J. Gaffey, P. Mann, Spectral reflectance properties of carbonaceous chondrites: 2. CM chondrites. *Icarus* **216**, 309-346 (2011).
17. E. A. Cloutis, P. Hudon, T. Hiroi, M. J. Gaffey, P. Mann, Spectral reflectance properties of carbonaceous chondrites: 8. “Other” carbonaceous chondrites: CH, ungrouped, polymict, xenolithic inclusions, and R chondrites. *Icarus* **221**, 984-1001 (2012).
- 20 18. P. Beck *et al.*, Hydrous mineralogy of CM and CI chondrites from infrared spectroscopy and their relationship with low albedo asteroids. *GeoChim. Cosmo. Acta* **74**, 4881-4892 (2010).
19. D. Takir *et al.*, Nature and degree of aqueous alteration in CM and CI carbonaceous chondrites. *Meteorit. Planet. Sci.* **48**, 1618-1637 (2013).
- 25 20. M. C. De Sanctis *et al.*, Ammoniated phyllosilicates with a likely outer Solar System origin on (1) Ceres. *Nature* **528**, 241-244 (2015).
21. F. Usui, S. Hasegawa, T. Ootsubo, T. Onaka, AKARI/IRC near-infrared asteroid spectroscopic survey: AcuA-spec. *Publ. Astron. Soc. Japan*, psy125 (2018).
22. H. Campins *et al.*, Water ice and organics on the surface of the asteroid 24 Themis. *Nature* **464**, 1320-1321 (2010).
- 30 23. A. S. Rivkin, J. P. Emery, Detection of ice and organics on an asteroidal surface. *Nature* **464**, 1322-1323 (2010).
24. D. Takir, J. P. Emery, Outer main belt asteroids: identification and distribution of four 3- μ m spectral groups. *Icarus* **219** 641-654 (2012).
- 35 25. S. Watanabe *et al.*, Hayabusa2 arrives at the carbonaceous asteroid 162173 Ryugu – a spinning-top-shaped rubble pile. *Science* (this issue).
26. P. Michel, M. Delbo, Orbital and thermal evolutions of four potential targets for a sample return space mission to a primitive near-Earth asteroid. *Icarus* **209**, 520-534 (2010).
27. C. Lantz *et al.*, Ion irradiation of carbonaceous chondrites: a new view of space weathering on primitive asteroids. *Icarus* **285**, 43-57 (2017).

28. E. A. Cloutis *et al.*, Spectral reflectance “deconstruction” of the Murchison CM2 carbonaceous chondrite and implications for spectroscopic investigations of dark asteroids. *Icarus* **305**, 203-224 (2018).
29. J. P. Bradley, L. P. Keller, D. E. Brownlee, K. L. Thomas, Reflectance spectroscopy of interplanetary dust particles. *Meteorit. Planet. Sci.* **31**, 394-402 (1996).
30. R. N. Clark, Spectral properties of mixtures of montmorillonite and dark carbon grains: implications for remote sensing minerals containing chemically and physically adsorbed water. *J. Geophys. Res.* **88**, 10635-10644 (1983).
31. T. Kouyama *et al.*, Development of an application scheme for the SELENE/SP lunar reflectance model for radiometric calibration of hyperspectral and multispectral sensors. *Planet. Space Sci.* **124**, 76-83 (2016).
32. M. Ishiguro *et al.*, Optical properties of (162173) 1999 JU3: in preparation for the JAXA Hayabusa 2 sample return mission. *Astrophys. J.* **792**, 74 (2014).
33. L. G. Henyey, J. L. Greenstein, Diffuse radiation in the Galaxy. *Astrophys. J.* **93**, 70-83 (1941).
34. C. H. Acton, Ancillary data services of NASA’s Navigation and Ancillary Information Facility. *Planet. Space Sci.* **44**, 65-70 (1996).
35. R. N. Clark, C. M. Pieters, R. O. Green, J. W. Boardman, N. E. Petro, Thermal removal from near-infrared imaging spectroscopy data of the Moon. *J. Geophys. Res.* **116**, E00G16 (2011).
36. NASA Reflectance Experiment LABORatory (RELAB) database, Brown University (<http://www.planetary.brown.edu/relab/>).
37. T. Hiroi, C. M. Pieters, M. E. Zolensky, M. Prinz, Reflectance spectra (UV-3 μ m) of heated Ivuna (CI) meteorite and newly identified thermally metamorphosed CM chondrites. *27th Lunar Planet. Sci. Conf.* abstr. 551 (1996).
38. Y. Nakauchi *et al.*, Reflectance spectra of carbonaceous chondrites measured under asteroid-like conditions: implications for Hayabusa2’s NIRS3 instrument. *49th Lunar Planet. Sci. Conf.* abstr. 2083 (2018).
39. T. Nakamura *et al.*, Chondrule flattening of MET01072 CM chondrite: possible evidence for long-duration slow impact. *Goldschmidt* abstr. 2228 (2016).

Acknowledgments: We thank the Hayabusa2 engineering teams for their efforts in making the mission successful and enhancing science operations, in particular T. Masuda, S. Yasuda, K. Matsushima, and T. Ohshima. We also thank H. Murao, Y. Sakata, and K. Taguchi for the development of NIRS3. T.N. thanks M. Takahashi for support in the meteorite analysis.

Funding: The NIRS3 instrument was funded and coordinated by JAXA and built by Meisei Electric, Genesis, and Hamamatsu Photonics. K.K. and the Japanese coauthors acknowledge the support from JAXA and the Hayabusa2 project. The French coauthors acknowledge the support from CNES. The Italian coauthors acknowledge the support from ASI. D.T. and D.L.D. have been supported by the NASA Hayabusa2 Participating Scientist Program (NNX17AL02G, NNX16AL34G). M.A., T.N., N.H., T.O., T.M., and S.W. have been supported by Grant-in-Aid for Scientific Research from JSPS (16H04044, 17H06459, 17K05639). Y.N., M.M., E.T., N.S., and S.S. have been supported by the JSPS Core-to-Core Program (17H01175). The work of T.H.

is funded by the NASA's Emerging Worlds Program (NNX17AK46G). The work of D.P. is funded by the European Union's H2020 programme (664931). **Author contributions:** K.K. led the operation and calibration of NIRS3, analyzed the data, and wrote the manuscript. R.E.M. contributed to the calibration and data analysis, and assisted in the writing. T.I. led the development of NIRS3. M.A., M.O., S.M., T.A., Y.N., and K.T. contributed the development and operation. T.N., M.M., and T.H. provided information about laboratory spectra of meteorites. H.S., C.P., R.B., F.P., and L.R. contributed to the data analysis. D.T., D.L.D., M.A.B., D.P., E.P., and A.G. participated in the interpretation of the data. All authors discussed the results and commented on the manuscript. **Competing interests:** The authors declare no competing interests. **Data and materials availability:** The NIRS3 calibrated data are available through the JAXA Data Archives and Transmission System website (https://darts.isas.jaxa.jp/pub/hayabusa2/nirs3_bundle/browse/).

Supplementary Materials:

Materials and Methods

Figures S1-S7

Table S1

References (31-39)




Fig. 1. NIRS3 near-infrared spectra of Ryugu. (A) Reflectance spectra including thermal emission. The range of variation in spectra acquired from the equatorial scan on July 10, 2018 are shown (gray) along with the average spectrum (red). I/F is the measured radiance divided by the solar flux. (B) An example of a thermally-corrected spectrum (observation date and time: 2018-07-10T06:02:22 UTC). Error bars are calculated based on the uncertainties in the element-to-element radiometric calibration (9). Shaded areas indicate regions with large calibration residuals. The absorption band indicated by the arrow at 2.72 μm is due to OH.




Fig. 2. NIRS3 spectral features projected onto the shape model of Ryugu (25). The left- and right-side views correspond to the western and eastern hemispheres, respectively. Yellow-shaded areas represent regions that were not observed. (A and B) Reflectance factor at $2.0\ \mu\text{m}$ corrected to a standard viewing geometry (9). (C and D) Surface temperature (K) derived from the thermal emission component. (E and F) Normalized intensity of the $2.72\ \mu\text{m}$ feature. The Z and X arrows indicate Ryugu's rotation axis and zero-longitude definition, respectively.




Fig. 3. Comparison of Ryugu with meteorite spectra. (A) The globally-averaged NIRS3 spectrum of Ryugu compared with laboratory spectra of meteorite samples: Ivuna (CI1), Cold Bokkeveld (CM2), MET 01072 (shocked CM2), and Allende (CV3). Details and references for the meteorite spectra are listed in Table S1. A ground-based visible and near-infrared spectrum of Ryugu (6) is also plotted. (B) Enlargement of (A) in the NIRS3 wavelength range.