



Publication Year	2021
Acceptance in OA	2022-03-29T13:56:57Z
Title	Seeing-sorted Visible Multi-Object Spectrograph U-band Imaging of the GOODS-south Field*
Authors	Otteson, Lillian, McCabe, Tyler, Ashcraft, Teresa A., Windhorst, Rogier A., Jansen, Rolf A., Cohen, Seth H., NONINO, Mario, PARIS, Diego, GRAZIAN, Andrea, FONTANA, Adriano, GIALLONGO, Emanuele, SPEZIALI, Roberto, TESTA, Vincenzo, Boutsia, Konstantina, Koekemoer, Anton, O'Connell, Robert W., Rutkowski, Michael J., Ryan, Russell E., Scarlata, Claudia, Teplitz, Harry
Publisher's version (DOI)	10.3847/2515-5172/ac1db6
Handle	http://hdl.handle.net/20.500.12386/32049
Journal	RESEARCH NOTES OF THE AAS
Volume	5

OPEN ACCESS

Seeing-sorted Visible Multi-Object Spectrograph *U*-band Imaging of the GOODS-south Field*

Lillian Otteson¹ , Tyler McCabe¹, Teresa A. Ashcraft²,
 Rogier A. Windhorst¹ , Rolf A. Jansen¹ , Seth H. Cohen¹ ,
 Mario Nonino³, Diego Paris⁴, Andrea Grazian⁵, Adriano Fontana⁴ 

+ Show full author list

Published August 2021 • © 2021. The Author(s). Published by the American Astronomical Society.

Research Notes of the AAS, Volume 5, Number 8

Citation Lillian Otteson *et al* 2021 *Res. Notes AAS* **5** 190

lotteson@asu.edu

¹ School of Earth and Space Exploration, Arizona State University, Tempe, AZ 85287-1404, USA; lotteson@asu.edu

² The University of Tampa, Tampa, FL 33606, USA

³ INAF-Osservatorio Astronomico di Trieste, Via Bazzoni 2, I-34124 Trieste, Italy

⁴ INAF—Osservatorio Astronomico di Roma, via Frascati 33, I-00078 Monte Porzio Catone, Italy

⁵ INAF—Osservatorio Astronomico di Padova Vicolo dell'Osservatorio, 5 Padova (PD) I-35122, Italy

⁶ Carnegie Observatories, Las Campanas Observatory, Colina El Pino, Casilla 601, La Serena, Chile

⁷ Space Telescope Science Institute, Baltimore, MD 21218, USA

⁸ Department of Astronomy, University of Virginia, Charlottesville, VA 22904-4325, USA

⁹ Minnesota State University, Mankato, MN 56001, USA

This site uses cookies. By continuing to use this site you agree to our use of

 Downloads

aa36791-...

Acharyya...

staa3796....

aa39738-...

IOEXPOR...

100%

Clear

Street SE, Minneapolis, MN 55455, USA

¹¹ Infrared Processing and Analysis Center, MS 100-22, Caltech, Pasadena, CA 91125, USA

Lillian Otteson  <https://orcid.org/0000-0001-9665-3003>

Rogier A. Windhorst  <https://orcid.org/0000-0001-8156-6281>

Rolf A. Jansen  <https://orcid.org/0000-0003-1268-5230>

Seth H. Cohen  <https://orcid.org/0000-0003-3329-1337>

Adriano Fontana  <https://orcid.org/0000-0003-3820-2823>

Emanuele Giallongo  <https://orcid.org/0000-0003-0734-1273>

Vincenzo Testa  <https://orcid.org/0000-0003-1033-1340>

Konstantina Boutsia  <https://orcid.org/0000-0003-4432-5037>

Anton Koekemoer  <https://orcid.org/0000-0002-6610-2048>

Michael J. Rutkowski  <https://orcid.org/0000-0001-7016-5220>

Russell E. Ryan  <https://orcid.org/0000-0003-0894-1588>

Claudia Scarlata  <https://orcid.org/0000-0002-9136-8876>

Harry Teplitz  <https://orcid.org/0000-0002-7064-5424>

Received August 2021

Revised August 2021

Accepted August 2021

Published August 2021

<https://doi.org/10.3847/2515-5172/ac1db6>

Ultraviolet astronomy; Astronomical seeing; Astronomy data analysis

Create or edit your corridor alerts

What are corridors? 

Create citation alert

Abstract

We present the optimal resolution and optimal depth *U*-filter mosaics using the seeing-sorted method of Ashcraft et al. on deep, ground-based *U*-band imaging of the Great Observatories Origins Deep Survey South field as part of the near-UV imaging program UVCANDELS. We use the

This site uses cookies. By continuing to use this site you agree to our use of  [U-band images obtained with the Visible Multi-Object Spectrograph on the](#)

European Southern Observatory Very Large Telescope by Nonino et al. Our best resolution mosaic includes images with a seeing full-width half maximum (FWHM) $\leq 0''.8$, and encompasses 50% of the data. Our best depth mosaic includes images with FWHM $\leq 1''.5$, corresponding to 100% of the data. Prior to being combined, the source fluxes in each individual background-subtracted image are corrected to match a 3D-HST photometric catalog of the same field to correct variations in the *U*-band zero-points. These mosaics provide deep *U*-band data complementary to the UVCANDELS HST WFC3 F275W and ACS F435W images. We assess the depth of both *U*-band mosaics.

Export citation and abstract

[BibTeX](#)

[RIS](#)

◀ **Previous** article in issue

Next article in issue ▶



Original content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

1. Introduction

Nonino et al. (2009) presented the results of deep *U*-band and *R*-band imaging of the GOODS-south field. In the current paper, we present the seeing-sorted version of their *U*-band images, which also have a closer zero-point match to other GOODS multi-wavelength photometry. By utilizing a number of image processing techniques, Nonino et al. (2009) were able to create an image mosaic out of the 552 single chip images taken from the VIMOS instrument on the Very Large Telescope (VLT) in Chile. The final mosaic from this work had a full width half maximum of $0''.8$ and reached a depth of $m_{AB} \approx 29.8$ mag (Nonino et al. 2009).

Ashcraft et al. (2018) used observations from the Large Binocular Telescope

This site uses cookies. By continuing to use this site you agree to our use of

Downloads
 aa36791-...
 Acharyya...
 staa3796....
 aa39738-...
 IOEXPOR...
 100%
 Clear

image mosaics: (1) the best resolution mosaic, generated with images using seeing $\text{FWHM} \leq 0''.8$, and (2) the best depth mosaic, cut off at $1''.8$ seeing (Ashcraft et al. 2018). This stacking method is useful to mitigate atmospheric effects associated with ground-based imaging. The optimal resolution mosaic is best for studying bright galaxies, and shows structure in galaxies more clearly than the best depth mosaic. The optimal depth stack, however, is more sensitive to lower surface brightness objects and structures. Ashcraft et al. (2018) showed using galaxy number counts that the optimal depth mosaic detects more of the faintest galaxies.

The work presented here focuses on the GOODS-south field, builds on the data of Nonino et al. (2009) and uses the seeing-sorted stacking method of Ashcraft et al. (2018). An additional step was incorporated to reduce uncertainty and variability in the zero-points of individual images following a process similar to one described in T. McCabe et al. (2021, in preparation).

2. Observations

The 552 individual U_V -filter images used in this work were taken between 2004 August and 2006 October by the VISIBLE Multi-Object Spectrograph (VIMOS) on ESO's VLT (Nonino et al. 2009). VIMOS is a four-chip charge coupled device (CCD) camera with a pixel scale of $0''.205/\text{pixel}$ and a field of view of $4' \times 7' \times 8'$. In imaging mode, each chip uses a 2048×2440 pixels EEV 44–82 backside illuminated CCD. The chips are separated by $2'$ gaps. Special care was taken when dithering to get uniform coverage of the field (Nonino et al. 2009).

3. Analysis

Ashcraft et al. (2018) found a ~ 0.2 mag difference in the U -band photometric zero-points between their optimal resolution and optimal depth LBT/LBC mosaics. This difference was attributed to variations in transparency among different exposures and across different nights. Here, we therefore correct the flux scale of each image to match the U -band photometry in the 3D-HST photometric catalog of Skelton et al. (2014). Figures 1(e) and (f) show an example of the uncorrected and corrected flux ratios between the VIMOS images and the 3D-HST catalog for the first four nights of observations.

This site uses cookies. By continuing to use this site you agree to our use of

 Downloads
 aa36791-...
 Acharyya...
 staa3796....
 aa39738-...
 IOEXPOR...
 100%
 Clear

0.6%. After applying these *U*-band zero-point corrections, this ratio increased to $99.8\% \pm 0.1\%$, better matching the 3D-HST catalog.

This site uses cookies. By continuing to use this site you agree to our use of

 Downloads

aa36791-...

Acharyya...

staa3796....

aa39738-...

IOEXPOR...

100%

Clear






Figure 1. VLT/VIMOS U_V -filter exposure maps for the best resolution mosaic (a) and the best depth mosaic (b) in GOODS-south. The color indicates the total exposure time in hours. Figure (c) is a histogram of the seeing for the 552 single chip VIMOS images. The median seeing, FWHM $\sim 0.78''$, is indicated by the dashed red line. Exposures with FWHM $\leq 0.8''$ are included in the best resolution mosaic. All exposures are used in the best depth mosaic. Figure (d)

This site uses cookies. By continuing to use this site you agree to our use of cookies. [Learn more](#)

correspond to the optimal resolution mosaic counts and the crosses correspond to the optimal depth mosaic. The counts are color-coded based on the exposure time areas they were taken from (i.e., shallow corresponds to least exposure time, deep corresponds to most exposure time; please magnify the PDF as needed). Figure (e) and (f) show uncorrected and corrected flux ratio distributions for the first four nights of observations, respectively. Nights are separated by solid vertical lines. The horizontal, dashed black line represents flux equal to 3D-HST (Skelton et al. 2014). The solid red line with confidence bands (shaded red) represents the global average for all 139 exposures before and after U_V zero-point corrections to match the 3D-HST U catalog.

All 552 single chip exposures were sorted based on their seeing (Figure 1(c)). SWARP (Bertin et al. 2002; Bertin 2010) was used to combine the individual images. Our best resolution mosaic, cut off at $\text{FWHM} \lesssim 0''.8$, used 288 images, just over 50% of the VIMOS U_V data (Figure 1(a)). Our best depth mosaic used images with seeing $\text{FWHM} \lesssim 1''.5$, corresponding to 100% of the VIMOS U_V data (Figure 1(b)). SExtractor (Bertin & Arnouts 1996) was used to create source catalogs of the mosaics. The mosaics shared the same SExtractor configuration parameters, which are similar to those in Ashcraft et al. (2018). The galaxies were separated from stars as described in, e.g., Windhorst et al. (2011). Figure 1(d) shows the resulting galaxy number counts from our U -band catalogs. The best resolution mosaic U -band counts start to turn over around $m_{\text{AB}} \approx 26.8$ mag, and fall off more quickly than the best depth U -band mosaic, which turns over at $m_{\text{AB}} \approx 27$ mag in U -band. For examples of the differences in resolution and depth between our best resolution and best depth mosaics we refer the reader to Figures 2–5 of Ashcraft et al. (2018), which are also representative for our data.

4. Summary

We present the results of deep U -band imaging of the GOODS-south field

This site uses cookies. By continuing to use this site you agree to our use of cookies. [View our cookie policy](#)

(2009). Two image mosaics were made following the seeing-sorted stacking method of Ashcraft et al. (2018). All 552 single chip images were sorted based on FWHM. The optimal resolution mosaic was assembled using only images with $\text{FWHM} \leq 0\farcs8$. The optimal depth mosaic was made from images with $\text{FWHM} \leq 1\farcs5$. Before creating the mosaics, 139 four-chip exposures were corrected to better match the 3D-HST *U*-bandflux (Skelton et al. 2014). With the completion of these mosaics, three of the four UVCANDELS fields will have ground-based images complementary to HST data. New LBT *U*-banddata for the EGS and COSMOS fields are currently being reduced, thereby completing the seeing-sorted stacking analysis for the UVCANDELS program. In the coming years, these mosaics will complement JWST 1–5 μm observations and help us better understand galaxy assembly over the past 10 billion years. The image mosaics and source catalogs from this work will be released as part of the UVCANDELS program later this year, found at <http://uvcandels.ipac.caltech.edu>.

This work used data from the 3D-HST Treasury Program (HST-GO-12177 and 12328). We also acknowledge support from UVCANDELS grant HST-GO-15647 provided by NASA through the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. We acknowledge support from NASA JWST Interdisciplinary Scientist grants NAG5-12460, NNX14AN10G and 80NSSC18K0200 from GSFC. M.N. acknowledges INAF 1.05.01.86.20.

Footnotes

- * Based on data acquired using the Very Large Telescope (VLT) of the European Southern Observatory (ESO).