



Publication Year	2015
Acceptance in OA	2020-04-06T07:36:37Z
Title	Photometric correction for VIRTIS-M data of comet 67P/CG
Authors	CIARNIELLO, Mauro, CAPACCIONI, FABRIZIO, FILACCHIONE, GIANRICO, Erard, Stéphan, Leyrat, Cedric, Bockelée-Morvan, Dominique, RAPONI, Andrea, DE SANCTIS, MARIA CRISTINA, TOSI, Federico, CAPRIA, MARIA TERESA, PICCIONI, GIUSEPPE, PALOMBA, Ernesto, LONGOBARDO, ANDREA, Drossart, Pierre, Quirico, Eric, Beck, Pierre, Schmitt, Bernard
Handle	http://hdl.handle.net/20.500.12386/23858
Journal	GEOPHYSICAL RESEARCH ABSTRACTS
Volume	17



Photometric correction for VIRTIS-M data of comet 67P/CG

Mauro Ciarniello (1), Fabrizio Capaccioni (1), Gianrico Filacchione (1), Stéphan Erard (2), Cedric Leyrat (2), Dominique Bockelée-Morvan (2), Andrea Raponi (1), Maria Cristina De Sanctis (1), Federico Tosi (1), Maria Teresa Capria (1), Giuseppe Piccioni (1), Ernesto Palomba (1), Andrea Longobardo (1), Pierre Drossart (2), Eric Quirico (3), Pierre Beck (3), and Bernard Schmitt (3)

(1) INAF, IAPS, Rome, Italy (mauro.ciarniello@iaps.inaf.it), (2) LESIA, Observatoire de Paris/CNRS/UPMC/Université Paris-Diderot, Meudon, France, (3) Univ. Grenoble Alpes, CNRS, IPAG, Grenoble, France

Photometric correction for VIRTIS-M data of comet 67P/CG

M. Ciarniello¹ (mauro.ciarniello@iaps.inaf.it), F. Capaccioni¹, G. Filacchione¹, S. Erard², C. Leyrat², D. Bockelée-Morvan², A. Raponi¹, M. C. De Sanctis¹, F. Tosi¹, M. T. Capria¹, G. Piccioni¹, E. Palomba¹, A. Longobardo¹, P. Drossart², E. Quirico³, P. Beck³, B. Schmitt³ and the Rosetta-VIRTIS team. ¹INAF-IAPS, Via del Fosso del Cavaliere 100, I-00133 Rome, Italy, mauro.ciarniello@iaps.inaf.it. ²LESIA, Observatoire de Paris/CNRS/UPMC/Université Paris-Diderot, Meudon, France. ³Univ. Grenoble Alpes, CNRS, IPAG, Grenoble, France.

VIRTIS, the Visible Infrared Thermal Imaging Spectrometer onboard the Rosetta orbiter [1], has acquired so far millions of spectra of the comet 67P/Churyumov-Gerasimenko [2]. The instrument is composed of two subsystems: a high-resolution channel (VIRTIS-H) which is a punctual spectrometer (2.0-5.0 μm) and the mapper (VIRTIS-M) able to produce hyper-spectral images of the target (0.25-5.1 μm). The huge amount of data produced by VIRTIS has been acquired under different observation and illumination conditions. This induces photometric effects on the measured signal that need to be quantified and removed, in order to characterize the intrinsic spectral variability of the surface. To achieve this task we computed a photometric correction from VIRTIS-M data (Ciarniello et al, 2015), starting from August 2014, when the nucleus was largely resolved (MTP006-MT007 observation sequences) by means of a simplified Hapke model [3]. The global surface single particle phase function (SPPF) and the single scattering albedo (SSA) are determined as well as the effect of sub-pixel roughness is discussed. Comparisons with photometric properties of other comets are shown.

This work is supported by the Italian Space Agency (ASI). We acknowledge funding from French and German space agency.

References

- 1- Coradini et al, SSR, 2007
- 2- Capaccioni et al., Science, in Press, 2015
- 3- Hapke, Theory of reflectance and emittance spectroscopy. Cambridge University Press, 2012