



Publication Year	2018
Acceptance in OA	2020-10-20T12:15:51Z
Title	(1) Ceres: Study of Thermal Convection in the Mantle and its Mechanical Effects
Authors	FORMISANO, Michelangelo, Federico, Costanzo, Magni, Gianfranco, DE SANCTIS, MARIA CRISTINA
Handle	http://hdl.handle.net/20.500.12386/27897

Space Studies of the Earth-Moon System, Planets, and Small Bodies of the Solar System (B)
The Golden Age of Small Bodies, Science and Exploration (B1.1)

(1) CERES: STUDY OF THERMAL CONVECTION IN THE MANTLE AND ITS MECHANICAL EFFECTS

Michelangelo Formisano, michelangelo.formisano@iaps.inaf.it

INAF - IAPS Rome, Rome, Italy

Costanzo Federico, costanzo.federico44@gmail.com

INAF-IAPS, Rome, Italy

Maria Cristina De Sanctis, mariacristina.desanctis@iaps.inaf.it

IAPS/INAF, Rome, Italy

Gianfranco Magni, gianfranco.magni@iaps.inaf.it

INAF, Rome, Italy

Ceres is the largest body of the Main Belt, which is characterized by a huge abundance of water ice in its interior. This feature is suggested by its relatively low bulk density (2162 kg m⁻³, Russell et al. 2016, Park et al. 2016) and by several geological and geochemical evidences (specific minerals or salts produced by aqueous alteration, icy patches on the surface, lobate morphologies interpretable as surface flows (De Sanctis et al. 2016, Carrozzo et al. 2018, Raponi et al. 2018, Zolotov 2017 and Schmidt et al., 2017)). Ceres is partially differentiated as suggested by its normalized moment of inertia, 0.37 (Park et al. 2016). A typical internal structure proposed for Ceres is: a rocky core (300-350 km), an icy (or muddy) mantle (100-150 km) and a rocky crust some kilometers in depth (eg. McCord Sotin 2005, Neveu Desch, 2015). The temperature gradient across the mantle, estimated through numerical modelling (e.g. McCord Sotin 2005, Neveu Desch 2015) would be large enough to initiate a thermal convection in the mantle. Since the mantle is not uniquely defined from a composition point of view, in this work we explore how the composition and, in particular the "degree" of muddiness of the mantle, can influence the characteristic of thermal convection. We also estimate the thickness of the top conductive boundary layer and the mechanical stress, which can cause its deformation.

- De Sanctis, M., et al. (2015) [doi:10.1038/nature16172](https://doi.org/10.1038/nature16172).

- Russell, C., et al. (2016), [doi:10.1126/science.aaf4219](https://doi.org/10.1126/science.aaf4219). - Park, R., et al. (2016), Lunar and Planetary Science Conference, vol. 47, p. 1781. - Schmidt, B. E., et al. (2017), [doi:10.1038/ngeo2936](https://doi.org/10.1038/ngeo2936) - Zolotov, M. Y. (2017), [doi:https://doi.org/10.1016/j.icarus.2017.06.018](https://doi.org/10.1016/j.icarus.2017.06.018). - Carrozzo, F., et al. (2018), Nature, formation and distribution of carbonates on ceres, Science Advances. - Raponi, A., et al. (2018), Variations in the amount of water ice on ceres' surface suggest a seasonal water cycle, Science Advances. - McCord, T., and C. Sotin (2005),

doi:10.1029/2004JE002244. - Neveu, M., and S. Desch (2015), Geochemistry, thermal evolution, and cryovolcanism on Ceres with a muddy ice mantle, *Geophys. Res. Lett.*