



Publication Year	2018
Acceptance in OA	2023-01-16T16:11:04Z
Title	Erratum: Tracing black hole accretion with SED decomposition and IR lines: from local galaxies to the high-z Universe
Authors	GRUPPIONI, Carlotta, Berta, S., SPINOGLIO, Luigi Giuseppe Maria, Pereira-Santaella, M., Pozzi, F., Andreani, P., BONATO, MATTEO, De Zotti, G., Malkan, M., Negrello, M., Vallini, L., Vignali, C.
Publisher's version (DOI)	10.1093/mnras/sty772
Handle	http://hdl.handle.net/20.500.12386/32870
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Volume	477

Erratum: Tracing black hole accretion with SED decomposition and IR lines: from local galaxies to the high- z Universe

by C. Gruppioni,¹★ S. Berta,² L. Spinoglio,³ M. Pereira-Santaella,⁴ F. Pozzi,⁵ P. Andreani,⁶ M. Bonato,⁷ G. De Zotti,⁸ M. Malkan,⁹ M. Negrello,¹⁰ L. Vallini^{1,5} and C. Vignali⁵

¹*Istituto Nazionale di Astrofisica - Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127 Bologna, Italy*

²*Max-Planck-Institut für Extraterrestrische Physik (MPE), Postfach 1312, D-85741 Garching, Germany*

³*Istituto Nazionale di Astrofisica - Istituto di Astrofisica e Planetologia Spaziali, via Fosso del Cavaliere 100, 00133 Roma*

⁴*Centro de Astrobiología (CSIC/INTA), Ctra de Torrejón a Ajalvir, km 4, E-28850, Torrejón de Ardoz, Madrid, Spain*

⁵*Dipartimento di Fisica e Astronomia, Università di Bologna, viale Berti Pichat 6, I-40127 Bologna, Italy*

⁶*European Southern Observatory, Karl-Schwarzschild-Straße 2, D-85748 Garching, Germany*

⁷*Department of Physics and Astronomy, Tufts University, 574 Boston Ave., Medford, MA 02155, USA*

⁸*Istituto Nazionale di Astrofisica - Osservatorio Astronomico di Padova, vicolo dell'Osservatorio 5, I-35122 Padova, Italy*

⁹*Department of Physics & Astronomy, UCLA, Los Angeles, CA 90095-1547, USA*

¹⁰*School of Physics and Astronomy, Cardiff University, Queen's Buildings, The Parade, Cardiff CF24 3AA, UK*

Key words: errata, addenda – galaxies: active – galaxies: evolution – galaxies: Seyfert – galaxies: luminosity function – infrared: galaxies.

This is an erratum to the paper ‘Tracing black hole accretion with SED decomposition and IR lines: from local galaxies to the high- z Universe’, by C. Gruppioni et al., published in MNRAS, 458, 4297 (2016).

Here we replace Table 4. In the original version of the Table, a few reported values of the coefficients were not correct (and did not match with those reported in Fig. 10). The values printed in the Figures are the correct ones and were also used for the all the calculations in the paper. The rest of the paper and our conclusions are unaffected by the Table correction.

Table 4. Best-fitting parameters of the line-luminosity relations: $\log_{10}(L_{\text{line}}) = a \cdot \log_{10}(L_{\text{IR}}) + b$ [equation 1]; $\log_{10}(L_{\text{line}}) = c \cdot \log_{10}(L_{\text{bol}}^{\text{AGN}}) + d$ [equation 2]; $\log_{10}(L_{\text{line}}) = e \cdot \log_{10}(L_{\text{IR}}^{\text{SF}}) + f$ [equation 3].

Line	a	b	c	d	e	f	$f_{\text{AGN}}(5\text{--}40\ \mu\text{m})$
PAH 6.2	0.92 ± 0.02	-1.91 ± 0.27	0.58 ± 0.03	2.17 ± 0.27	0.94 ± 0.02	-2.09 ± 0.25	<40 per cent
			0.83 ± 0.03	-1.26 ± 0.34			>40 per cent
PAH 11.2	0.95 ± 0.03	-2.41 ± 0.31	0.52 ± 0.04	2.58 ± 0.30	0.93 ± 0.03	-2.12 ± 0.30	<40 per cent
			0.75 ± 0.03	-0.53 ± 0.36			>40 per cent
[Ne II] 12.8	0.87 ± 0.19	-2.17 ± 0.21	0.57 ± 0.02	1.61 ± 0.23	0.89 ± 0.02	-2.20 ± 0.19	<40 per cent
			0.72 ± 0.02	-0.72 ± 0.22			>40 per cent
[Ne V] 14.3	0.89 ± 0.03	-2.98 ± 0.40	0.66 ± 0.01	-0.17 ± 0.16	0.87 ± 0.03	-2.78 ± 0.38	<40 per cent
					0.93 ± 0.03	-2.68 ± 0.28	>40 per cent
[Ne III] 15.5	0.92 ± 0.04	-2.99 ± 0.40	0.58 ± 0.01	1.04 ± 0.13	0.92 ± 0.04	-2.99 ± 0.40	<40 per cent
					0.78 ± 0.02	-0.84 ± 0.23	>40 per cent
[S III] 18.7	0.82 ± 0.02	-1.93 ± 0.22	0.52 ± 0.02	1.66 ± 0.24	0.78 ± 0.02	-1.38 ± 0.23	<40 per cent
			0.66 ± 0.02	-0.27 ± 0.28			>40 per cent
[Ne V] 24.3	0.78 ± 0.04	-1.49 ± 0.39	0.53 ± 0.02	1.40 ± 0.17	0.77 ± 0.04	-1.40 ± 0.39	<40 per cent
					0.88 ± 0.03	-2.02 ± 0.31	>40 per cent
[O IV] 25.9	0.91 ± 0.03	-2.69 ± 0.35	0.59 ± 0.01	1.19 ± 0.15	0.90 ± 0.03	-2.50 ± 0.35	<40 per cent
					0.84 ± 0.03	-1.10 ± 0.28	>40 per cent
[S III] 33.5	0.72 ± 0.05	-0.53 ± 0.56	0.27 ± 0.04	4.85 ± 0.43	0.76 ± 0.06	-0.78 ± 0.61	<40 per cent
			0.49 ± 0.30	2.00 ± 3.49			>40 per cent
[Si II] 34.8	0.94 ± 0.02	-2.83 ± 0.26	0.51 ± 0.02	2.21 ± 0.25	0.87 ± 0.02	-1.94 ± 0.25	<40 per cent
			0.69 ± 0.03	-0.37 ± 0.31			>40 per cent

* E-mail: carlotta.gruppioni@oabo.inaf.it

Table 4 – *continued*

Line	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>	<i>e</i>	<i>f</i>	$f_{\text{AGN}}(5\text{--}40\ \mu\text{m})$
[N III] 57.3	0.71 ± 0.09	-0.67 ± 0.50	0.70 ± 0.09	0.08 ± 1.04	0.71 ± 0.04	-0.47 ± 0.46	<40 per cent
			0.35 ± 0.06	2.87 ± 0.67			>40 per cent
[O I] 63	0.72 ± 0.04	-0.08 ± 0.35	0.80 ± 0.04	-0.72 ± 0.48	0.70 ± 0.03	0.32 ± 0.35	<40 per cent
			0.54 ± 0.04	1.77 ± 0.41			>40 per cent
[O III] 88	0.88 ± 0.04	-2.11 ± 0.40	1.09 ± 0.05	-4.20 ± 0.53	0.83 ± 0.035	-1.40 ± 0.39	<40 per cent
			0.61 ± 0.05	0.79 ± 0.56			>40 per cent
[N II] 122	0.88 ± 0.04	-2.76 ± 0.37	0.91 ± 0.04	-2.62 ± 0.48	0.86 ± 0.03	-2.48 ± 0.36	<40 per cent
			0.53 ± 0.05	0.73 ± 0.54			>40 per cent
[O I] 145	0.84 ± 0.03	-2.38 ± 0.38	0.93 ± 0.04	-3.11 ± 0.48	0.79 ± 0.03	-1.70 ± 0.37	<40 per cent
			0.68 ± 0.05	-0.60 ± 0.55			>40 per cent
[C II] 158	0.96 ± 0.03	-2.73 ± 0.31	0.79 ± 0.04	-0.37 ± 0.46	0.94 ± 0.03	-2.39 ± 0.30	<40 per cent
			0.80 ± 0.03	-1.24 ± 0.38			>40 per cent

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.