



Publication Year	2015
Acceptance in OA	2020-03-30T14:34:19Z
Title	Corrigendum to "Optical constants of ethylene glycol over an extremely wide spectral range" [Opt. Mater. 37 (2014) 36-41]
Authors	Sani, Elisa, DELL'ORO, Aldo
Publisher's version (DOI)	10.1016/j.optmat.2015.06.039
Handle	http://hdl.handle.net/20.500.12386/23706
Journal	OPTICAL MATERIALS
Volume	48

**Corrigendum to “Optical constants of ethylene glycol over an extremely wide spectral range”
[Optical Materials vol. 37 (2014) 36-41]**

Elisa Sani^a, Aldo Dell’Oro^b

^a INO-CNR, Istituto Nazionale di Ottica, Largo E. Fermi, 6, 50125 Firenze, Italy

^b INAF Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5, 50125 Firenze, Italy

The authors would like to integrate the information contained in the printed version of the above article by providing numerical data concerning Fig. 2 of the mentioned paper. A part of these data were already reported in Table 1 of the mentioned paper in a reduced form for space reasons.

The data are given as 5-column files and should be read as follows:

- First column: Wavenumbers (cm⁻¹)
- Second column: n
- Third column: Uncertainty on n ($\pm\Delta n$)
- Fourth column: k
- Fifth column: Uncertainty on k ($\pm\Delta k$)

As indicated in Table 1 of the mentioned paper, it exist a spectral region where the optical constant k is very low and its value lies below the experimental sensitivity. Thus to provide usable data files (i.e. data in agreement with the required formatting) also in this spectral region, which roughly goes, discontinuously, from 11451 to about 37000 cm⁻¹, we have chosen to impose $k=\Delta k$ in the attached data. From physical point-of-view this means that the correct value of k is below the indicated value.

Finally it should be noticed that the printed version of the above paper showed three graphs in Figure 2. However, to provide numerical data with enough accuracy to appreciate all the spectral features of optical constants, and considering that a single downloadable data file is forced to contain 250 rows maximum, we have chosen here to split the 5000-54000 cm⁻¹ interval in two parts, namely 5000-15000 and 15000-54000 cm⁻¹.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.optmat.2015.06.039>

DOI of original article: 10.1016/j.optmat.2014.04.035

DOI of supplementary material: 10.1016/j.optmat.2015.06.039

Corresponding Author name: Elisa Sani

E-mail address: elisa.sani@ino.it