



Publication Year	2021
Acceptance in OA	2022-03-28T15:29:49Z
Title	Late-time radio observations of the short GRB200522A: constraints on the magnetar model
Authors	Bruni, Gabriele, O'Connor, B., Matsumoto, T., Troja, E., Piran, T., PIRO, LUIGI, RICCI, ROBERTO
Publisher's version (DOI)	10.1093/mnrasl/slab046
Handle	http://hdl.handle.net/20.500.12386/31982
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY. LETTERS
Volume	505

Late-time radio observations of the short GRB 200522A: constraints on the magnetar model

G. Bruni¹★, B. O'Connor^{2,3,4,5}★, T. Matsumoto^{6,7,8}★, E. Troja^{2,3}, T. Piran⁶, L. Piro¹ and R. Ricci^{9,10}

¹INAF – Istituto di Astrofisica e Planetologia Spaziali, via Fosso del Cavaliere 100, I-00133 Roma, Italy

²Astrophysics Science Division, NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, MD 20771, USA

³Department of Astronomy, University of Maryland, College Park, MD 20742-4111, USA

⁴Department of Physics, The George Washington University, 725 21st Street NW, Washington, DC 20052, USA

⁵Astronomy, Physics, and Statistics Institute of Sciences (APSIS), The George Washington University, Washington, DC 20052, USA

⁶Racah Institute of Physics, Edmund J. Safra Campus, Hebrew University of Jerusalem, Jerusalem 91904, Israel

⁷Research Center for the Early Universe, Graduate School of Science, University of Tokyo, Tokyo 113-0033, Japan

⁸Department of Physics, Graduate School of Science, University of Tokyo, Tokyo 113-0033, Japan

⁹Istituto Nazionale di Ricerche Metrologiche – Torino, Strada delle Cacce 91, I-10135 Torino, Italy

¹⁰INAF – Istituto di Radioastronomia, via Gobetti 101, I-40129 Bologna, Italy

Accepted 2021 May 3. Received 2021 April 29; in original form 2021 April 1

ABSTRACT

GRB 200522A is a short duration gamma-ray burst (GRB) at redshift $z=0.554$ characterized by a bright infrared counterpart. A possible, although not unambiguous, interpretation of the observed emission is the onset of a luminous kilonova powered by a rapidly rotating and highly magnetized neutron star, known as magnetar. A bright radio flare, arising from the interaction of the kilonova ejecta with the surrounding medium, is a prediction of this model. Whereas the available data set remains open to multiple interpretations (e.g. afterglow, r-process kilonova, magnetar-powered kilonova), long-term radio monitoring of this burst may be key to discriminate between models. We present our late-time upper limit on the radio emission of GRB 200522A, carried out with the Karl G. Jansky Very Large Array at 288 d after the burst. For kilonova ejecta with energy $E_{\text{ej}} \approx 10^{53}$ erg, as expected for a long-lived magnetar remnant, we can already rule out ejecta masses $M_{\text{ej}} \lesssim 0.03 M_{\odot}$ for the most likely range of circumburst densities $n \gtrsim 10^{-3} \text{ cm}^{-3}$. Observations on timescales of $\approx 3\text{--}10$ yr after the merger will probe larger ejecta masses up to $M_{\text{ej}} \sim 0.1 M_{\odot}$, providing a robust test to the magnetar scenario.

Key words: gamma-ray bursts: individual: GRB 200522A – stars: magnetars – stars: neutron.

1 INTRODUCTION

Short duration gamma-ray bursts (GRBs) are brief flashes of high-energy radiation produced, at least in some cases, by the coalescence of two neutron stars (NSs; Eichler et al. 1989), as demonstrated by the association between the binary NS merger GW170817 and the short GRB 170817A (Abbott et al. 2017a, b). The fate of the merger remnant, either a massive NS or a black hole (BH), is still poorly constrained by either gravitational wave or electromagnetic observations (Ai et al. 2018; Pooley et al. 2018; Piro et al. 2019; Abbott et al. 2019; van Putten & Della Valle 2019), and remains a central question of the field with important implications for fundamental physics such as the nuclear equation of state (EoS).

After the merger, the lifetime of the surviving NS primarily depends on its mass and the unknown EoS of cold dense matter (Lattimer & Prakash 2016). Soft EoSs favour a prompt collapse to a BH, whereas stiffer EoSs allow for a dynamically stable NS (Giacomazzo & Perna 2013; Piro, Giacomazzo & Perna 2017). If the remnant NS is also highly magnetized, its large reservoir of rotational

energy, up to 10^{53} erg for an NS mass of $2.1 M_{\odot}$, could be transferred to the merger ejecta by the NS dipole spin-down radiation (Metzger & Bower 2014). These magnetar-driven winds accelerate the ejecta to transrelativistic speeds and substantially decrease its opacity by mixing (Metzger, Thompson & Quataert 2018). Note also that any long-lived NS would decrease the opacity by neutrino irradiation (e.g. Metzger & Fernandez 2014). When the ejecta undergo radioactive decays, the emitted signal, commonly referred to as either kilonova or macronova, is expected to peak at optical/ultraviolet energies within a few days after the merger, appearing bluer and brighter than a purely r-process powered kilonova (e.g. Yu, Zhang & Gao 2013; Wollaeger et al. 2019). On longer time-scales, this fast ejecta interacts with the surrounding medium, producing a luminous radio afterglow, visible months to years after the merger (Nakar & Piran 2011). In some cases, radio emission from the nascent pulsar wind nebula may also become visible a few months after the merger (Piro & Kulkarni 2013).

Based on these models, events associated with bright and blue kilonovae are promising targets to search for a late-time radio transient and test whether the luminosity of the optical and near-infrared (nIR) emission is the result of a long-lived magnetar. Past late-time radio surveys of short GRB locations all resulted in a non-

* E-mail: gabriele.bruni@inaf.it (GB); oconnorb@gwmail.gwu.edu (BO); tatsuya.matsumoto@mail.huji.ac.il (TM)

detection, thus disfavouring the presence of a stable magnetar in a large fraction of short GRBs (Fong et al. 2016; Horesh et al. 2016; Klose et al. 2019; Schroeder et al. 2020; Grandorf et al. 2021; Ricci et al. 2021), and future all-sky radio surveys will provide additional constraints independent of short GRBs (Beniamini & Lu 2021). The tightest upper limit for a short GRB with a kilonova candidate was derived for GRB 160821B (Lamb et al. 2019; Troja et al. 2019), and excluded ejecta masses $\lesssim 0.14 M_{\odot}$ for densities $n \gtrsim 10^{-3} \text{ cm}^{-3}$ and an energy deposition of $E \approx 10^{53} \text{ erg}$ (Ricci et al. 2021). An upper limit $M_{\text{ej}} \lesssim 0.06 M_{\odot}$ was also placed for GRB 150101B, a nearby short burst associated with a blue kilonova (Troja et al. 2018). More recently, another event, GRB 200522A at redshift $z = 0.554$, was associated with a luminous nIR counterpart whose properties are open to multiple interpretations, including a bright kilonova emission. Fong et al. (2021) suggested that the observed nIR signal could be the result of a magnetar-boosted kilonova, peaking at optical wavelengths. A bright late-time radio flare is therefore a natural expectation of this scenario.

Additional observational evidence, presented by O’Connor et al. (2021), favoured other interpretations. In particular, optical imaging simultaneous to the nIR observations does not detect a bright counterpart, whereas late-time X-ray observations find evidence of a long-lasting, non-thermal emission. Overall, this evidence points to a substantial afterglow contribution up to late times. This may account for most, if not all, of the nIR signal if a modest amount of dust is present along the line of sight. In any case, the observations of a bright, regular short GRB and afterglow modeling strongly suggest that GRB 200522A was viewed along its jet’s axis. For this particular orientation, other factors may brighten the kilonova emission without invoking an additional energy source (Korobkin et al. 2020; Heinzel et al. 2021). A late-time radio flare may still be produced but it would peak at a later time and at a lower luminosity than in the magnetar scenario.

Recently we carried out radio observations of this event, obtaining a deep upper limit on radio emission almost a year after the event. Whereas the optical and nIR data set is consistent with different interpretations, we show here that late time radio observations are a suitable tool to determine whether a powerful magnetar has operated in this event and has boosted the nIR signal. We discuss the observations in Section 2 and afterglow and kilonova constraints in Section 3. We describe the model used to interpret the afterglow observations and the results in Section 4. We conclude with implications and the importance of further observations in Section 5.

2 RADIO OBSERVATIONS

At early times, GRB 200522A was observed by Fong et al. (2021) with the Karl G. Jansky Very Large Array (VLA) in C configuration. A weak radio counterpart was detected in C band (6 GHz) a few hours after the burst and seen to fade below detection threshold within a few days. This radio signal may be produced either by a standard forward shock (FS) emission, a rapidly fading reverse shock (RS) component, or a combination of both. In any case, it was no longer detected at $T_0 + 11 \text{ d}$ with a 3σ upper limit of $14 \mu\text{Jy}$.

We re-observed the field with the VLA on 2021 February 19, ($T_0 + 288 \text{ d}$), with the array in A configuration (P.I. Troja). A single 2-h run was carried out, applying a modified C band (6 GHz) setup to avoid radio frequency interferences from the Clarke belt satellites, and involving all the 28 antennas of the array. The net time on source was 1h 22 min. The source 3C 147, observed at the beginning of the run, was used as both a bandpass and flux density scale calibrator, while the phase calibrator paired to the target was J0022+0014. A

phase-referencing cycle of 8 min, alternating between calibrator and target, was applied.

Data were reduced with the the Common Astronomy Software Applications package (CASA,¹ McMullin et al. 2007). We processed the raw data with VLA pipeline version 5.6.2-3. Imaging was performed with the `tclean` task, applying natural weighting. This led to an angular resolution of $0.53 \text{ arcsec} \times 0.36 \text{ arcsec}$. No emission was detected at the position of the radio afterglow detected by Fong et al. (2021). We measure an RMS noise of $2.6 \mu\text{Jy}$ per beam, resulting in a 3σ upper limit of $7.8 \mu\text{Jy}$ per beam. At a distance $z \sim 0.554$ (O’Connor et al. 2021), this corresponds to a luminosity $L_{\nu} \lesssim 10^{29} \text{ erg s}^{-1} \text{ Hz}^{-1}$ at 185 d (rest frame) since the merger.

3 AFTERGLOW AND KILONOVA CONSTRAINTS

We use the afterglow models of O’Connor et al. (2021) to bracket the range of possible circumburst densities. These models provide a good description of the broadband data set, which includes X-ray data from *Swift* and *Chandra*, radio data from the VLA, optical photometry from Gemini, and nIR photometry from the *Hubble Space Telescope*. Since we wish to test the hypothesis of a luminous kilonova emission, we only consider models of a forward shock (FS) plus black-body (BB) emission.

Multiple solutions can fit the available data equally well: (a) a bright early-peaking FS expanding into a homogeneous medium with density $10^{-4} \text{ cm}^{-3} \lesssim n \lesssim 10^{-1} \text{ cm}^{-3}$ (1σ), and (b) a bright RS plus an FS expanding in a medium with a slightly lower density $2 \times 10^{-5} \text{ cm}^{-3} \lesssim n \lesssim 10^{-2} \text{ cm}^{-3}$. In both cases, the total (beaming corrected) jet energy is $E_j \approx 4 \times 10^{49} \text{ erg}$. This set of solutions also includes the models presented in Fong et al. (2021), who fix the fraction of the electron energy to the total kinetic energy $\epsilon_e = 0.1$ and the fraction of magnetic field energy $\epsilon_B = 0.01$. Extremely low-density solutions with $n \approx 2 \times 10^{-5} \text{ cm}^{-3}$ appear unlikely given the location of the GRB within its host ($\sim 1.1 \text{ kpc}$ from the host galaxy’s nucleus; Fong et al. 2021; O’Connor et al. 2021), and we therefore focus our work on the range $10^{-4} \text{ cm}^{-3} \lesssim n \lesssim 10^{-1} \text{ cm}^{-3}$ (O’Connor et al. 2021).

A kilonova component, although not required, can be constrained by the data as shown in Fig. 1. After subtracting the afterglow contribution, the residual optical and nIR emission can be modelled with a simple BB with rest-frame temperature $T \sim 4300 \text{ K}$ (O’Connor et al. 2021). This is consistent with a model of a radioactively powered kilonova with large ejecta mass $0.03 M_{\odot} \lesssim M_{\text{ej}} \lesssim 0.1 M_{\odot}$ (Wollaeger et al. 2018; Korobkin et al. 2020). In the magnetar-boosted scenario, comparably massive ejecta, $0.01 M_{\odot} \lesssim M_{\text{ej}} \lesssim 0.1 M_{\odot}$, can reproduce the observed nIR luminosity. However, the peak of the kilonova emission lies in the optical range ($T \sim 6400\text{--}7000 \text{ K}$; Fong et al. 2021) and overpredicts the optical flux by a factor of $\gtrsim 8$. In order for the optical emission to be consistent with this model, an intrinsic extinction of $A_V \gtrsim 1 \text{ mag}$ should be considered.

Due to the limited data set, neither scenario can be firmly ruled out on the basis of the optical/nIR data alone. However, the predictions for a late-time radio flare differ greatly. In the magnetar model, a remnant NS with $M \sim 2.1 M_{\odot}$ and initial spin period $P_0 \sim 0.7 \text{ ms}$ will impart to the ejecta, a large kinetic energy, $E_{\text{ej}} \approx 10^{53} \text{ erg}$. Whereas for a radioactively powered kilonova with ejecta mass $M_{\text{ej}} \sim 0.03\text{--}0.1 M_{\odot}$ and velocity $v_{\text{ej}} \sim 0.15c$, as presented by O’Connor et al. (2021), the energy is significantly smaller ($\approx 10^{51} \text{ erg}$). Late-time

¹<https://casa.nrao.edu/>




Figure 1. Flux density in the optical/nIR bands (after subtraction of the afterglow component) for GRB 200522A (O’Connor et al. 2021). Data are compared to the predicted emission from a magnetar-boosted kilonova (blue shaded region; Fong et al. 2021), and a radioactively powered kilonova from a lanthanide-poor ($Y_e = 0.37$) wind ejected along the polar axis (grey shaded region; O’Connor et al. 2021). The red dashed line shows a simple BB spectrum with $T \sim 4300$ K (rest frame).

radio observations can discern between these two scenarios as we discuss below.

4 IMPLICATION OF THE LATE RADIO OBSERVATIONS

To test the magnetar-boosted model, we calculate the radio flare arising from the energized kilonova ejecta. The kilonova ejecta with a kinetic energy E_{ej} and mass M_{ej} propagates into an ISM with a density n . At the shock front, the magnetic field is amplified and electrons are accelerated to a power-law distribution, radiating synchrotron emission (Nakar & Piran 2011). Our calculation method follows Ricci et al. (2021). In this particular case, we do not consider the afterglow suppression by the relativistic jet (Margalit & Piran 2020), whose time-scale is much shorter ($\lesssim 0.2$ yr) than the time of our observations. Following the magnetar-boosted model presented in Fong et al. (2021), we focus mainly on a kinetic energy of $E_{ej} = 10^{53}$ erg.

Fig. 2 depicts the radio light curves for different ejecta masses of $M_{ej} = 0.01 - 0.1 M_\odot$, consistent with the kilonova modeling and densities of $n = 10^{-3}$ and 10^{-2} cm^{-3} , representative of the median values found through broad-band afterglow fitting. We set the power-law index of the electron distribution $p = 2.5$, and microphysical parameters $\epsilon_e = 0.1$ and $\epsilon_B = 0.01$ identical to Fong et al. (2021). We adopt these parameters throughout unless otherwise specified. The observed frequency, 6 GHz, is typically larger than both the synchrotron frequency ν_m and the self-absorption frequency ν_a resulting from the adopted parameters (see Ricci et al. 2021). The light-curve peaks at the deceleration time, with more massive ejecta leading to a later peak and a lower peak flux.

By imposing that the flux F_ν is smaller than the observed upper limit, we derive a lower limit on the ejecta mass. With $E_{ej} = 10^{53}$ erg the outflow is in the relativistic (or at least mildly relativistic) regime in which the flux depends most sensitively on the Lorentz factor Γ




Figure 2. Light curves of GRB 200522A for various ejecta masses of $M_{ej} = 0.01, 0.03$, and $0.1 M_\odot$ (the corresponding initial Lorentz factors $\Gamma_{in} = 6.6, 2.9$, and 1.6 or $\beta_{in} = 0.77$ for the last case). Solid and dashed lines correspond to densities 10^{-2} and 10^{-3} cm^{-3} , respectively. The other parameters are $\epsilon_e = 0.1$, $\epsilon_B = 0.01$, and $p = 2.5$. Gray triangles show 3σ flux density upper limits from earlier observations by Fong et al. (2021), and our late-time monitoring.

and hence on the mass:

$$F_\nu \simeq 40 \mu\text{Jy} \epsilon_{e,-1}^{p-1} \epsilon_{B,-2}^{(p+1)/4} n_{-3}^{(p+5)/4} d_{28}^{-2} \nu_{\text{GHz}}^{(1-p)/2} t_{\text{yr}}^3 \left(\frac{\Gamma}{2} \right)^{2(p+3)}, \quad (1)$$

where d , ν , and t are the distance (cm), observing frequency (GHz), and time since the merger (yr), respectively. We use the notation $Q_x = Q/10^x$ in cgs units for the other quantities. The upper limit on the observed flux gives an upper limit on the Lorentz factor which corresponds to a lower limit on the ejecta mass by $E_{ej} \simeq \Gamma M_{ej} c^2$:

$$M_{ej} \gtrsim 0.03 M_\odot E_{ej,53} \left[\epsilon_{e,-1}^{p-1} \epsilon_{B,-2}^{(p+1)/4} n_{-3}^{(p+5)/4} d_{28}^{-2} t_{\text{yr}}^3 \nu_{\text{GHz}}^{(1-p)/2} F_{10\mu\text{Jy}}^{-1/2} \right]^{\frac{1}{p+3}}, \quad (2)$$

where $F_{10\mu\text{Jy}} = F_\nu/10 \mu\text{Jy} \sim 0.78$ is our limit. Note that this mass limit depends linearly on the energy but it is almost independent from the uncertain parameters (n , ϵ_e , and ϵ_B).

Fig. 3 shows the parameter space of ejecta mass and energy ruled out by our upper limit of $7.8 \mu\text{Jy}$. For an ejecta energy of $E_{ej} = 10^{53}$ erg, the current limit reasonably rejects ejecta mass lower than $M_{ej} \lesssim 0.03 M_\odot$ for the most likely range of densities $n \gtrsim 10^{-3} \text{ cm}^{-3}$. Note that equation (2) underestimates the mass limit by a factor of a few for $n = 0.1 \text{ cm}^{-3}$. This is because ejecta with $E_{ej} \lesssim M_{ej} c^2$ are in the Newtonian regime, so equations (1) and (2) should be replaced with those for the Newtonian limit:

$$F_\nu \simeq 150 \mu\text{Jy} \epsilon_{e,-1}^{p-1} \epsilon_{B,-2}^{(p+1)/4} n_{-1}^{(p+5)/4} d_{28}^{-2} \nu_{\text{GHz}}^{(1-p)/2} t_{\text{yr}}^3 \left(\frac{\beta}{0.8} \right)^{\frac{5p+3}{2}}, \quad (3)$$

$$M_{ej} \gtrsim 0.3 M_\odot E_{ej,53} \left[\epsilon_{e,-1}^{p-1} \epsilon_{B,-2}^{(p+1)/4} n_{-3}^{(p+5)/4} d_{28}^{-2} t_{\text{yr}}^3 \nu_{\text{GHz}}^{(1-p)/2} F_{10\mu\text{Jy}}^{-1/2} \right]^{\frac{8}{5p+3}}, \quad (4)$$

where β is the ejecta velocity normalized by the speed of light. See also equation (26) in Ricci et al. (2021) for the deep-Newtonian case, where $\beta \lesssim 0.2$.




Figure 3. Constraints on the kinetic energy for the allowed range of densities (10^{-4} to 10^{-1} cm^{-3}) for GRB 200522A. For each density value, the range of energies lying above the curve is ruled out by the radio data. The other parameters are fixed to $\varepsilon_e = 0.1$, $\varepsilon_B = 0.01$, and $p = 2.5$. The blue shaded region represents the relevant range of ejecta mass.




Figure 4. Ejecta mass ruled out by null detection $F_v < 7.8 \mu\text{Jy}$ as a function of time. For a fixed density, only values above each curve are allowed by the data. The vertical orange line denotes the observation time in this work (288 d). The shaded region represents the relevant range of ejecta mass.

Later observations are needed to exclude more massive ejecta. Fig. 4 depicts the ejecta mass ruled out by a similar null-detection as a function of time. Our limit can rule out $M_{\text{ej}} \lesssim 0.14 M_{\odot}$ for an environment denser than $n \sim 0.1 \text{ cm}^{-3}$, which is consistent with the small projected offset from the galaxy’s center and the afterglow observations, although with low posterior probability ($\lesssim 15$ per cent). For lower densities of $n = 10^{-2}$ and $n = 10^{-3} \text{ cm}^{-3}$, favoured by the afterglow data, observations in a few years will test ejecta of $0.03\text{--}0.1 M_{\odot}$. An observation at 3 yr after the merger would be effective in excluding masses $M_{\text{ej}} \lesssim 0.1 M_{\odot}$ and $M_{\text{ej}} \lesssim 0.05 M_{\odot}$ for $n = 10^{-2}$ and $n = 10^{-3} \text{ cm}^{-3}$, respectively. For even lower densities $n < 10^{-4} \text{ cm}^{-3}$, the peak flux falls below the expected upper limit and late observations will not result in a significantly better constraint.

Finally, we note that merger ejecta with kinetic energy of $E_{\text{ej}} \approx 10^{51} \text{ erg}$, as suggested by O’Connor et al. (2021), is allowed by the observations. This is more explicitly demonstrated in Fig. 5.




Figure 5. The same as Fig. 2 but for the standard kilonova scenario ($E_{\text{ej}} = 10^{51} \text{ erg}$ and $M_{\text{ej}} = 0.03$ and $0.1 M_{\odot}$ corresponding to $\beta_{\text{in}} = 0.19$ and 0.11 , respectively). Black solid line represents the expected 3σ detection threshold of ngVLA in a ~ 30 h exposure.

Within this model, the peak timescale would be $\gtrsim 30$ yr after the merger, and the peak flux is $\lesssim 0.2 \mu\text{Jy}$. Even with an optimistic choice of $\varepsilon_B = 0.1$ and the increased sensitivity of future observatories, the detection of a radio flare would be challenging for this burst. For example, considering the nominal RMS of $0.23 \mu\text{Jy}$ per beam in a 1-h observation with the next generation VLA (ngVLA, Selina et al. 2018), it would take over 30 h to reach a 3σ detection (see also Corsi et al. 2019 for an exhaustive discussion of radio counterparts studies with the ngVLA). This is expected in view of the large distance of this event as compared to those discussed in Nakar & Piran (2011).

5 CONCLUSIONS

We present VLA observations of the short duration GRB 200522A at 288 d post merger. We do not detect a radio source at the GRB position, placing an upper limit of $\nu L_{\nu} \lesssim 9 \times 10^{38} \text{ erg s}^{-1}$ at 9 GHz (rest frame). This is comparable to the limits obtained for the larger sample of cosmological short GRBs at $z \lesssim 0.5$ (Schroeder et al. 2020; Ricci et al. 2021). In order to break the degeneracy in the interpretation of the GRB optical/nIR counterpart, we compare our radio limit to the theoretical predictions for a late-time radio flare produced by the merger ejecta.

We first consider the scenario of a long-lived magnetar re-energizing the merger ejecta to $E_{\text{ej}} = 10^{53} \text{ erg}$. This model was invoked by Fong et al. (2021) to interpret the nIR counterpart as a magnetar-boosted kilonova. Based on our VLA observation, we can exclude ejecta masses $M_{\text{ej}} \lesssim 0.03 M_{\odot}$ for a wide range of densities $n \gtrsim 10^{-3} \text{ cm}^{-3}$, also preferred by the GRB afterglow modeling (O’Connor et al. 2021).

Additional late-time observations will provide even stronger constraints on the magnetar model. Assuming a similar sensitivity of $\approx 8 \mu\text{Jy}$ at 6 GHz, we find that future observations at ~ 3 and ~ 10 yr post-merger would rule out $M_{\text{ej}} \lesssim 0.1 M_{\odot}$ for $n \gtrsim 10^{-2} \text{ cm}^{-3}$ and $n \gtrsim 10^{-3} \text{ cm}^{-3}$, respectively.

Finally, we discuss the standard scenario of a radioactively powered kilonova, also consistent with the optical/nIR data set (O’Connor et al. 2021). This model implies a significantly lower ejecta energy $E_{\text{ej}} \approx 10^{51} \text{ erg}$ as well as large masses, $0.03 M_{\odot} \leq M_{\text{ej}} \leq 0.1 M_{\odot}$, and its associated radio transient would be too faint for detection by both current and future observatories. This is not unexpected in view

of the distance of this object and the prediction for the late radio flares (Nakar & Piran 2011). Events at much closer distance, such as the sample of short bursts identified by Dichiara et al. (2020) and obviously GW170817 (Makhathini et al. 2020; Troja et al. 2020, 2021; Hajela et al. 2021), will be pivotal in constraining the late-time radio emission from the merger ejecta.

ACKNOWLEDGEMENTS

The authors thank the referee for useful comments that improved the manuscript. The authors also thank Chris L. Fryer for providing the kilonova spectra used to model GRB 200522A. ET and BO were supported in part by the National Aeronautics and Space Administration through grant GO21065A. TP was supported by an advanced ERC grant TReX. TM was supported by JSPS Postdoctoral Fellowship, Kakenhi No. 19J00214. LP acknowledges support from the European Union's Horizon 2020 Programme under the AHEAD2020 project (grant agreement no. 871158) and from MIUR, PRIN 2017 (grant 20179ZF5KS). The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Abbott B. P. et al., 2017a, *ApJ*, 848, L12
 Abbott B. P. et al., 2017b, *ApJ*, 848, L13
 Abbott B. P. et al., 2019, *Phys. Rev. X*, 9, 011001
 Ai S., Gao H., Dai Z.-G., Wu X.-F., Li A., Zhang B., Li M.-Z., 2018, *ApJ*, 860, 57
 Beniamini P., Lu W., 2021, preprint (arXiv:2104.01181)
 Corsi A. et al., 2019, *BAAS*, 51, 209
 Dichiara S., Troja E., O'Connor B., Marshall F. E., Beniamini P., Cannizzo J. K., Lien A. Y., Sakamoto T., 2020, *MNRAS*, 492, 5011
 Eichler D., Livio M., Piran T., Schramm D. N., 1989, *Nature*, 340, 126
 Fong W., Metzger B. D., Berger E., Özel F., 2016, *ApJ*, 831, 141
 Fong W. et al., 2021, *ApJ*, 906, 127
 Giacomazzo B., Perna R., 2013, *ApJ*, 771, L26.
 Grandorf C. et al., 2021, *ApJ*, 908, 63
 Hajela A. et al., 2021, preprint (arXiv:2104.02070)
 Heinzel J. et al., 2021, *MNRAS*, 502, 3057
 Horesh A., Hotokezaka K., Piran T., Nakar E., Hancock P., 2016, *ApJ*, 819, L22
 Klose S. et al., 2019, *ApJ*, 887, 206
 Korobkin O. et al., 2020, *ApJ*, 910, 116
 Lamb G. P. et al., 2019, *ApJ*, 883, 48
 Lattimer J. M., Prakash M., 2016, *Phys. Rev.*, 621, 127
 Makhathini S. et al., 2020, preprint (arXiv:2006.02382)
 Margalit B., Piran T., 2020, *MNRAS*, 495, 4981
 McMullin J. P., Waters B., Schiebel D., Young W., Golap K., 2007, *ASP Conf Series*, Vol 376., Astron. Soc. Pac., San Francisco. 127
 Metzger B. D., Bower G. C., 2014, *MNRAS*, 437, 1821
 Metzger B. D., Fernández R., 2014, *MNRAS*, 441, 3444
 Metzger B. D., Thompson T. A., Quataert E., 2018, *ApJ*, 856, 101
 Nakar E., Piran T., 2011, *Nature*, 478, 82
 O'Connor B. et al., 2021, *MNRAS*, 502, 1279
 Piro A. L., Kulkarni S. R., 2013, *ApJ*, 762, L17
 Piro A. L., Giacomazzo B., Perna R., 2017, *ApJ*, 844, L19
 Piro L. et al., 2019, *MNRAS*, 483, 1912
 Pooley D., Kumar P., Wheeler J. C., Grossan B., 2018, *ApJ*, 859, L23
 Ricci R. et al., 2021, *MNRAS*, 500, 1708
 Schroeder G. et al., 2020, *ApJ*, 902, 82
 Selina R. J. et al., 2018, *Proc. SPIE Conf. Ser. Vol. 10700*. *SPIE*, Bellingham, p. 107001O
 Troja E. et al., 2018, *Nat. Commun.*, 9, 4089
 Troja E. et al., 2019, *MNRAS*, 489, 2104
 Troja E. et al., 2020, *MNRAS*, 498, 5643
 Troja E., O'Connor B., Ryan G., Piro L., Ricci R., Zhang B. et al., 2021, *MNRAS*, submitted, preprint (arXiv:2104.13378)
 van Putten M. H. P. M., Della Valle M., 2019, *MNRAS*, 482, L46
 Wollaeger R. T. et al., 2018, *MNRAS*, 478, 3298
 Wollaeger R. T., Fryer C. L., Fontes C. J., Lippuner J., Vestrand W. T., Mumpower M. R., Korobkin O. et al., 2019, *ApJ*, 880, 22
 Yu Y.-W., Zhang B., Gao H., 2013, *ApJL*, 776, L40

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.