



Publication Year	2018
Acceptance in OA	2020-10-05T09:15:28Z
Title	Chromospheric and Transition Region Emission Properties of G, K, and M dwarf Exoplanet Host Stars
Authors	France, Kevin, Arulanantham, Nicole, Fossati, Luca, LANZA, Antonino Francesco, Linsky, Jeffrey L., Redfield, Seth, Loyd, Robert, Schneider, Christian
Handle	http://hdl.handle.net/20.500.12386/27569
Volume	231

Chromospheric and Transition Region Emission Properties of G, K, and M dwarf Exoplanet Host Stars

American Astronomical Society, AAS Meeting #231, id. 349.06

- [France, Kevin](#) (*University of Colorado*);
- [Arulanantham, Nicole](#) (*University of Colorado*);
- [Fossati, Luca](#) (*Space Research Institute, Austrian Academy of Sciences*);
- [Lanza, A. F.](#) (*INAF-Osservatorio Astrofisico di Catania*);
- [Linsky, Jeffrey L.](#) (*University of Colorado*);
- [Redfield, Seth](#) (*Weslyan University*);
- [Loyd, Robert](#) (*Arizona State University*);
- [Schneider, Christian](#) (*University of Hamburg*)

Abstract

Exoplanet magnetic fields have proven notoriously hard to detect, despite theoretical predictions of substantial magnetic field strengths on close-in extrasolar giant planets. It has been suggested that stellar and planetary magnetic field interactions can manifest as enhanced stellar activity relative to nominal age-rotation-activity relationships for main sequence stars or enhanced activity on stars hosting short-period massive planets. In a recent study of M and K dwarf exoplanet host stars, we demonstrated a significant correlation between the relative luminosity in high-temperature stellar emission lines ($L(\text{ion})/L_{\text{Bol}}$) and the “star-planet interaction strength”, $M_{\text{plan}}/a_{\text{plan}}$. Here, we expand on that work with a survey of G, K, and M dwarf exoplanet host stars obtained in two

recent far-ultraviolet spectroscopic programs with the Hubble Space Telescope. We have measured the relative luminosities of stellar lines C II, Si III, Si IV, and N V (formation temperatures from 30,000 - 150,000 K) in a sample of ~60 exoplanet host stars and an additional ~40 dwarf stars without known planets. We present results on star-planet interaction signals as a function of spectral type and line formation temperature, as well as a statistical comparison of stars with and without planets.