



Publication Year	2003
Acceptance in OA	2024-01-25T12:26:10Z
Title	QSO Absorption Lines as a Cosmological Probe: Exploring the Lyman Forest with VLT/UVES
Authors	BIANCHI, Simone, CRISTIANI, Stefano, Kim, Tae-Sun, D'Odorico, Sandro
Publisher's version (DOI)	10.1007/10857580_35
Handle	http://hdl.handle.net/20.500.12386/34625
Journal	ESO ASTROPHYSICS SYMPOSIA
Serie	ESO ASTROPHYSICS SYMPOSIA

Quasar absorption lines as a cosmological probe: exploring the Lyman forest with VLT/UVES

Simone Bianchi¹, Stefano Cristiani², Tae-Sun Kim¹, and Sandro D’Odorico¹

¹ ESO, Karl-Schwarzschild-Strasse 2, D-85748 Garching, Germany

² ST-ECF, ESO, Karl-Schwarzschild-Strasse 2, D-85748 Garching, Germany

1 Observations and Data Reduction

The remarkable efficiency of the Ultra-Violet Echelle Spectrograph (UVES) at the VLT has made it possible to push high-resolution, high-S/N ground observations of the Ly α forest down to $z \sim 1.5$, gaining new insight into the physical state of the intergalactic medium (IGM) and its evolution over more than 90% of the cosmic time. The results presented here are based on recent UVES observations of eight QSOs spectra, covering the Ly α forest at $1.5 < z_{\text{Ly}\alpha} < 3.6$ with S/N ~ 40 –50 and resolution $R \sim 45\,000$ [1,2]. The absorption lines of the normalized spectra were fit to the Voigt profiles to derive the three line parameters: absorption redshift, z , the HI column density, N_{HI} in cm^{-2} , and the Doppler parameters, b in km s^{-1}).

2 The Number Density Evolution of the Ly α forest

Fig. 1 shows the evolution of the line number density per unit redshift, dn/dz , in the interval $N_{\text{HI}} = 10^{13.6-17} \text{ cm}^{-2}$. The maximum-likelihood fit to the $z > 1.5$ data is $dn/dz = 6.1 (1+z)^{2.5 \pm 0.2}$ (dashed line). The evolution of the Ly α forest with z is mainly governed by two physical processes: the Hubble expansion and the ionizing ultraviolet background flux (UVB,[3]). At higher z , the Hubble expansion and the non-decreasing UVB cause a rapid evolution of dn/dz . At lower z , HST observations have shown a slow-down in dn/dz (solid line in Fig. 1,[4]), which can be explained with a decrease of the UVB flux in the local universe. The UVES observations imply that the turn-off in the evolution occurs at $z \sim 1$. The evolution and the redshift of the turn-off are consistent with an UVB to which galaxies contribute as much as QSOs or more, as long as a fraction $f_{\text{esc}} \geq 0.05$ of the UV flux can escape the internal absorption in a galaxy [5].

3 The cosmic baryon density

A lower-bound to the cosmic baryon density can be derived from the distribution of the Ly α optical depths [6]. For a UVB with a contribution from galaxies ($f_{\text{esc}} = 0.1$), the effective optical depths measured in the UVES spectra at $1.5 < z < 4$ implies $\Omega_b h^{1.5} > 0.028$ (assuming IGM temperature $T = 2 \cdot 10^4 \text{ K}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$). This value is consistent with the BBN value for a low D/H




Fig. 1. The number density evolution of the Ly α forest

primordial abundance. Most of the baryons reside in the Lyman forest at $1.5 < z < 4$ with little change in the contribution to Ω as a function of z . Conversely, given the observed opacity, a higher UVB requires a higher Ω_b . As pointed out by Haehnelt et al. [7], values of f_{esc} as large as 0.4 [8], would result in too large Ω_b values.

4 The Temperature of the IGM

Absorption lines in the Ly α forest are broadened by gas thermal motion and other processes, therefore the minimum b value can provide an upper limit on the temperature. The minimum b is found to increase as z decreases [9]. When the column densities are converted into over-densities, this implies that the temperature at the mean ISM density decreases with z . The large fluctuations in the minimum b value and in its dependence on N_{HI} , even at a similar redshift, suggest that the temperature of the intergalactic medium might fluctuate. A large fluctuation seen at $z \sim 3.1$ is probably due to the HeII reionization [10,9].

References

1. T.-S. Kim, S. Cristiani, S. D'Odorico: A&A, **373**, 757 (2001)
2. T.-S. Kim, et al.: A&A, submitted (2002)
3. T. Theuns, A. Leonard, G. Efstathiou: MNRAS, **297**, 49 (1998)
4. R. J. Weymann et al.: ApJ, **506**, 1 (1998)
5. S. Bianchi, S. Cristiani, T.-S. Kim: A&A, **376**, 1 (2001)
6. D. H. Weinberg et al.: ApJ, **490**, 564 (1997)
7. M. Haehnelt et al.: ApJ, **549**, L151 (2001)
8. C. Steidel, M. Pettini, K. Adelberger: ApJ, **546**, 66 (2001)
9. T.-S. Kim, S. Cristiani, S. D'Odorico: A&A, **383**, 747 (2001)
10. J. Schaye et al.: MNRAS, **318**, 817 (2000)