



Publication Year	2017
Acceptance in OA	2023-05-30T14:06:19Z
Title	Solar system dynamics in general relativity
Authors	Battista, Emmanuele, Dell'Agnello, Simone, Esposito, Giampiero, Di Fiore, Luciano, Simo, Jules, GRADO, ANIELLO
Publisher's version (DOI)	10.1142/S0219887817501171
Handle	http://hdl.handle.net/20.500.12386/34226
Journal	INTERNATIONAL JOURNAL OF GEOMETRIC METHODS IN MODERN PHYSICS
Volume	14

Solar system dynamics in general relativity

Emmanuele Battista,^{*} Giampiero Esposito,[†] and Luciano Di Fiore[‡]

*Istituto Nazionale di Fisica Nucleare, Sezione di Napoli,
Complesso Universitario di Monte S. Angelo, Via Cintia Edificio 6, 80126 Napoli, Italy*

Simone Dell'Agnello[§]

Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali di Frascati, 00044 Frascati, Italy

Jules Simo^{**}

*Aerospace Engineering, Computing & Technology Building,
School of Engineering, University of Central Lancashire, Preston, PR1 2HE, United Kingdom*

Aniello Grado^{††}

INAF, Osservatorio Astronomico di Capodimonte, 80131 Napoli, Italy

(Dated: July 22, 2016)

Recent work in the literature has advocated using the Earth-Moon-planetoid Lagrangian points as observables, in order to test general relativity and effective field theories of gravity in the solar system. However, since the three-body problem of classical celestial mechanics is just an approximation of a much more complicated setting, where all celestial bodies in the solar system are subject to their mutual gravitational interactions, while solar radiation pressure and other sources of nongravitational perturbations also affect the dynamics, it is conceptually desirable to improve the current understanding of solar system dynamics in general relativity, as a first step towards a more accurate theoretical study of orbital motion in the weak-gravity regime. For this purpose, starting from the Einstein equations in the de Donder-Lanczos gauge, this paper arrives first at the Levi-Civita Lagrangian for the geodesic motion of celestial bodies, showing in detail under which conditions the effects of internal structure and finite extension get cancelled in general relativity to first post-Newtonian order. The resulting nonlinear ordinary differential equations for the motion of planets and satellites are solved for the Earth's orbit about the Sun, written down in detail for the Sun-Earth-Moon system, and investigated for the case of planar motion of a body immersed in the gravitational field produced by the other bodies (e.g. planets with their satellites). At this stage, we prove an exact property, according to which the fourth-order time derivative of the original system leads to a linear system of ordinary differential equations. This opens an interesting perspective on forthcoming research on planetary motions in general relativity within the solar system, although the resulting equations remain a challenge for numerical and qualitative studies. Last, the evaluation of quantum corrections to location of collinear and noncollinear Lagrangian points for the planar restricted three-body problem is revisited, and a new set of theoretical values of such corrections for the Earth-Moon-planetoid system is displayed and discussed. On the side of classical values, the general relativity corrections to Newtonian values for collinear and noncollinear Lagrangian points of the Sun-Earth-planetoid system are also obtained.

PACS numbers: 04.60.Ds, 95.10.Ce

I. INTRODUCTION

Recent work by the authors [1–4], motivated, on the quantum side, by modern developments in effective field theories of gravity [5–13], and, on the classical side, by the beautiful discoveries in celestial mechanics [14–54], aerospace engineering [55–62] and (lunar) laser ranging techniques [63–75], has studied in detail the libration points of the restricted three-body problem in Newtonian gravity, general relativity and effective field theories of gravity. In

^{*}E-mail: ebattista@na.infn.it

[†]E-mail: gesposit@na.infn.it

[‡]E-mail: luciano.difiore@na.infn.it

[§]E-mail: simone.dellagnello@lnf.infn.it

^{**}E-mail: JSimo@uclan.ac.uk

^{††}E-mail: aniello.grado@gmail.com

particular, we have found (see Ref. [4] and Erratum therein) that general relativity corrects by 0.19 mm, -0.32 mm and -0.04 mm the location of the collinear Lagrangian points L_1, L_2 and L_3 respectively, for the circular restricted three-body problem where a planetoid (e.g. a satellite) moves in the gravitational field of the Earth and the Moon. Moreover, for the planar (x, y) coordinates of noncollinear Lagrangian points L_4 and L_5 , the Einstein correction to Newtonian values is 2.73 mm for x and -1.59 mm for y . The possible quantum corrections¹ to the Lagrangian points $L_1 \dots L_5$ are just below or just above 1 mm. These tiny theoretical values are conceptually interesting but, unfortunately, too small to put on firm ground the hope that one might arrive in the near future to a new test of general relativity or to a clearcut discrimination between corrections predicted by general relativity and those predicted by effective field theories of gravity.

Nevertheless, since the three-body problem is just an approximation of a much more complicated setting, where all celestial bodies in the solar system (planets, satellites, asteroids, ...) interact with each other while solar radiation pressure [76] and other sources of nongravitational perturbations may also come into play, it remains conceptually desirable to improve the current understanding of solar system dynamics in general relativity, as a first step towards a more accurate theoretical study of orbital motion, possibly including the outstanding open problem of solar system stability.

For this purpose, Sec. II performs a concise but detailed review of the N -body problem in general relativity, following the thorough analysis in Ref. [19]. First, the Einstein equations are studied in the de Donder-Lanczos gauge so as to obtain a Lagrangian for the geodesic motion of celestial bodies. Second, the physics of gravitating bodies with finite extension is studied, showing in detail under which conditions one arrives at a cancellation effect also in classical general relativity. Section III applies the Lagrangian of Sec. II to the Earth's motion around the Sun and then to the Sun-Earth-Moon system, and eventually to the planar motion of a body subject to the gravitational attraction of several bodies, when all their mutual distances are large enough that we are in the weak-gravity regime. Section IV derives in detail the linear (not linearized!) system of ordinary differential equations associated with our nonlinear equations of motion. Section V considers another possible definition of quantum corrections to Lagrangian points. Results and open problems are discussed in Sec. VI.

II. LAGRANGIAN OF THE N-BODY PROBLEM IN GENERAL RELATIVITY

In General Relativity, the N -body Lagrangian of celestial mechanics is derived by appealing to the geodesic principle for the motion of each celestial body. The first part of the analysis deals with the field equations and derives an approximate form of the metric components and of the resulting line element under suitable assumptions. The second part of the analysis relies upon three basic assumptions on the physics of gravitating systems, which make it precise what sort of cancellation principle still holds on passing from Newtonian to relativistic celestial mechanics. Hence a Lagrangian of N material points is eventually derived.

A. The Einstein equations

The Einstein equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{8\pi G_N}{c^4}T_{\mu\nu} \quad (2.1)$$

are a quasi-linear set of partial differential equations. As is well known, this means that they are linear in second-order derivatives of the metric $g_{\mu\nu}$, whereas the nonlinearity results from the squares of first derivatives of the spacetime metric. Such a quasi-linear system is not in normal form unless a suitable supplementary condition (more frequently called gauge-fixing) is imposed. In the Levi-Civita analysis we rely upon, such a condition is the so-called de Donder-Lanczos gauge. The Lanczos gauge sets to zero the action of the scalar wave operator $\square$ on spacetime coordinates, i.e.

$$\square x^\nu = - \sum_{\lambda, \mu=0}^3 g^{\lambda\mu} \Gamma^\nu_{\lambda\mu} = 0. \quad (2.2)$$

¹ These might be seen as low-energy effects in quantum gravity, more easily accessible than the elusive high-energy effects of the Planck era.

Such a set of spacetime coordinates, if they exist, were said to be isometric because Eqs. (2.2) generalize the harmonic character of Cartesian coordinates in a Euclidean metric. The de Donder formulation takes linear combinations of such equations with coefficients given by covariant components of the metric, i.e.

$$Z_\mu = \sum_{\nu=0}^3 g_{\mu\nu} \square x^\nu. \quad (2.3)$$

Now a remarkable identity relates the Ricci tensor and the symmetrized partial derivatives of the Z_μ , i.e.

$$R_{\mu\nu} - \frac{1}{2} \left(\frac{\partial Z_\mu}{\partial x^\nu} + \frac{\partial Z_\nu}{\partial x^\mu} \right) = \frac{1}{2} \sum_{\lambda,\rho=0}^3 g^{\lambda\rho} \frac{\partial^2 g_{\mu\nu}}{\partial x^\lambda \partial x^\rho} + F \left(g_{\alpha\beta}, \frac{\partial g_{\mu\nu}}{\partial x^\rho} \right). \quad (2.4)$$

The term F is of lower order in that it depends only on the metric and its first partial derivatives. In the desired isometric coordinates, for which the Z_μ vanish, the Einstein equations, jointly with the conservation equations for the energy-momentum tensor resulting from the Bianchi identity, take therefore the normal form [19] with respect to the x^0 variable.

In the following analysis, the dimensionless terms with an order of magnitude of

$$\beta^2 \equiv \frac{v^2}{c^2}, \quad \gamma \equiv \frac{G_N}{c^2} \int_S \frac{\mu}{r} dS = \frac{G_N}{c^2} \sum_{h=1}^n \int_h \frac{\mu}{r} dC_h, \quad (2.5)$$

where S is the region occupied by all bodies C_h , $h = 1, \dots, n$, and μ is the function representing² the local density, are said to be of first order and are denoted by $O(I)$. The use of the de Donder-Lanczos gauge implies that the metric components take the approximate form [19]

$$g_{00} = 1 - 2\gamma + 2\gamma^2 - 2\zeta, \quad (2.6)$$

$$g_{0i} = g_{i0} = 4\gamma_i + O(II), \quad \forall i = 1, 2, 3, \quad (2.7)$$

$$g_{ii} = -1 - 2\gamma, \quad \forall i = 1, 2, 3. \quad (2.8)$$

With this notation, γ is the Newtonian potential in a domain S of Euclidean space³ with Cartesian coordinates x^1, x^2, x^3 , and $O(II)$ is a term of second order in β^2 and γ . If $x^0 \equiv ct$, the associated dimensionless velocities are

$$\beta^i \equiv \frac{dx^i}{dx^0} \implies \beta^2 \equiv \sum_{i=1}^3 (\beta^i)^2. \quad (2.9)$$

One can therefore define the three potentials

$$\gamma_i \equiv \frac{G_N}{c^2} \int_S \frac{\mu \beta_i}{r} dS, \quad (2.10)$$

while the function ζ can be split into the sum of three functions, i.e. [19]

$$\zeta \equiv \varphi + \psi + \nu, \quad (2.11)$$

where φ and ψ solve a Poisson-type equation:

$$\varphi \equiv -\frac{G_N}{c^2} \int_S \frac{\mu \gamma}{r} dS, \quad (2.12)$$

² No confusion should arise when we write instead μ for a coordinate index in tensor equations, and similarly for other Greek letters used hereafter, whose meaning will be clear from the context.

³ It is a nontrivial property that three-dimensional space is taken to be Euclidean. From the point of view of general formalism, some authors regard this assumption as Procuste's bed [26] and hence a severe drawback, but for the analysis of planetary motions this remains a legitimate approach, whereas it would be totally inappropriate in the relativistic astrophysics of binary systems.

$$\psi \equiv \frac{3}{2} \frac{G_N}{c^2} \int_S \frac{\mu \beta^2}{r} dS, \quad (2.13)$$

whereas

$$\nu \equiv \frac{1}{2} \frac{G_N}{c^2} \frac{\partial^2}{\partial (x^0)^2} \int_S \mu r dS. \quad (2.14)$$

By virtue of (2.6)-(2.14), the squared line element takes the approximate form

$$\left(\frac{ds}{dx^0} \right)^2 = 1 - 2 \left(\frac{\beta^2}{2} + \gamma \right) + 2\gamma^2 - 2\zeta - 2\gamma\beta^2 + 8 \sum_{i=1}^3 \gamma_i \beta^i. \quad (2.15)$$

The formulation of the geodesic principle for celestial bodies needs actually the square root of (2.15), i.e.

$$\delta \int ds = \delta \int \frac{ds}{dx^0} dx^0 = 0. \quad (2.16)$$

From the standard second-order Taylor expansion of $(1 + \varepsilon)^{\frac{1}{2}}$ in the neighborhood of $\varepsilon = 0$ one gets, upon denoting by $\mathcal{N}$ the Newtonian gravitational Lagrangian

$$\mathcal{N} \equiv \frac{\beta^2}{2} + \gamma, \quad (2.17)$$

the useful approximate formula

$$\begin{aligned} \frac{ds}{dx^0} &= \sqrt{1 - 2 \left(\frac{\beta^2}{2} + \gamma \right) + 2\gamma^2 - 2\zeta - 2\gamma\beta^2 + 8 \sum_{i=1}^3 \gamma_i \beta^i} \\ &\approx 1 - \mathcal{N} + \gamma^2 - \zeta - \gamma\beta^2 + 4 \sum_{i=1}^3 \gamma_i \beta^i - \frac{1}{2} \mathcal{N}^2. \end{aligned} \quad (2.18)$$

The approximate calculation leading from (2.15) to (2.18) is simple, but it represents a crucial conceptual step. A rigorous analysis of stability of the solar system would require working with the square root on the first line of Eq. (2.18) without any expansion, and then using the modern qualitative methods of the calculus of variations. The constant 1 is of course inessential, and this is made precise by pointing out that also the following variation vanishes [77]:

$$\delta \int dx^0 = 0, \quad (2.19)$$

because we can avoid letting x^0 to vary since the left-hand side of the equation expressing the geodesic principle undergoes a variation that vanishes by virtue of the conditions resulting from variation of the three coordinates x^1, x^2, x^3 . In light of (2.16) and (2.19), we get eventually the dimensionless Lagrangian

$$\mathcal{L} = 1 - \frac{ds}{dx^0} = \mathcal{N} + \mathcal{D}, \quad (2.20)$$

where $\mathcal{N}$, defined in (2.17), is the Lagrangian of a dimensionless material element in Newtonian mechanics, whereas $\mathcal{D}$ is the Einstein modification of this Newtonian Lagrangian, and is given by [19]

$$\mathcal{D} \equiv \frac{1}{2} \mathcal{N}^2 - \gamma^2 + \zeta + \gamma\beta^2 - 4 \sum_{i=1}^3 \gamma_i \beta^i. \quad (2.21)$$

In the course of this first part of the analysis one assumes that the gradient of pressure vanishes at the center of gravity of each body, which implies that the motion remains geodesic and hence it is legitimate to limit ourselves to the consideration of unbundled media [19].

B. Physics of gravitating bodies with finite extension

The Einstein modification $\mathcal{D}$ in (2.21) contains the functions γ, γ_i, ζ obtained by integrating over all bodies, and hence is related to finite size and internal structure of such bodies, that we will later identify with Sun, Earth, Moon, all planets with their satellites, and a mechanical satellite sent off from the Earth. Following Ref. [19], for each body we can think of the domain S as the disjoint union of C with S' , where C denotes the domain occupied by the body, while S' is the residual portion of S . We shall use hereafter the notation according to which

$$\gamma' \equiv \gamma|_{S'}, \quad \gamma'' \equiv \gamma|_C, \quad (2.22)$$

$$\zeta' \equiv \zeta|_{S'}, \quad \zeta'' \equiv \zeta|_C, \quad (2.23)$$

so that we can write the decompositions

$$\gamma = \gamma' + \gamma'', \quad \zeta = \zeta' + \zeta'', \quad (2.24)$$

and also, for the two parts (Newton and Einstein) of the full Lagrangian, the resulting splits

$$\mathcal{N} = \mathcal{N}' + \mathcal{N}'', \quad \mathcal{D} = \mathcal{D}' + \mathcal{D}'', \quad (2.25)$$

where $\mathcal{N}', \mathcal{D}'$ are the part that we would have if the body C were suppressed, whereas $\mathcal{N}'', \mathcal{D}''$ characterize the influence of the body C on the motion of a point $P \in C$. The explicit formulas we need are

$$\mathcal{N}' = \frac{1}{2}\beta^2 + \gamma', \quad \mathcal{N}'' = \gamma'', \quad (2.26)$$

$$\mathcal{D}' = \frac{1}{2}\mathcal{N}'^2 - \gamma'^2 + \zeta' + \gamma'\beta^2 - 4 \sum_{i=1}^3 \gamma'_i \beta^i, \quad (2.27)$$

$$\mathcal{D}'' = \mathcal{N}'\gamma'' - \frac{1}{2}\gamma''^2 - 2\gamma'\gamma'' + \varphi'' + \psi'' + \nu'' + \gamma''\beta^2 - 4 \sum_{i=1}^3 \gamma''_i \beta^i. \quad (2.28)$$

Indeed, the internal forces resulting from γ'' are in general stronger than the external forces resulting from the potential γ' . Nevertheless, in the equation ruling the motion of the center of gravity, the contributions of derivatives of γ'' cancel pairwise exactly. It is here that resides, conceptually, the origin of the cancellation principle in classical mechanics [19].

C. Center of gravity, quasi-translational motions, size vs. distance

At this stage, since the mere use of Eqs. (2.22)-(2.28) would lead to unmanageable equations and would not shed enough light on the N-body problem in General Relativity, some further physical assumptions come into play that are indeed satisfied *approximately* by all known celestial bodies in the solar system. They are as follows [19].

A1 The center of gravity P_G of each body C is *substantial*, i.e., it always adheres to the same material element. This implies that its motion will be characterized, as it occurs for any other material point P , by a Lagrangian $\mathcal{L} = \mathcal{N} + \mathcal{D}$, with the Einstein perturbation being given by the term $\mathcal{D}$. Furthermore, we shall assume that the center of gravity P_G is always a center of gravitation. The latter condition means that P_G is a point where the Newtonian attractions of material elements of the body⁴ (i.e., the internal forces) add up to zero. In other words, P_G coincides with the mass center of the body.

⁴ Recall that the Newtonian potential γ is bounded everywhere and it vanishes at infinity, hence there exists a point of minimum for γ at which its gradient vanishes, which means that the force vanishes at this point.

A2 The body performs a quasi-translational motion. Indeed, in a translational motion, all points of the body have, at any instant t , the same vector speed, e.g., the speed $\vec{v}_g$ of the center of gravity. We can still regard as a translation every motion for which, defining

$$|\Delta \vec{v}(t)| \equiv |\vec{v}_{P_i}(t) - \vec{v}_{P_j}(t)| \quad \forall P_i, P_j \in C, \quad (2.29)$$

one has always

$$\frac{|\Delta \vec{v}|}{|\vec{v}_g|} \ll 1. \quad (2.30)$$

We need sufficiently small values of the ratio in (2.30), e.g., of order 10^{-2} , so that one can neglect, as a quantity of order greater than 1, every product of the type

$$\beta^2 \frac{|\Delta \vec{v}|}{|\vec{v}_g|}, \quad \gamma \frac{|\Delta \vec{v}|}{|\vec{v}_g|}.$$

This is precisely what happens for planetary motions. Their deformations are initially negligible and they behave, as a consequence, as essentially rigid bodies. Their motion is actually a composition of translation and rotation. However, for every point of the body, the speed resulting from rotation attains only a few percent of the common speed of translation. For example, in the case of the Earth, one has

$$\frac{|\Delta \vec{v}|}{|\vec{v}_g|} \approx 3 \cdot 10^{-2}.$$

A3 On denoting by d the maximal size of the body C , and by R the minimal Euclidean distance d_E between points of C and points of the residual portion of S' :

$$R \equiv \min d_E(P_j(C), P_k(S')), \quad (2.31)$$

the quantity $\left(\frac{d}{R}\right)^2$ is negligible. For the Sun-Earth system, one has indeed $\left(\frac{d}{R}\right)^2 \approx 10^{-4}$.

By virtue of **A1**, the gradient of the potential γ'' vanishes at the center of gravity, and hence γ'' behaves as a constant. Moreover, by virtue of **A2**, the dimensionless velocities β^i defined in (2.9) are constant within the body C . On defining

$$\gamma'' \equiv \tilde{\omega} = \text{constant}, \quad (2.32)$$

one finds from (2.10)-(2.14)

$$-\varphi'' = \tilde{\omega} \gamma'' = \tilde{\omega}^2, \quad (2.33)$$

$$\psi'' = \frac{3}{2} \beta^2 \gamma'' = \frac{3}{2} \tilde{\omega} \beta^2, \quad (2.34)$$

$$\gamma''_i = \beta^i \gamma'' = \tilde{\omega} \beta^i, \quad (2.35)$$

$$\nu'' = \frac{1}{2} \frac{G_N}{c^2} \frac{\partial^2}{\partial (x^0)^2} \int_C \mu r dC = 0, \quad r = d_E(P_j, P_G) \quad \forall P_j \in C. \quad (2.36)$$

We stress that ν'' vanishes because the integral in (2.36) is constant during the motion.

D. The χ_k coefficients

Hereafter we denote with P_k the center of gravity of the body C_k , for all $k = 0, 1, \dots, n-1$. We also denote with l_k the gravitational radius of the k -th body having mass m_k (we assume, on experimental ground, the equality of inertial and gravitational mass, and also of active and passive gravitational mass), i.e.,

$$l_k \equiv \frac{G_N m_k}{c^2}, \quad (2.37)$$

which, in the solar system, does not exceed the 1.5 km for the gravitational radius of the Sun. The assumption **A3** implies that the potential of the body C_k acting on the center of gravity P_h of C_h is given, as if the mass of C_k were completely concentrated at the center of gravity P_k , in the form $\frac{G_N m_k}{r(P_k, P_h)}$, where the denominator is the Euclidean distance $d_E(P_k, P_h)$. The dimensionless form of such a potential is obtained dividing by c^2 , i.e.

$$\frac{G_N m_k}{c^2 r(P_k, P_h)} = \frac{l_k}{r(P_k, P_h)}.$$

Now we need extra labels in the notation, since we are going to derive the Lagrangian in the form $\mathcal{N} + \mathcal{D}$ for each (celestial) body. For this purpose, following again Ref. [19], we shall denote by β_h^2 the square of the velocity of P_h , by $(\beta_h)_i$ the component along the axis x^i of β_h , and by γ'_h the potential at P_h resulting from all bodies C_k described by an index $k \neq h$. This latter condition means that

$$\gamma'_h = \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{l_k}{r(P_k, P_h)}, \quad (2.38)$$

with the understanding that the Kronecker δ plays the role of giving vanishing weight to the divergent term $\frac{l_k}{r(P_k, P_k)}$, which is therefore ruled out from the sum (sometimes this is expressed by the $\sum'$ notation, which is here made clearer).

In light of (2.32), we can regard as being constant the integral

$$\gamma''_h(P) = \frac{G_N}{c^2} \int_{C_h} \frac{\mu(Q)}{r(Q, P)} dC_h = \tilde{\omega}_h, \quad (2.39)$$

which is the potential of the body C_h at any point P of C_h itself. The constant $\tilde{\omega}_h$ is majorized by the ratio $\frac{\bar{l}}{\bar{r}}$, where (∂C_h) being the boundary of C_h

$$\bar{r} \equiv \max d_E(P_h, \partial C_h), \quad (2.40)$$

while $\bar{l} \equiv \frac{G_N \bar{m}}{c^2}$, $\bar{m}$ being the mass contained within a homogeneous sphere of density $\bar{\mu}$ and radius $\bar{r}$, having set

$$\bar{\mu} = \sup(\mu) \text{ within } C_h. \quad (2.41)$$

Equations (2.39)-(2.41) tell us that $\tilde{\omega}$ is a quantity of first order, being close to a quantity proportional to a Newtonian potential. This simple property will be nicely exploited below.

Next, we consider an integration domain S' consisting of all bodies C_k with the exception of the body C_h , in formulas

$$S' \equiv \cup_k C_k - C_h, \quad (2.42)$$

and, for points $Q \in S'$ and $P \in C_h$, we consider the decomposition of the function φ_h for the body C_h in the form

$$\varphi_h = \varphi'_h + \varphi''_h, \quad (2.43)$$

where

$$\varphi'_h = -\frac{G_N}{c^2} \int_{S'} \frac{\mu \gamma}{r(Q, P)} dS'. \quad (2.44)$$

In light of assumption **A3**, we can re-express (2.44) in the form

$$\begin{aligned} \varphi'_h &= -\frac{G_N}{c^2} \sum_{k=0}^{n-1} \frac{(1 - \delta_{k,h})}{r(P_k, P)} \int_{C_k} \mu(Q) \gamma_k(Q) dC_k \\ &= -\frac{G_N}{c^2} \sum_{k=0}^{n-1} \frac{(1 - \delta_{k,h})}{r(P_k, P)} \left(\sum_{j=0}^{n-1} (1 - \delta_{j,k}) \frac{l_j}{r(P_j, P_k)} \int_{C_k} \mu(Q) dC_k + \frac{G_N}{c^2} \int_{C_k} \mu(Q) dC_k \int_{C_k} \frac{\mu(Q')}{r(Q, Q')} dC_k \right) \\ &= -\sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{l_k}{r(P_k, P)} \sum_{j=0}^{n-1} (1 - \delta_{j,k}) \frac{l_j}{r(P_j, P_k)} - \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{l_k \chi_k}{r(P_k, P)}, \end{aligned} \quad (2.45)$$

having defined [19]

$$\chi_k \equiv \frac{1}{l_k} \left(\frac{G_N}{c^2} \right)^2 \int_{C_k} \mu(Q) dC_k \int_{C_k} \frac{\mu(Q')}{r(Q', Q)} dC_k = \frac{G_N}{c^2} \tilde{\omega}_k = \text{constant}, \quad (2.46)$$

and where each γ_k potential in (2.45) has been split as in (2.24), i.e., $\gamma'_k(Q) + \gamma''_k(Q)$, and use of (2.38) and (2.39) has been made in order to express $\gamma'_k(Q)$ and $\gamma''_k(Q)$, respectively. The existence of the constant coefficients χ_k is conceptually interesting, but their values are extremely small, because the integrals occurring in (2.46) are finite but are multiplied by the square of the ratio $\frac{G_N}{c^2}$.

E. The effacement property

At this stage, one can obtain the desired decomposition of the Lagrangian for the h -th body C_h in the form [19]

$$\mathcal{L}_h = \mathcal{N}_h + \mathcal{D}'_h + \mathcal{D}''_h, \quad (2.47)$$

where $\mathcal{N}_h$ is the Newtonian term

$$\mathcal{N}_h = \frac{1}{2}(\beta_h)^2 + \gamma'_h = \mathcal{N}'_h, \quad (2.48)$$

while $\mathcal{D}'_h$ is the pointlike Einstein perturbation

$$\mathcal{D}'_h = \frac{1}{2}(\mathcal{N}'_h)^2 - (\gamma'_h)^2 + \zeta'_h + \gamma'_h(\beta_h)^2 - 4 \sum_{i=1}^3 (\Gamma'_h)_i (\beta_h)_i, \quad (2.49)$$

$(\beta_h)_i$ denoting the i -th component of the velocity of the h -th body as we said before (2.38), and

$$(\Gamma'_h)_i \equiv \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{l_k (\beta_k)_i}{r(P_k, P_h)}. \quad (2.50)$$

The expression (2.47) of the Lagrangian is completed by $\mathcal{D}''_h$, i.e. the Einstein perturbation resulting from the extension of bodies. Upon defining

$$m \equiv \sum_{k=0}^{n-1} m_k, \quad l \equiv \frac{G_N m}{c^2}, \quad l_k \equiv \frac{G_N m_k}{c^2}, \quad \lambda_k \equiv \frac{m_k}{m} = \frac{l_k}{l}, \quad (2.51)$$

the perturbation $\mathcal{D}''_h$ reads as

$$\mathcal{D}''_h = -\tilde{\omega}_h (\beta_h)^2 - l \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{\lambda_k (\chi_k + 2\tilde{\omega}_h)}{r(P_k, P_h)}. \quad (2.52)$$

We can now recall that Lagrangians differing by a multiplicative constant give rise to equivalent equations of motion. The simple and profound idea of Levi Civita was to consider a first-order quantity σ_h , and to multiply $\mathcal{L}_h$ by $(1 + \sigma_h)$. After doing this, one can try to choose σ_h in such a way that the occurrence of the constant $\tilde{\omega}_h$ gets exactly cancelled. This is indeed feasible because, up to higher order terms here negligible, one finds

$$(1 + \sigma_h) \mathcal{L}_h \sim (1 + \sigma_h) \mathcal{N}_h + \mathcal{D}'_h + \mathcal{D}''_h, \quad (2.53)$$

where, in particular,

$$(1 + \sigma_h) \mathcal{N}_h + \mathcal{D}''_h = \frac{1}{2}(\beta_h)^2 + \left(\frac{1}{2} \sigma_h - \tilde{\omega}_h \right) (\beta_h)^2 + l \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{\lambda_k [(1 - \chi_k) + (\sigma_h - 2\tilde{\omega}_h)]}{r(P_k, P_h)}. \quad (2.54)$$

This formula suggests choosing

$$\sigma_h = 2\tilde{\omega}_h, \quad (2.55)$$

to achieve the desired cancellation. The result is also consistent with what we know already about the first-order nature of the constant $\tilde{\omega}_h$. We can further define

$$\Lambda_k \equiv \lambda_k(1 - \chi_k), \quad (2.56)$$

Note also that the pointlike Einstein perturbation $\mathcal{D}'_h$ is still expressed in terms of the dimensionless λ_k coefficients, but we can insert also therein the Λ_k defined in (2.56), because

$$\mathcal{D}'_h(\lambda_k) = \mathcal{D}'_h(\Lambda_k) + \text{higher-order terms}. \quad (2.57)$$

One finds therefore that each (celestial) body is ruled by a pointlike Lagrangian L_h where the Einstein perturbation is no longer split into pointlike plus finite-size part, and one can write [19]

$$L_h = \mathcal{N}_h + \mathcal{D}_h, \quad (2.58)$$

where, having defined

$$\gamma_h \equiv l \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{\Lambda_k}{r(P_k, P_h)}, \quad (2.59)$$

$$(\Gamma_h)_i \equiv l \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{\Lambda_k(\beta_k)_i}{r(P_k, P_h)}, \quad (2.60)$$

$$\zeta_h \equiv \varphi_h + \psi_h + \nu_h, \quad (2.61)$$

$$\varphi_h \equiv -l^2 \sum_{k=0}^{n-1} \left[(1 - \delta_{k,h}) \frac{\Lambda_k}{r(P_k, P_h)} \sum_{s=0}^{n-1} (1 - \delta_{s,k}) \frac{\Lambda_s}{r(P_s, P_k)} \right], \quad (2.62)$$

$$\psi_h \equiv \frac{3}{2} l \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \frac{\Lambda_k(\beta_k)^2}{r(P_k, P_h)}, \quad (2.63)$$

$$\nu_h \equiv \frac{1}{2} l \frac{\partial^2}{\partial(x^0)^2} \sum_{k=0}^{n-1} (1 - \delta_{k,h}) \Lambda_k r(P_k, P_h), \quad (2.64)$$

the Newtonian term takes the familiar form

$$\mathcal{N}_h \equiv \frac{1}{2} (\beta_h)^2 + \gamma_h, \quad (2.65)$$

while the Einstein perturbation is eventually expressed by the sum of functions

$$\mathcal{D}_h \equiv \frac{1}{2} (\mathcal{N}_h)^2 - (\gamma_h)^2 + \zeta_h + \gamma_h (\beta_h)^2 - 4 \sum_{i=1}^3 (\Gamma_h)_i (\beta_h)_i. \quad (2.66)$$

Equations (2.58)-(2.66) lead to an accurate scheme for writing down and studying the equations of motion of each (celestial) body, and provide a precise statement of the cancellation principle in General Relativity: on going from Newtonian to relativistic celestial mechanics, the effects of extension and internal structure of bodies are encoded in the family of Λ_k parameters, Eq. (2.56), which differ only by a tiny amount (see Eq. (2.46)) from the dimensionless mass ratios $\frac{m_k}{m}$ of Eq. (2.51). Thus, the effects of finite extension of bodies get eventually dissolved neatly, and it is as if we were dealing with material points which do not affect at all their center of gravity. This holds also for the solar system.

As is stressed in Ref. [27], a more rigorous proof of such a cancellation principle can be found in Ref. [18], to which we refer the reader interested in a broader understanding.

III. SUN-EARTH, SUN-EARTH-MOON AND N -BODY DYNAMICS

Driven by the concepts outlined in the previous sections, we now aim at investigating the system consisting of the Sun, Earth and as many additional celestial bodies as possible by means of the Lagrangian (2.58). An important comment should be made at this stage. In fact, in the most general case, acceleration terms appear in (2.58). However, by bearing in mind that $\chi_k \ll 1$ and $\lambda_k = \text{constant}$ (cf. Eqs. (2.46) and (2.51)), it easily follows that

$$\nu_h = 0, \quad \forall h, \quad (3.1)$$

since the Euclidean distance $r(P_k, P_h)$ occurring in Eq. (2.64) depends on time only implicitly, through the coordinates $x^i(t)$, i.e.,

$$\frac{\partial}{\partial t} r(P_k, P_h) = \frac{\partial}{\partial t} r(x^1(t), x^2(t), x^3(t)) = 0. \quad (3.2)$$

In other words, under our assumption the Lagrangian (2.58) turns out to be a function of the Euclidean coordinates $x^1(t), x^2(t), x^3(t)$ and their first time derivatives only. Moreover, Eq. (3.1) is valid also in the more general case of a Lagrangian function depending explicitly on the time variable t , since the value of the integral

$$\int_D \mu r \, dC, \quad (3.3)$$

evaluated in the region D made up of all those spatial points which are very distant from P_h reduces nearly to zero because it turns out to be of order $\left(\frac{d}{R}\right)^2$, which, according to hypothesis **A3**, represents a negligible quantity.

As a first step, we have analysed the system made up of just two bodies by considering the case of one celestial body orbiting a fixed massive object (i.e., the Sun). We have found that perihelion shift predicted by the Levi-Civita Lagrangian is in accordance with the well-known results expected within the usual 1PN picture of general relativity. In fact by employing the latter approximation, the orbit of the revolving body (in the equatorial plane $\theta = \pi/2$) is described by the well-known relations [78]

$$r(\phi) = \frac{(1 - e^2)a}{1 + e \cos[(1 - \delta\phi_0/2\pi)\phi]}, \quad (3.4)$$

and

$$\begin{aligned} x &= e a + r(\phi) \cos \phi, \\ y &= r(\phi) \sin \phi, \end{aligned} \quad (3.5)$$

with

$$\delta\phi_0 = \frac{6\pi G_N M}{c^2 a (1 - e^2)}, \quad (3.6)$$

e being the eccentricity of the orbit of the orbiting body, a the semi-major axis and M the mass of the massive object. As demonstrated in Figs. 1 and 2, the outcomes achieved within the Levi-Civita framework are in agreement with the ones expected through the 1PN approximation method, witnessing that Levi-Civita actually made a mistake in Ref. [19] when concluding that his pattern predicts a more pronounced shift of the perihelion in the orbit of the revolving body. This point is in accordance with the analysis of Ref. [27].

Moreover, the Newtonian relations expressing the eccentricity e of the orbit as a function of the initial velocity v_0 of the revolving body and its distance $\mathcal{R}$ from the massive one according to

$$\begin{aligned} v_0 &= \sqrt{\frac{G_N M}{\mathcal{R}}}, && \text{circular orbit,} \\ 0 < v_0 < \sqrt{\frac{2G_N M}{\mathcal{R}}}, && \left(v_0 \neq \sqrt{\frac{G_N M}{\mathcal{R}}} \right), \quad \text{elliptic orbit,} \\ v_0 &= \sqrt{\frac{2G_N M}{\mathcal{R}}}, && \text{parabolic orbit,} \\ v_0 &> \sqrt{\frac{2G_N M}{\mathcal{R}}}, && \text{hyperbolic orbit,} \end{aligned} \quad (3.7)$$




FIG. 1: Circular orbit around the Sun of a body having the mass of the Earth as described by the Lagrangian function (2.58). The same orbit is obtained also when Eqs. (3.4)–(3.6) are employed.




FIG. 2: Elliptic orbit around the Sun of a body having the mass of the Earth as described by the Lagrangian function (2.58). The same orbit is obtained also when Eqs. (3.4)–(3.6) are employed.

are found to be respected by applying the Levi-Civita framework.

After that, the system consisting of three bodies (i.e., the Earth and the Moon orbiting the Sun) has been considered. We have recovered the orbit of the Moon around the Earth and around the Sun. Once again, the Levi-Civita Lagrangian produces negligible differences with respect to both the 1PN model and Newtonian theory. As an example, Fig. 3 describes the motion around the Sun (and in the presence of the Earth describing the usual Newtonian elliptic orbit around the Sun) of a body having the same mass as the Moon in the hypothesis of circular motion. Also the cases of elliptic, parabolic, and hyperbolic orbits have been analysed.

The presence in our model of such periodic solutions will become crucial at the end of Sec. IV (see also appendix B and the remarks therein on the Sturm-Liouville problem).

Many complications arise when we deal with the Sun-Jupiter-Earth-Moon-planetoid system so that some further simplification must be assumed for the Lagrangian function (2.58). If we suppose that the velocity β_k of the Sun, Earth, Moon, and Jupiter is negligible if compared to the planetoid velocity, we can then set (cf. Eqs. (2.60) and (2.63))

$$\begin{aligned} (\Gamma_h)_i &= 0, \\ \psi_h &= 0, \end{aligned} \tag{3.8}$$

where the index h must be understood as labelling the planetoid. In this way the resulting Lagrangian is such that

$$\mathcal{D}_h = \frac{1}{2}(\mathcal{N}_h)^2 - (\gamma_h)^2 + \varphi_h + \gamma_h(\beta_h)^2, \tag{3.9}$$

and hence it assumes a form that can be studied more easily.




FIG. 3: Circular orbit around the Sun in the gravitational field produced by the Sun itself and the Earth of a body having the mass of the Moon as described by the Levi-Civita Lagrangian (2.58). Note that this Figure is similar to Fig. 1 since the distances of the Earth and the Moon from the Sun are quite the same.

A. Sun-Earth system

In the simple system made up of just the Earth orbiting the Sun, the joint effect of Eqs. (2.58) and (3.9) gives rise to a Lagrangian function describing the planar motion of the Earth

$$m_E c^2 \mathcal{L}_E(x, y, \dot{x}, \dot{y}) = m_E c^2 (\mathcal{N}_E + \mathcal{D}_E), \quad (3.10)$$

(m_E being the mass of the Earth) such that

$$\mathcal{N}_E = \frac{1}{2} (\beta_E)^2 + \gamma_E = \frac{1}{2c^2} (\dot{x}^2 + \dot{y}^2) + \frac{Gm_S}{c^2} \frac{1}{r}, \quad (3.11)$$

$$\mathcal{D}_E = \frac{1}{2} (\mathcal{N}_E)^2 - (\gamma_E)^2 + \varphi_E + \gamma_E (\beta_E)^2, \quad (3.12)$$

with (cf. (2.62))

$$\varphi_E = -\frac{l^2}{r^2} \Lambda_S \Lambda_E, \quad (3.13)$$

where m_S is the mass of the Sun and

$$r = r(t) = \sqrt{x^2(t) + y^2(t)}, \quad (3.14)$$

denoting the Euclidean planar distance between the point $\mathbf{r}(t) = (x(t), y(t))$ occupied at the time t by the Earth and the Sun, supposed at rest at the origin of the coordinate system. The resulting Euler-Lagrange equations (which, once integrated numerically, have led to Figs. 1 and 2) can be written in the form

$$\begin{cases} f(x, y, \dot{x}, \dot{y}) \ddot{x} + g(\dot{x}, \dot{y}) \ddot{y} + h_1(x, y, \dot{x}, \dot{y}) = 0, \\ g(\dot{x}, \dot{y}) \ddot{x} + f(x, y, \frac{\dot{x}}{\sqrt{3}}, \sqrt{3}\dot{y}) \ddot{y} + h_2(x, y, \dot{x}, \dot{y}) = 0, \end{cases} \quad (3.15)$$

where

$$f(x, y, \dot{x}, \dot{y}) = \frac{1}{2} m_E \left(2 + \frac{6l\Lambda_S}{r} + \frac{3\dot{x}^2 + \dot{y}^2}{c^2} \right), \quad (3.16)$$

$$g(\dot{x}, \dot{y}) = \frac{m_E}{c^2} \dot{x} \dot{y}, \quad (3.17)$$

$$h_1(x, y, \dot{x}, \dot{y}) = \frac{lm_E \Lambda_S [-4c^2 l \Lambda_E r x - 2c^2 l \Lambda_S r x + r^2 (-6y \dot{x} \dot{y} + 2c^2 x - 3x \dot{x}^2 + 3x \dot{y}^2)]}{2r^5}, \quad (3.18)$$

$$h_2(x, y, \dot{x}, \dot{y}) = \frac{lm_E \Lambda_S [-4c^2 l \Lambda_E r y - 2c^2 l \Lambda_S r y + r^2 (-6x \dot{x} \dot{y} + 2c^2 y + 3x \dot{x}^2 - 3y \dot{y}^2)]}{2r^5}. \quad (3.19)$$

Note that it is possible to obtain $h_2(x, y, \dot{x}, \dot{y})$ starting from $h_1(x, y, \dot{x}, \dot{y})$ (or vice-versa) by setting

$$\begin{aligned} x &\longleftrightarrow y, \\ \dot{x}^2 &\rightarrow -\dot{x}^2, \\ \dot{y}^2 &\rightarrow -\dot{y}^2, \end{aligned} \quad (3.20)$$

whereas the terms linear in the velocities have the freedom to change according to

$$\begin{aligned} \dot{x} &\rightarrow \pm \dot{x}, \\ \dot{y} &\rightarrow \pm \dot{y}. \end{aligned} \quad (3.21)$$

B. Sun-Earth-Moon system

If we analyse the Sun-Earth-Moon system, the Lagrangian for the Moon becomes

$$m_M c^2 \mathcal{L}_M(x, y, \dot{x}, \dot{y}) = m_M c^2 (\mathcal{N}_M + \mathcal{D}_M), \quad (3.22)$$

m_M being the mass of the Moon, while

$$\gamma_M = \frac{l_S}{r} + \frac{l_E}{r_{EM}}, \quad (3.23)$$

$$\varphi_M = -l^2 \left[\frac{\Lambda_S}{r} \left(\frac{\Lambda_E}{r_E} + \frac{\Lambda_M}{r} \right) + \frac{\Lambda_E}{r_{EM}} \left(\frac{\Lambda_M}{r_{EM}} + \frac{\Lambda_S}{r_E} \right) \right], \quad (3.24)$$

where now $r = r(t) = \sqrt{x^2(t) + y^2(t)}$ denotes the distance of the Moon from the Sun, $r_{EM} = |\mathbf{r} - \mathbf{r}_E| = \sqrt{(x_E(t) - x(t))^2 + (y_E(t) - y(t))^2}$ represents the distance between the Earth, having coordinates $\mathbf{r}_E(t) = (x_E(t), y_E(t))$, and the Moon, whereas r_E is the distance between the Sun and the Earth. The resulting Euler-Lagrange equations, giving rise to Fig. 3, assume the form

$$\begin{cases} \tilde{f}(x, y, x_E, y_E, \dot{x}, \dot{y}) \ddot{x} + \tilde{g}(\dot{x}, \dot{y}) \ddot{y} + \tilde{h}_1(x, y, x_E, y_E, \dot{x}, \dot{y}) = 0, \\ \tilde{g}(\dot{x}, \dot{y}) \ddot{x} + \tilde{f}(x, y, x_E, y_E, \frac{\dot{x}}{\sqrt{3}}, \sqrt{3}\dot{y}) \ddot{y} + \tilde{h}_2(x, y, x_E, y_E, \dot{x}, \dot{y}) = 0, \end{cases} \quad (3.25)$$

with

$$\tilde{f}(x, y, x_E, y_E, \dot{x}, \dot{y}) = \frac{m_M}{2c^2} \left(2c^2 + \frac{6c^2 l_S}{r} + \frac{6c^2 l_E}{r_{EM}} + 3\dot{x}^2 + \dot{y}^2 \right), \quad (3.26)$$

$$\tilde{g}(\dot{x}, \dot{y}) = \frac{m_M \dot{x} \dot{y}}{c^2}, \quad (3.27)$$

$$\begin{aligned}
\tilde{h}_1(x, y, x_E, y_E, \dot{x}, \dot{y}) = & \frac{1}{2} m_M \left\{ \frac{1}{r_{EM}^3} \left(\frac{2c^2 l_E^2 x_E}{r_{EM}} + \frac{4c^2 l^2 \Lambda_E \Lambda_M x_E}{r_{EM}} + 2c^2 l_E x_E + 2c^2 l^2 \Lambda_E \Lambda_S (x_E/r_E) \right) \right. \\
& + \frac{2c^2 l_E l_S x_E}{r} + 3l_E x_E \dot{x}^2 + 6l_E y_E \dot{x} \dot{y} - 3l_E x_E \dot{y}^2 \Big) + 6y \dot{x} \left[\frac{l_E \omega_E x_E}{r_{EM}^3} - \dot{y} \left(\frac{l_S}{r^3} + \frac{l_E}{r_{EM}^3} \right) \right] \\
& + x \left[2c^2 \left(-\frac{l_S^2}{r^4} - \frac{2l^2 \Lambda_M \Lambda_S}{r^4} - \frac{l^2 \Lambda_E \Lambda_S}{r_E r^3} - \frac{l_E^2}{r_{EM}^4} - \frac{2l^2 \Lambda_E \Lambda_M}{r_{EM}^4} + \frac{l_E}{r_{EM}^3} - \frac{l^2 \Lambda_E \Lambda_S}{r_E r_{EM}^3} \right) \right. \\
& \left. \left. + \frac{l_S}{r^3} + \frac{l_S l_E}{r r_{EM}^3} - \frac{l_S l_E}{r^3 r_{EM}} \right) - \frac{6l_E \omega_E y \dot{x}}{r_{EM}^3} + 3(\dot{y}^2 - \dot{x}^2) \left(\frac{l_S}{r^3} + \frac{l_E}{r_{EM}^3} \right) \right] \Big\}, \tag{3.28}
\end{aligned}$$

ω_E being the classical pulsation associated to the motion of the Earth around the Sun and, likewise the two-body case,

$$\tilde{h}_1(x, y, x_E, y_E, \dot{x}, \dot{y}) \longleftrightarrow \tilde{h}_2(x, y, x_E, y_E, \dot{x}, \dot{y}), \tag{3.29}$$

if

$$\begin{aligned}
x & \longleftrightarrow y, \\
x_E & \longleftrightarrow y_E, \\
\dot{x} & \rightarrow -\dot{x}, \\
\dot{y} & \rightarrow -\dot{y}, \\
\dot{x}^2 & \rightarrow -\dot{x}^2, \\
\dot{y}^2 & \rightarrow -\dot{y}^2.
\end{aligned} \tag{3.30}$$

C. Case of the h -th body immersed in the gravitational field produced by the other $(n-1)$ bodies

Driven by the features of the above analysis, it is possible to infer the presence of a recursive scheme according to which the Lagrange equations resulting from Eqs. (2.58) and (3.9), and describing the motion of the h -th body immersed in the gravitational field produced by the other $n-1$ bodies, are given by

$$\begin{cases} \mathcal{F}(x_h, y_h, x_k, y_k, \dot{x}_h, \dot{y}_h) \ddot{x}_h + \mathcal{G}(\dot{x}_h, \dot{y}_h) \ddot{y}_h + \mathcal{H}_1(x_h, y_h, x_k, y_k, \dot{x}_h, \dot{y}_h) = 0, \\ \mathcal{G}(\dot{x}_h, \dot{y}_h) \ddot{x}_h + \mathcal{F}(x_h, y_h, x_k, y_k, \frac{\dot{x}_h}{\sqrt{3}}, \sqrt{3} \dot{y}_h) \ddot{y}_h + \mathcal{H}_2(x_h, y_h, x_k, y_k, \dot{x}_h, \dot{y}_h) = 0, \end{cases} \quad (k = 0, 1, \dots, n-1), (k \neq h), \tag{3.31}$$

possessing the property

$$\mathcal{H}_1(x_h, y_h, x_k, y_k, \dot{x}_h, \dot{y}_h) \longleftrightarrow \mathcal{H}_2(x_h, y_h, x_k, y_k, \dot{x}_h, \dot{y}_h), \tag{3.32}$$

when

$$\begin{aligned}
x_h & \longleftrightarrow y_h, \\
x_k & \longleftrightarrow y_k, \\
\dot{x}_h & \rightarrow -\dot{x}_h, \\
\dot{y}_h & \rightarrow -\dot{y}_h, \\
\dot{x}_h^2 & \rightarrow -\dot{x}_h^2, \\
\dot{y}_h^2 & \rightarrow -\dot{y}_h^2.
\end{aligned} \tag{3.33}$$

IV. LINEAR SYSTEM OF ORDINARY DIFFERENTIAL EQUATIONS ASSOCIATED WITH THE NONLINEAR EQUATIONS OF MOTION

The expressions describing the relativistic motion of a massive object we have derived in the previous section (cf. Eqs. (3.15), (3.25), and (3.31)) clearly represent a coupled second-order system of nonlinear ordinary differential

equations. All the coefficients occurring in these equations can be seen as smooth and differentiable functions on $\mathbb{R}^2$ since the distance scales occurring in our framework prevent the bodies from colliding. This means that our model cannot be employed to investigate the binary systems analysed in relativistic astrophysics [80–89], or those systems made up of comets or asteroids hitting a planet or other celestial bodies.

We now aim at showing that it is possible to map such systems into a second-order system of *linear* ordinary differential equations by applying the time derivative operator $\frac{d}{dt}$ four times to the original equations. The resulting expressions contain the fourth-order time derivative of the functions $x(t)$ and $y(t)$ as their unknowns. Consider the system (3.31) written in matrix form

$$\mathcal{A} \begin{bmatrix} \ddot{x}_h \\ \ddot{y}_h \end{bmatrix} + \begin{bmatrix} \mathcal{H}_1(x_h, y_h, x_k, y_k, \dot{x}_h, \dot{y}_h) \\ \mathcal{H}_2(x_h, y_h, x_k, y_k, \dot{x}_h, \dot{y}_h) \end{bmatrix} = 0, \quad (4.1)$$

where

$$\mathcal{A} = \begin{bmatrix} \mathcal{F}(x_h, y_h, x_k, y_k, \dot{x}_h, \dot{y}_h) & \mathcal{G}(\dot{x}_h, \dot{y}_h) \\ \mathcal{G}(\dot{x}_h, \dot{y}_h) & \mathcal{F}(x_h, y_h, x_k, y_k, \frac{\dot{x}_h}{\sqrt{3}}, \sqrt{3}\dot{y}_h) \end{bmatrix}. \quad (4.2)$$

Thus, we can write Eq. (3.31) as (hereafter the convention on the summation over repeated indices is employed)

$$\mathcal{A}_{ir}(x_j, \dot{x}_j)\ddot{x}_r + \mathcal{H}_i(x_j, \dot{x}_j) = 0, \quad (i = 1, 2). \quad (4.3)$$

Bearing in mind the obvious relations

$$\frac{d}{dt}\mathcal{A}_{ir} = \dot{\mathcal{A}}_{ir} = \frac{\partial \mathcal{A}_{ir}}{\partial x_j}\dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j}\ddot{x}_j, \quad (4.4)$$

$$\frac{d}{dt}\mathcal{H}_i = \dot{\mathcal{H}}_i = \frac{\partial \mathcal{H}_i}{\partial x_j}\dot{x}_j + \frac{\partial \mathcal{H}_i}{\partial \dot{x}_j}\ddot{x}_j, \quad (4.5)$$

the first-order time derivative of (4.3) gives

$$\mathcal{A}_{ir}\frac{d^2}{dt^2}\dot{x}_r + \left(\frac{\partial \mathcal{A}_{ir}}{\partial x_j}\dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j}\ddot{x}_j\right)\frac{d}{dt}\dot{x}_r + \frac{\partial \mathcal{H}_i}{\partial x_j}\dot{x}_j + \frac{\partial \mathcal{H}_i}{\partial \dot{x}_j}\ddot{x}_j = 0, \quad (i = 1, 2), \quad (4.6)$$

which represents a nonlinear system of differential equations whose solutions are given by the functions $\dot{x}_r$ ($r = 1, 2$). The second-order time derivative of (4.3) leads to the system

$$\begin{aligned} \mathcal{A}_{ir}\frac{d^2}{dt^2}\ddot{x}_r + 2\left(\frac{\partial \mathcal{A}_{ir}}{\partial x_j}\dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j}\ddot{x}_j\right)\frac{d}{dt}\ddot{x}_r + \ddot{x}_r\frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k}\frac{d}{dt}\ddot{x}_k + \frac{\partial \mathcal{H}_i}{\partial \dot{x}_k}\frac{d}{dt}\ddot{x}_k + \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k}\dot{x}_l\dot{x}_k + 2\frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k}\ddot{x}_l\dot{x}_k \right. \\ \left. + \frac{\partial \mathcal{A}_{ir}}{\partial x_k}\ddot{x}_k + \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial \dot{x}_k}\ddot{x}_l\ddot{x}_k\right)\ddot{x}_r + \frac{\partial^2 \mathcal{H}_i}{\partial x_l \partial x_k}\dot{x}_l\dot{x}_k + 2\frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial x_k}\ddot{x}_l\dot{x}_k + \frac{\partial \mathcal{H}_i}{\partial x_k}\ddot{x}_k + \frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial \dot{x}_k}\ddot{x}_l\ddot{x}_k = 0, \quad (i = 1, 2), \end{aligned} \quad (4.7)$$

still representing a nonlinear set of differential equations in the unknowns $\ddot{x}_r$. However, by patiently applying the time derivative operator four times to Eq. (4.3) (see Appendix A for details) the fourth time derivatives $x_r^{(4)}$ are found to solve a *linear* system of ordinary differential equations of the form

$$\mathcal{A}_{ir}\frac{d^2}{dt^2}x_r^{(4)} + 4\left(\frac{\partial \mathcal{A}_{ir}}{\partial x_j}\dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j}\ddot{x}_j\right)\frac{d}{dt}x_r^{(4)} + \ddot{x}_r\left(\frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k}\right)\frac{d}{dt}x_k^{(4)} + \left(\frac{\partial \mathcal{H}_i}{\partial \dot{x}_k}\right)\frac{d}{dt}x_k^{(4)} + F_i(x_p, \dot{x}_p, \dots, x_p^{(4)}) = 0, \quad (4.8)$$

($i, p = 1, 2$).

The form of $F_i(x_p, \dot{x}_p, \dots, x_p^{(4)})$ can be read from Eq. (A3), where it is explicitly shown that this term does not depend on time derivatives of x_p having an order higher than the fourth and that its dependence on $x_p^{(4)}$ turns out to

be linear. Therefore, we have obtained the original result according to which the differential equations describing the motion of a body in our Solar System within the first post-Newtonian approximation can be put in linear form if the fourth time derivative of the original equation is employed. In our analysis, we have been inspired by the work in Ref. [90], where the author differentiated (see below) a system of nonlinear partial differential equations to arrive at their solution. More precisely, starting from a system of n nonlinear second-order hyperbolic partial differential equations of the form

$$A^{\lambda\mu} \frac{\partial^2 w_s}{\partial x^\lambda \partial x^\mu} + f_s = 0, \quad s = 1, 2, \dots, n, \quad \lambda, \mu = 1, 2, 3, 4, \quad (4.9)$$

the coefficients $A^{\lambda\mu}$ and f_s depending in a nonlinear way on the four variables x^α , the unknown functions w_s , and their first time derivatives $\frac{\partial w_s}{\partial x^\alpha}$, in Ref. [90] it is shown that by applying five times of the derivative operator with respect to any whatsoever variable x^α leads to the linear system

$$A^{\lambda\mu} \frac{\partial^2 U_S}{\partial x^\lambda \partial x^\mu} + \frac{\partial U_S}{\partial x^\lambda} + F_S = 0, \quad S = 1, 2, \dots, N, \quad \lambda, \mu = 1, 2, 3, 4,^5 \quad (4.10)$$

where U_S represents the partial derivatives of order five of w_s and F_S is a function of the four variables x^α , the unknown functions w_s , and of their partial derivatives up to the fifth order included, but not of the derivatives of higher order. In particular, when this framework is applied to Einstein's field equations (which represent a system of $n = 10$ quasilinear partial differential equations) it suffices to derive them four times in order to obtain a linear system, where the unknowns are given by the fourth derivatives of the metric tensor components [90].

Having obtained a linear system of differential equations of second order associated to (3.31), we might in principle exploit their reduction to canonical form (Appendix B) and the rich qualitative theory [91] of such equations. In fact, as we have seen in Sec. III, our model predicts the presence of periodic solutions (see for example Figs. 1, 2, and 3) in simple cases, and hence the qualitative theory just mentioned and the encouraging evidence for simple (but nontrivial) two-body systems suggest undertaking the much harder analysis of N -body systems.

V. QUANTUM EFFECTS ON LAGRANGIAN POINTS REVISITED

The work in Secs. II and III puts on firm ground the investigations initiated in Refs. [1–4]. In other words, the motion of celestial bodies in the solar system can be studied by employing a Lagrangian that is almost independent of the internal structure [19]. Within Einstein's theory, the *effacement* property of Newtonian theory is still valid at the first post-Newtonian approximation, because all large, direct self-action effects cancel each other or contribute terms in the equations of motion which can be removed, so that the final equations can be written in terms of only some centers of mass and some effective masses [19, 26, 27]. But then, to the extent that the effective-gravity prescriptions are correct, it becomes legitimate to consider Newtonian potential terms among large masses in the Lagrangian used to derive geodesic motion of planets and other bodies, and insert therein the quantum modifications worked out in Refs. [5, 6, 11–13].

In particular, the work of Ref. [4] has studied in detail how quantum corrections on the relativistic position of the Earth-Moon Lagrangian points can be evaluated. In fact the effective gravity picture modifies the Newtonian potential among bodies of masses m_A and m_B in the low-energy/long-distance domain through the asymptotic expansion

$$\begin{aligned} V_E(r) &\sim -\frac{G_N m_A m_B}{r} \left[1 + \left(\kappa_1 \frac{(l_A + l_B)}{r} + \kappa_2 \frac{(l_P)^2}{r^2} + O(G_N^2) \right) \right] \\ \Rightarrow \frac{V_E(r)}{c^2 m_B} &\sim -\frac{l_A}{r} \left[1 + \left(\kappa_1 \frac{(l_A + l_B)}{r} + \kappa_2 \frac{(l_P)^2}{r^2} + O(G_N^2) \right) \right], \end{aligned} \quad (5.1)$$

l_A and l_B being the gravitational radii of the bodies and l_P the Planck length, whereas κ_1 and κ_2 represent constants⁶ resulting from the calculation of Feynman diagrams involved in the particular definition adopted for the potential:

⁵ N represents the product by n of the number of partial derivatives of order five of a function of four variables.

⁶ In this conceptual scheme, physical phenomena are described of course in a classical way on large distances, but the precise values of some coefficients depend on the underlying quantum theory. Thus, the dimensionless κ_1 parameter is the effective-gravity weight of the purely classical term $\frac{(l_A + l_B)}{r}$.

one-particle reducible, scattering or bound-states (see Tab. I). The term $\kappa_1 \frac{(l_A + l_B)}{r}$ in (5.1) refers to a post-Newtonian correction to the classical potential, whereas $\kappa_2 \frac{(l_P)^2}{r^2}$ represents a fully quantum term, depending on the square of the Planck length.

TABLE I: The values assumed by κ_1 and κ_2 in the three different potentials.

κ_i	one-particle reducible	scattering	bound-states
κ_1	-1	3	$-\frac{1}{2}$
κ_2	$-\frac{167}{30\pi}$	$\frac{41}{10\pi}$	$\frac{41}{10\pi}$

In Ref. [4] we have proposed a framework where the aforementioned quantum corrections to Lagrangian points can be obtained by constructing a map whose form is inspired by the pattern enlightened in Eq. (5.1). In particular, we have applied the map

$$(U_\alpha, U_\beta) \rightarrow (V_\alpha, V_\beta), \quad (5.2)$$

with

$$U_\alpha(r) \equiv \frac{l_\alpha}{r} = U_\alpha, \quad (5.3)$$

$$U_\beta(s) \equiv \frac{l_\beta}{s} = U_\beta, \quad (5.4)$$

$$V_\alpha(r) \sim \left[1 + \kappa_2 \frac{(l_P)^2}{r^2} \right] U_\alpha(r) + \kappa_1 (U_\alpha(r))^2 + O(G_N^3), \quad (5.5)$$

$$V_\beta(s) \sim \left[1 + \kappa_2 \frac{(l_P)^2}{s^2} \right] U_\beta(s) + \kappa_1 (U_\beta(s))^2 + O(G_N^3), \quad (5.6)$$

to the Lagrangian describing the motion of the planetoid in the gravitational field generated by the Earth and the Moon, which, upon adopting the set of coordinates (ct, ξ, η, ζ) (c being the speed of light) reads as

$$L = \frac{1}{2} \sum_{\mu, \nu=0}^3 g_{\mu\nu} \frac{dx^\mu}{dt} \frac{dx^\nu}{dt}, \quad (5.7)$$

where

$$\begin{aligned} g_{00} = & 1 - 2\frac{l_\alpha}{r} - 2\frac{l_\beta}{s} - \frac{\Omega^2}{c^2}(\xi^2 + \eta^2) + 2 \left[\left(\frac{l_\alpha}{r} \right)^2 + \left(\frac{l_\beta}{s} \right)^2 \right] \\ & - 2 \frac{(l_\alpha + l_\beta)}{R^3} \left(\frac{l_\alpha}{r} + \frac{l_\beta}{s} \right) (\xi^2 + \eta^2) + 4 \frac{l_\alpha}{r} \frac{l_\beta}{s} \\ & + \frac{(2-\rho)}{(1+\rho)} \frac{l_\alpha}{r} \frac{l_\beta}{R} + \frac{(2\rho-1)}{(1+\rho)} \frac{l_\beta}{s} \frac{l_\alpha}{R} - 7 \frac{\xi}{R^2} \left(\frac{l_\alpha}{r} l_\beta - \frac{l_\beta}{s} l_\alpha \right) \\ & + (1+\rho)^{-1} \frac{\eta^2}{R} \left[\rho \left(\frac{l_\alpha}{r} \right)^3 \frac{l_\beta}{(l_\alpha)^2} + \left(\frac{l_\beta}{s} \right)^3 \frac{l_\alpha}{(l_\beta)^2} \right], \end{aligned} \quad (5.8)$$

$$2cg_{01} = \left(1 + 2\frac{l_\alpha}{r} + 2\frac{l_\beta}{s} \right) 2\Omega\eta, \quad (5.9)$$

$$2cg_{02} = - \left(1 + 2\frac{l_\alpha}{r} + 2\frac{l_\beta}{s} \right) 2\Omega\xi - 8\frac{\Omega^2 R}{(1+\rho)} \left(\rho\frac{l_\alpha}{r} - \frac{l_\beta}{s} \right), \quad (5.10)$$

$$g_{03} = 0, \quad (5.11)$$

$$g_{ij} = - \left(1 + 2\frac{l_\alpha}{r} + 2\frac{l_\beta}{s} \right) \delta_{ij}, \quad i, j = 1, 2, 3. \quad (5.12)$$

l_α and l_β being the gravitational radii of the Earth and the Moon, respectively, r and s their respective distances from the planetoid, ρ the ratio between the masses of the Earth and the Moon, R their distance, and

$$\Omega \equiv \omega \left[1 - \frac{3}{2} \frac{(l_\alpha + l_\beta)}{R} \left(1 - \frac{1}{3} \frac{\rho}{(1+\rho)^2} \right) \right], \quad (5.13)$$

where ω represents the Newtonian pulsation.

However, since effective field theories of gravity provide quantum corrections to the Newtonian potential among bodies but not to its powers, we now consider a more refined prescription where only the purely Newtonian terms (i.e., those which are linear or bilinear in U_α and U_β) are corrected through the map (5.5) and (5.6), while the remaining ones are left unchanged. According to this new perspective, the effective gravity Lagrangian L_V can be obtained from (5.7)-(5.12) by setting

$$\begin{aligned} U_\alpha &\rightarrow V_\alpha, \\ U_\beta &\rightarrow V_\beta, \\ U_\alpha U_\beta &\rightarrow V_\alpha V_\beta, \\ (U_\alpha)^n &\rightarrow (U_\alpha)^n, \quad n > 1, \\ (U_\beta)^n &\rightarrow (U_\beta)^n, \quad n > 1. \end{aligned} \quad (5.14)$$

The new map (5.14) is such that the quantum corrected Lagrangian reads as

$$\begin{aligned} L_V &= \frac{c^2}{2} \left\{ 1 - 2(V_\alpha + V_\beta) - \frac{\Omega^2}{c^2} (\xi^2 + \eta^2) + 2[(U_\alpha)^2 + (U_\beta)^2] \right. \\ &\quad - 2\frac{(l_\alpha + l_\beta)}{R^3} (\xi^2 + \eta^2)(V_\alpha + V_\beta) + 4V_\alpha V_\beta \\ &\quad + \frac{(2-\rho)}{(1+\rho)} \frac{l_\beta}{R} V_\alpha + \frac{(2\rho-1)}{(1+\rho)} \frac{l_\alpha}{R} V_\beta - 7\frac{\xi}{R^2} (l_\beta V_\alpha - l_\alpha V_\beta) \\ &\quad \left. + (1+\rho)^{-1} \frac{\eta^2}{R} \left[\rho \frac{l_\beta}{(l_\alpha)^2} (U_\alpha)^3 + \frac{l_\alpha}{(l_\beta)^2} (U_\beta)^3 \right] \right\} \\ &\quad - \frac{1}{2} (\dot{\xi}^2 + \dot{\eta}^2 + \dot{\zeta}^2) \left[1 + 2(V_\alpha + V_\beta) \right] + \Omega\eta\dot{\xi} \left[1 + 2(V_\alpha + V_\beta) \right] \\ &\quad - \Omega\xi\dot{\eta} \left[1 + 2(V_\alpha + V_\beta) \right] - 4\frac{\Omega^2 R}{(1+\rho)} \dot{\eta} (\rho V_\alpha - V_\beta). \end{aligned} \quad (5.15)$$

Therefore, new values of quantum corrections on the relativistic distances in the Earth-Moon system [4]

$$\begin{aligned} r_{1,GR} &= 3.2637628817407598555 \times 10^8 \text{ m}, \\ r_{2,GR} &= 4.4892056003414800050 \times 10^8 \text{ m}, \\ r_{3,GR} &= 3.8167471569392170594 \times 10^8 \text{ m}, \\ r_{4,GR} &= r_{5,GR} = 3.8439999999998611069 \times 10^8 \text{ m}, \end{aligned} \quad (5.16)$$

are found and displayed in Tab. II. Needless to say, the theoretical expression and value of such quantum corrections (if they exist) to Lagrangian points remain an open problem, because the classical effacement property holds only approximately [19, 26, 27], while 3 sets of quantum parameters κ_1 and κ_2 are conceivable in effective gravity [5, 6, 11–13]. In this respect, the quantum corrections to general relativity values for noncollinear points L_4 and L_5 look encouraging in the case of (κ_1, κ_2) values appropriate to scattering potential, because corrections just below a centimeter are comparable with the purely instrumental, time-of-flight uncertainty of the geodesic positioning techniques based laser-ranging [63–75]. However, the total error budget of satellite/lunar laser ranging [63–75] varies with the specific application and/or orbit, at the level of millimeter to centimeter.

TABLE II: Quantum corrections on the relativistic position of Earth-Moon Lagrangian points for three different types of potential obtained by considering the Lagrangian function (5.15).

Quantum corrections on Lagrangian points			
L_i	One-particle reducible	Scattering	Bound-states
L_1	$r_Q - r_{GR} = -1.27$ mm	$r_Q - r_{GR} = 3.67$ mm	$r_Q - r_{GR} = -0.65$ mm
L_2	$r_Q - r_{GR} = -0.75$ mm	$r_Q - r_{GR} = 2.39$ mm	$r_Q - r_{GR} = -0.35$ mm
L_3	$r_Q - r_{GR} = -2.96$ mm	$r_Q - r_{GR} = 8.89$ mm	$r_Q - r_{GR} = -1.48$ mm
L_4	$r_Q - r_{GR} = -2.98$ mm	$r_Q - r_{GR} = 8.85$ mm	$r_Q - r_{GR} = -1.50$ mm
L_5	$r_Q - r_{GR} = -2.98$ mm	$r_Q - r_{GR} = 8.85$ mm	$r_Q - r_{GR} = -1.50$ mm

VI. CONCLUDING REMARKS AND OPEN PROBLEMS

In the first part of our paper, to prepare the ground for future work, we have provided an original synthesis of the Levi-Civita analysis of the problem of motion of N bodies in general relativity, a very difficult problem that was studied, among the others, by Einstein himself with Infeld and Hoffman [17], Fock [18], Levi-Civita [19], Damour, Soffel and Xu [29–32].

The Sun-Earth, Sun-Earth-Moon and N -body dynamics have been investigated in Sec. III, while Sec. IV contains our original proof that the nonlinear ordinary differential equations for planetary motions can be always mapped into an exact, linear system of ordinary differential equations, where the unknowns are the fourth-order time derivatives of the original unknown functions. In Sec. V, the evaluation of quantum corrections to location of collinear and noncollinear Lagrangian points for the planar restricted three-body problem has been revisited, and a new set of theoretical values of such corrections for the Earth-Moon-planetoid system has been displayed. It is clear from Tab. II that the few millimeters quantum corrections regarding the relativistic position of Lagrangian points represent a huge obstacle for future experimental measurements. Nevertheless, Einstein theory produces more pronounced (classical) effects on larger distances than those involved in the Earth-Moon system, but at the cost of increasing the efforts for reaching more distant planets. As an example, by applying the framework based on the Lagrangian function (5.7)-(5.12) (see Ref. [4] for further details) to the Sun-Earth and Sun-Jupiter systems we obtain the corrections reported in Tabs. III and IV [79].

Notably, the values of Tab. III agree with those of Ref. [34], and those of Tab. IV with the ones reported in Refs. [45, 49]. Moreover, we are aware of the fact that many satellites are currently situated near the Sun, but unluckily none of them is planned to approach the Lagrangian point L_1 in order to test our theoretical model. Eventually, the situation becomes far more complicated for the Sun-Jupiter system because of the large distances involved.

In the years to come, we hope that our result in Sec. IV, jointly with the qualitative methods [91] for linear second-order ordinary differential equations in canonical form (see (B4) and (B5)), will lead to improved theoretical calculations of planetary motions in the solar system, with a wide range of applications in fundamental and applied science.

Last but not least, it will be also very interesting to compare planetary motions according to Ref. [18], where the energy-momentum tensor inside and outside bodies is obtained, with the planetary motions according to Ref. [19], where the effacement property plays instead a key role as we have seen in our Sec. II. None of the investigations of N -body dynamics published so far in the literature can indeed claim complete superiority over the others.

TABLE III: General relativity corrections on the position of Newtonian Lagrangian points for the Sun-Earth system obtained by considering the general relativity Lagrangian (5.7-5.12). The differences involved refer to the distances of the Sun from the planetoid.

General relativity corrections on the Sun-Earth system	
L_i	Corrections
L_1	$r_{GR} - r_{cl} = 4.8 \text{ m}$
L_2	$r_{GR} - r_{cl} = -5.0 \text{ m}$
L_3	$r_{GR} - r_{cl} = -0.3 \text{ cm}$
$L_{4,5}$	$r_{GR} - r_{cl} = -0.3 \text{ cm}$

TABLE IV: General relativity corrections on the position of Newtonian Lagrangian points for the Sun-Jupiter system obtained by employing the general relativity Lagrangian (5.7-5.12). The differences involved refer to the distances of the Sun from the planetoid.

General relativity corrections on the Sun-Jupiter system	
L_i	Corrections
L_1	$r_{GR} - r_{cl} = 30 \text{ m}$
L_2	$r_{GR} - r_{cl} = -38 \text{ m}$
L_3	$r_{GR} - r_{cl} = -1 \text{ m}$
$L_{4,5}$	$r_{GR} - r_{cl} = -1 \text{ m}$

Appendix A: Linear differential equations associated to nonlinear ones

In this appendix we provide the details of the calculations leading to (4.8) starting from Eq (3.31) written in the form given by (4.3). As explained in Sec. IV, bearing in mind Eqs. (4.4) and (4.5), the first- and second-order time derivatives of (4.3) are given by Eqs. (4.6) and (4.7), respectively.

The third time derivative of (4.3) gives a nonlinear system of differential equations with unknowns $x_p^{(3)}$ ($p = 1, 2$)

having the form

$$\begin{aligned}
& \mathcal{A}_{ir} \frac{d^2}{dt^2} x_r^{(3)} + 3 \left(\frac{\partial \mathcal{A}_{ir}}{\partial x_j} \dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \ddot{x}_j \right) \frac{d}{dt} x_r^{(3)} + \ddot{x}_r \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k} \frac{d}{dt} x_k^{(3)} + \frac{\partial \mathcal{H}_i}{\partial \dot{x}_k} \frac{d}{dt} x_k^{(3)} + 2 \left[\left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial x_j} \right) \dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial x_j} \ddot{x}_j \right. \\
& + \left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \right) \ddot{x}_j + \left(\frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \right) x_j^{(3)} \left. \right] x_r^{(3)} + x_r^{(3)} \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k} x_k^{(3)} + \ddot{x}_r \left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k} \right) x_k^{(3)} + \left(\frac{d}{dt} \frac{\partial \mathcal{H}_i}{\partial \dot{x}_k} \right) x_k^{(3)} + \left[\left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \right) \dot{x}_l \dot{x}_k \right. \\
& + \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \right) (\ddot{x}_l \dot{x}_k + \dot{x}_l \ddot{x}_k) + 2 \left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \right) \ddot{x}_l \dot{x}_k + 2 \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \right) (x_l^{(3)} \dot{x}_k + \dot{x}_l x_k^{(3)}) + \left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial x_k} \right) \ddot{x}_k \\
& + \left(\frac{\partial \mathcal{A}_{ir}}{\partial x_k} \right) x_k^{(3)} + \left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_k \partial \dot{x}_l} \right) \ddot{x}_l \ddot{x}_k + \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_k \partial \dot{x}_l} \right) (x_l^{(3)} \ddot{x}_k + \ddot{x}_l x_k^{(3)}) \left. \right] \ddot{x}_r + \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \dot{x}_l \dot{x}_k + 2 \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \ddot{x}_l \dot{x}_k \right. \\
& + \frac{\partial \mathcal{A}_{ir}}{\partial x_k} \ddot{x}_k + \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial \dot{x}_k} \ddot{x}_l \ddot{x}_k \left. \right) x_r^{(3)} + \left(\frac{d}{dt} \frac{\partial^2 \mathcal{H}_i}{\partial x_l \partial x_k} \right) \dot{x}_l \dot{x}_k + \frac{\partial^2 \mathcal{H}_i}{\partial x_l \partial x_k} (\ddot{x}_l \dot{x}_k + \dot{x}_l \ddot{x}_k) + 2 \left(\frac{d}{dt} \frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial x_k} \right) \ddot{x}_l \dot{x}_k \\
& + 2 \left(\frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial x_k} \right) (x_l^{(3)} \dot{x}_k + \ddot{x}_l \ddot{x}_k) + \left(\frac{d}{dt} \frac{\partial \mathcal{H}_i}{\partial x_k} \right) \ddot{x}_k + \frac{\partial \mathcal{H}_i}{\partial x_k} x_k^{(3)} + \left(\frac{d}{dt} \frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial \dot{x}_k} \right) \ddot{x}_k \ddot{x}_l + \left(\frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial \dot{x}_k} \right) (x_l^{(3)} \ddot{x}_k + \ddot{x}_l x_k^{(3)}) = 0, \\
& (i = 1, 2).
\end{aligned} \tag{A1}$$

The nonlinearities occurring in (A1) vanish if we compute the fourth time derivative. In fact, as anticipated in Sec. IV, by differentiating once again Eq. (A1) we end up with a *linear* system of coupled ordinary differential equations for the unknown functions $x_p^{(4)}$ ($p = 1, 2$) which can be written as

$$\begin{aligned}
& \mathcal{A}_{ir} \frac{d^2}{dt^2} x_r^{(4)} + 4 \left(\frac{\partial \mathcal{A}_{ir}}{\partial x_j} \dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \ddot{x}_j \right) \frac{d}{dt} x_r^{(4)} + \ddot{x}_r \left(\frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k} \right) \frac{d}{dt} x_k^{(4)} + \left(\frac{\partial \mathcal{H}_i}{\partial \dot{x}_k} \right) \frac{d}{dt} x_k^{(4)} + F_i \left(x_p, \dot{x}_p, \dots, x_p^{(4)} \right) = 0, \\
& (i, p = 1, 2).
\end{aligned} \tag{A2}$$

where

$$\begin{aligned}
F_i \left(x_p, \dot{x}_p, \dots, x_p^{(4)} \right) = & 3 \left[\frac{d}{dt} \left(\frac{\partial \mathcal{A}_{ir}}{\partial x_j} \dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \ddot{x}_j \right) \right] x_r^{(4)} + 2 \left\{ \frac{d}{dt} \left[\left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial x_j} \right) \dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial x_j} \ddot{x}_j + \left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \right) \ddot{x}_j \right. \right. \\
& + \left. \left. \left(\frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \right) x_j^{(3)} \right] \right\} x_r^{(3)} + 2 \left[\left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial x_j} \right) \dot{x}_j + \frac{\partial \mathcal{A}_{ir}}{\partial x_j} \ddot{x}_j + \left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \right) \ddot{x}_j + \left(\frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_j} \right) x_j^{(3)} \right] x_r^{(4)} \\
& + \left(\frac{d^2}{dt^2} \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k} \right) \ddot{x}_r x_k^{(3)} + 2 \left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k} \right) \left(x_r^{(3)} x_k^{(3)} + \ddot{x}_r x_k^{(4)} \right) + \left(\frac{\partial \mathcal{A}_{ir}}{\partial \dot{x}_k} \right) \left(x_r^{(4)} x_k^{(3)} + 2 x_r^{(3)} x_k^{(4)} \right) \\
& + \left(\frac{d^2}{dt^2} \frac{\partial \mathcal{H}_i}{\partial \dot{x}_k} \right) x_k^{(3)} + 2 \left(\frac{d}{dt} \frac{\partial \mathcal{H}_i}{\partial \dot{x}_k} \right) x_k^{(4)} + \left\{ \frac{d}{dt} \left[\left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \right) \dot{x}_l \dot{x}_k + \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \right) (\dot{x}_l \dot{x}_k + \dot{x}_l \ddot{x}_k) \right. \right. \\
& + 2 \left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \right) \ddot{x}_l \dot{x}_k + 2 \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \right) (x_l^{(3)} \dot{x}_k + \ddot{x}_l \ddot{x}_k) + \left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial x_k} \right) \ddot{x}_k + \left(\frac{\partial \mathcal{A}_{ir}}{\partial x_k} \right) x_k^{(3)} \\
& + \left. \left. \left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_k \partial \dot{x}_l} \right) \ddot{x}_l \ddot{x}_k + \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_k \partial \dot{x}_l} \right) (x_l^{(3)} \ddot{x}_k + \ddot{x}_l x_k^{(3)}) \right] \right\} \ddot{x}_r + \left[\left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \right) \dot{x}_l \dot{x}_k \right. \\
& + \left. \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \right) (\dot{x}_l \dot{x}_k + \dot{x}_l \ddot{x}_k) + 2 \left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \right) \ddot{x}_l \dot{x}_k + 2 \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \right) (x_l^{(3)} \dot{x}_k + \ddot{x}_l \ddot{x}_k) \right. \\
& + \left. \left(\frac{d}{dt} \frac{\partial \mathcal{A}_{ir}}{\partial x_k} \right) \ddot{x}_k + \left(\frac{\partial \mathcal{A}_{ir}}{\partial x_k} \right) x_k^{(3)} + \left(\frac{d}{dt} \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_k \partial \dot{x}_l} \right) \ddot{x}_l \ddot{x}_k + \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_k \partial \dot{x}_l} \right) (x_l^{(3)} \ddot{x}_k + \ddot{x}_l x_k^{(3)}) \right] x_r^{(3)} \\
& + \left[\frac{d}{dt} \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \dot{x}_l \dot{x}_k + 2 \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \ddot{x}_l \dot{x}_k + \frac{\partial \mathcal{A}_{ir}}{\partial x_k} \ddot{x}_k + \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial \dot{x}_k} \dot{x}_l \ddot{x}_k \right) \right] x_r^{(3)} + \left(\frac{\partial^2 \mathcal{A}_{ir}}{\partial x_l \partial x_k} \dot{x}_l \dot{x}_k \right. \\
& + 2 \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial x_k} \ddot{x}_l \dot{x}_k + \frac{\partial \mathcal{A}_{ir}}{\partial x_k} \ddot{x}_k + \frac{\partial^2 \mathcal{A}_{ir}}{\partial \dot{x}_l \partial \dot{x}_k} \dot{x}_l \ddot{x}_k \left. \right) x_r^{(4)} + \left(\frac{d^2}{dt^2} \frac{\partial^2 \mathcal{H}_i}{\partial x_l \partial x_k} \right) \dot{x}_l \dot{x}_k \\
& + 2 \left(\frac{d}{dt} \frac{\partial^2 \mathcal{H}_i}{\partial x_l \partial x_k} \right) (\dot{x}_l \dot{x}_k + \dot{x}_l \ddot{x}_k) + \left(\frac{\partial^2 \mathcal{H}_i}{\partial x_l \partial x_k} \right) (x_l^{(3)} \dot{x}_k + 2 \ddot{x}_l \ddot{x}_k + \dot{x}_l x_k^{(3)}) + 2 \left(\frac{d^2}{dt^2} \frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial x_k} \right) \ddot{x}_l \dot{x}_k \\
& + 4 \left(\frac{d}{dt} \frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial x_k} \right) (x_l^{(3)} \dot{x}_k + \ddot{x}_l \ddot{x}_k) + 2 \left(\frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial x_k} \right) (x_l^{(4)} \dot{x}_k + 2 x_l^{(3)} \ddot{x}_k + \ddot{x}_l x_k^{(3)}) + \left(\frac{d^2}{dt^2} \frac{\partial \mathcal{H}_i}{\partial x_k} \right) \ddot{x}_k \\
& + 2 \left(\frac{d}{dt} \frac{\partial \mathcal{H}_i}{\partial x_k} \right) x_k^{(3)} + \frac{\partial \mathcal{H}_i}{\partial x_k} x_k^{(4)} + \left(\frac{d^2}{dt^2} \frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial \dot{x}_k} \right) \ddot{x}_l \ddot{x}_k + 2 \left(\frac{d}{dt} \frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial \dot{x}_k} \right) (x_l^{(3)} \ddot{x}_k + \ddot{x}_l x_k^{(3)}) \\
& + \frac{\partial^2 \mathcal{H}_i}{\partial \dot{x}_l \partial \dot{x}_k} \left(x_l^{(4)} \ddot{x}_k + 2 x_k^{(3)} x_l^{(3)} + \ddot{x}_l x_k^{(4)} \right), \quad (i, p = 1, 2).
\end{aligned} \tag{A3}$$

It is thus clear from last equation that the functions $F_i \left(x_p, \dot{x}_p, \dots, x_p^{(4)} \right)$ ($i = 1, 2$) depend linearly on $x_p^{(4)}$ ($p = 1, 2$) and that no derivatives of order higher than four appear, which clearly means that Eq. (A2) (or equivalently (4.8)) is linear with respect to $x_p^{(4)}$.⁷

⁷ Recall that the most general form of a linear ordinary differential equation of order n is given by

$$\left(r_n(t) \frac{d^n}{dt^n} + r_{n-1}(t) \frac{d^{n-1}}{dt^{n-1}} + \dots + r_1(t) \frac{d}{dt} + r_0(t) \right) x(t) = f(t), \tag{A4}$$

the coefficients $r_n(t), r_{n-1}(t), \dots, r_1(t), r_0(t)$ and the term $f(t)$ being continuous real-valued functions of t in the interval $a \leq t \leq b$ and $r_n(t)$ a function that does not vanish at any point of the aforementioned interval. The operator

$$L \equiv r_n(t) \frac{d^n}{dt^n} + r_{n-1}(t) \frac{d^{n-1}}{dt^{n-1}} + \dots + r_1(t) \frac{d}{dt} + r_0(t), \tag{A5}$$

is called linear differential operator of order n .

Appendix B: Linear differential equations of second order. Sturm-Liouville problem

Within the framework of ordinary differential equations, every second-order linear differential equation can be written in the form

$$\left[\frac{d^2}{dx^2} + p(x) \frac{d}{dx} + q(x) \right] u(x) = 0, \quad (\text{B1})$$

where x is taken to lie in the closed interval $[a, b]$, while p and q are suitably smooth functions. This equation can be brought into the Liouville form, where the coefficient of the first-order derivative vanishes. For this purpose, one sets $u(x) = \varphi(x)\psi(x)$, so that Eq. (B1) reads as

$$\varphi\psi'' + (2\varphi' + p\varphi)\psi' + (\varphi'' + p\varphi' + q\varphi)\psi = 0. \quad (\text{B2})$$

Our task is achieved if the function φ solves the first-order equation

$$\frac{\varphi'}{\varphi} = -\frac{p}{2} \implies \varphi(x) = \exp\left(-\frac{1}{2} \int p(x) dx\right). \quad (\text{B3})$$

At this stage, since we can divide by φ , we can re-express Eq. (B2) in the form

$$\psi'' + W\psi = 0, \quad (\text{B4})$$

where the potential terms turns out to be

$$W = \frac{\varphi''}{\varphi} + p\frac{\varphi'}{\varphi} + q = -\frac{1}{2}p' - \frac{p^2}{4} + q. \quad (\text{B5})$$

Once we have reduced ourselves to studying Eq. (B4), one can deduce important qualitative properties. For example, if the function W is continuous for $x \in [a, b]$, and if there exist real constants ω and Ω such that

$$0 < \omega^2 < W(x) < \Omega^2, \quad (\text{B6})$$

one can compare the zeros of solutions of Eq. (B4) with the zeros of solutions of the equations

$$\psi'' + \alpha^2\psi = 0, \quad \alpha = \omega \text{ or } \Omega. \quad (\text{B7})$$

Equations (B7) are solved by periodic functions $\sin \alpha(x - x_0)$ which have zeros at $x_0 + \frac{k\pi}{\alpha}$, k being an integer and α taking one of the two values in (B7). One can then prove that the difference δ between two adjacent zeros of a solution of Eq. (B4) satisfies the conditions [92]

$$\frac{\pi}{\Omega} \leq \delta \leq \frac{\pi}{\omega}. \quad (\text{B8})$$

Now the first of our Eqs. (3.15) (or the more general (3.31)) can be written, after using the procedure used in Sec. IV, i.e. four differentiations with respect to t of the original equation, in the form

$$g(t) [Y'' + p_1(t)Y' + q_1Y] + f(t) [X'' + p_2(t)X' + q_2X] = 0, \quad (\text{B9})$$

where, for each pair (p_1, q_1) and (p_2, q_2) , we can evaluate the corresponding potential W_1 (respectively W_2) according to Eq. (B5). The letters X and Y here denote fourth-order time derivative of the original functions $x(t)$ and $y(t)$, respectively.

Acknowledgments

E. B. and G. E. are grateful to the Dipartimento di Fisica “Ettore Pancini” of Federico II University for hospitality and support. The work of E. B., S. D. and G. E. has been supported by the INFN funding of the NEWREFLECTIONS experiment.

[1] E. Battista and G. Esposito, Restricted three-body problem in effective-field-theory models of gravity, Phys. Rev. D **89**, 084030 (2014).

- [2] E. Battista and G. Esposito, Full three-body problem in effective-field-theory models of gravity, *Phys. Rev. D* **90**, 084010 (2014); *Phys. Rev. D* **93**, 049901(E) (2016).
- [3] E. Battista, S. Dell’Agnello, G. Esposito and J. Simo, Quantum effects on Lagrangian points and displaced periodic orbits in the Earth-Moon system, *Phys. Rev. D* **91**, 084041 (2015); *Phys. Rev. D* **93**, 049902(E) (2016).
- [4] E. Battista, S. Dell’Agnello, G. Esposito, L. Di Fiore, J. Simo and A. Grado, Earth-Moon Lagrangian points as a test bed for general relativity and effective field theories of gravity, *Phys. Rev. D* **92**, 064045 (2015); *Phys. Rev. D* **93**, 109904(E) (2016).
- [5] J. F. Donoghue, Leading quantum correction to the Newtonian potential, *Phys. Rev. Lett.* **72**, 2996 (1994).
- [6] J. F. Donoghue, General relativity as an effective field theory: the leading quantum corrections, *Phys. Rev. D* **50**, 3874 (1994).
- [7] I. J. Muzinich and S. Vokos, Long range forces in quantum gravity, *Phys. Rev. D* **52**, 3472 (1995).
- [8] H. W. Hamber and S. Liu, On the quantum corrections to the Newtonian potential, *Phys. Lett. B* **357**, 51 (1995).
- [9] A. A. Akhundov, S. Bellucci, and A. Shiekh, Gravitational interaction to one loop in effective quantum gravity, *Phys. Lett. B* **395**, 16 (1997).
- [10] I. B. Khriplovich and G. G. Kirilin, Quantum power correction to the Newton law, *Sov. Phys. JETP* **95**, 981 (2002).
- [11] N. E. J. Bjerrum-Bohr, J. F. Donoghue, and B. R. Holstein, Quantum gravitational corrections to the nonrelativistic scattering potential of two masses, *Phys. Rev. D* **67**, 084033 (2003).
- [12] N. E. J. Bjerrum-Bohr, J. F. Donoghue, and B. R. Holstein, Quantum corrections to the Schwarzschild and Kerr metrics, *Phys. Rev. D* **68**, 084005 (2003).
- [13] J. F. Donoghue, The effective field theory treatment of quantum gravity, *AIP Conf. Proc.* **1483**, 73 (2012).
- [14] F. F. Tisserand, *Treatise of Celestial Mechanics*, Vols. 1 to 4 (Gauthier-Villars, Paris, 1889-1896; Jacques Gabay, Paris, 1990).
- [15] H. Poincaré, The three-body problem and the equations of dynamics, *Acta Mathematica* **13**, 1 (1890); On the three-body problem, *Bull. Astronomique* **8**, 12 (1891).
- [16] H. Poincaré, *Les Methodes Nouvelles de la Mecanique Celeste* (Gauthier-Villars, Paris, 1892), reprinted as *New Methods of Celestial Mechanics*, edited by D. L. Goroff (American Institute of Physics, College Park, 1993).
- [17] A. Einstein, L. Infeld, and B. Hoffman, The gravitational equations and the problem of motion, *Ann. Math.* **39**, 65 (1938).
- [18] V. A. Fock, On the motion of finite masses after the Einstein theory of gravitation, *J. Phys. (Moscow)* **1**, 81 (1939).
- [19] T. Levi-Civita, *The N-Body Problem in General Relativity* (Gauthier-Villars, Paris, 1941; Reidel, Dordrecht, 1964).
- [20] B. D. Tapley and J. M. Lewallen, Solar influence on satellite motion near the stable earth-moon libration points, *AIAA J.* **2**, 728 (1964).
- [21] L. A. Pars, *A Treatise on Analytical Dynamics* (Heinemann, London, 1965).
- [22] V. Szebehely, *Theory of Orbits: the Restricted Problem of Three Bodies* (Academic Press, New York, 1967).
- [23] E. Krefetz, Restricted three-body problem in the post-Newtonian approximation, *Astron. J.* **72**, 471 (1967).
- [24] V. A. Brumberg, *Relativistic Celestial Mechanics* (Nauka, Moscow, 1972).
- [25] R. A. Freitas and F. Valdes, A search for natural or artificial objects located at the Earth-Moon libration points, *Icarus* **42**, 442 (1980).
- [26] T. Damour, The problem of motion in Newtonian and Einsteinian gravity, in *300 Years of Gravitation*, eds. S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1987).
- [27] T. Damour and G. Schafer, Levi-Civita and the general relativistic problem of motion, in *Studies in the History of General Relativity, Einstein Studies Vol. 3*, eds. J. Eisenstaedt, A. J. Kox (Birkhauser, Boston, 1992).
- [28] A. Celletti and A. Giorgilli, On the stability of the Lagrangian points in the spatial restricted problem of three bodies, *Cel. Mech. Dyn. Astron.* **50**, 31 (1991).
- [29] T. Damour, M. Soffel, and C. Xu, General-relativistic celestial mechanics. I. Method and definition of reference systems, *Phys. Rev. D* **43**, 3273 (1991).
- [30] T. Damour, M. Soffel, and C. Xu, General-relativistic celestial mechanics. II. Translational equations of motion, *Phys. Rev. D* **45**, 1017 (1992).
- [31] T. Damour, M. Soffel, and C. Xu, General-relativistic celestial mechanics. III. Rotational equations of motion, *Phys. Rev. D* **47**, 3124 (1993).
- [32] T. Damour, M. Soffel, and C. Xu, General-relativistic celestial mechanics. IV. Theory of satellite motion, *Phys. Rev. D* **49**, 618 (1994).
- [33] T. I. Maindl and R. Dvorak, On the dynamics of the relativistic restricted three-body problem, *Astronomy & Astrophys.* **290**, 335 (1994).
- [34] T. I. Maindl, The solar system’s Lagrangian points in the framework of the relativistic restricted three-body problem, *ASP Conference Series* **107**, 147 (1996).
- [35] A. Celletti and L. Chierchia, On the stability of realistic three-body problems, *Commun. Math. Phys.* **186**, 413 (1997).
- [36] G. Dell’Antonio, Noncollision periodic solutions of the N-body system, *Nonlinear Differ. Equ. Appl.* **5**, 117 (1998).
- [37] K. B. Bhatnagar and P. P. Hallan, Existence and stability of $L_{4,5}$ in the relativistic restricted three-body problem, *Cel. Mech. Dyn. Astron.* **69**, 271 (1998).
- [38] C. D. Murray and S. F. Dermott, *Solar System Dynamics* (Cambridge University Press, Cambridge, 1999).
- [39] C. N. Douskos and E. A. Perdios, On the stability of equilibrium points in the relativistic restricted three-body problem, *Cel. Mech. Dyn. Astron.* **82**, 317 (2002).
- [40] A. Morbidelli, *Modern Celestial Mechanics: Aspects of Solar System Dynamics* (Taylor & Francis, London, 2002).
- [41] V. A. Brumberg, Special solutions in a simplified restricted three-body problem with gravitational radiation taken into

- account, *Cel. Mech. Dyn. Astron.* **85**, 269 (2003).
- [42] L. F. Wanex, Chaotic amplification in the relativistic restricted three-body problem, *Z. Naturforsch.* **58a**, 13 (2003).
 - [43] S. Kopeikin and I. Vlasov, The effacing principle in the post-Newtonian celestial mechanics, in *Proceedings XI Marcel Grossmann Meeting*, 2475-2477 (World Scientific, Singapore, 2008).
 - [44] H. Asada, Gravitational wave forms for a three-body system in Lagrange's orbit: Parameter determinations and a binary source test, *Phys. Rev. D* **80**, 064021 (2009).
 - [45] K. Yamada and H. Asada, Collinear solution to the general relativistic three-body problem, *Phys. Rev. D* **82**, 104019 (2010).
 - [46] H. Asada, T. Futamase and P. Hogan, *Equations of Motion in General Relativity*, Int. Ser. Monogr. Phys. **148** (Oxford University Press, Oxford, 2010).
 - [47] M. Connors, P. Wiegert, and C. Veillet, Earth's Trojan asteroid, *Nature (London)* **475**, 481 (2011).
 - [48] B. Bertotti, P. Farinella and D. Vokrouhlicky, *Physics of the Solar System: Dynamics and Evolution, Space Physics, and Spacetime Structure* (Springer, Berlin, 2012).
 - [49] K. Yamada and H. Asada, Triangular solution to the general relativistic three-body problem for general masses, *Phys. Rev. D* **86**, 124029 (2012).
 - [50] *List of Jupiter Trojans*, Minor Planet Center, 25 February 2014.
 - [51] K. Yamada and H. Asada, Post-Newtonian effects on the stability of the triangular solution in the three-body problem for general masses, *Phys. Rev. D* **91**, 124016 (2015).
 - [52] K. Yamada and H. Asada, Non-chaotic evolution of triangular configuration due to gravitational radiation reaction in the three-body problem, *Phys. Rev. D* **93**, 084027 (2016).
 - [53] T. Y. Zhou, W. G. Cao, and Y. Xie, Collinear solution to the three-body problem under a scalar-tensor gravity, *Phys. Rev. D* **93**, 064065 (2016).
 - [54] F. L. Dubeibe, F. D. Lora-Clavijo, and G. A. Gonzalez, Post-Newtonian circular restricted 3-body problem: Schwarzschild primaries, arXiv:1605.06204 [gr-qc].
 - [55] C. R. McInnes, *Solar Sailing: Technology, Dynamics and Mission Applications* (Springer Praxis, London, 1999).
 - [56] J. Simo and C. R. McInnes, Solar sail trajectories at the earth-moon Lagrange points, in *59th International Astronomical Congress, Glasgow, Scotland*, 2008.
 - [57] J. Simo and C. R. McInnes, Displaced periodic orbits with low-thrust propulsion in the earth-moon system, in *19th AAS/AIAA Space Flight Mechanics Meeting, Savannah, Georgia*, 2009.
 - [58] J. Simo and C. R. McInnes, Solar sail orbits at the earth-moon libration points, *Comm. Nonlinear Sci. Numer. Simulat.* **14**, 4191 (2009).
 - [59] J. Simo and C. R. McInnes, Asymptotic analysis of displaced lunar orbits, *J. of Guidance, Control, and Dynamics* **32**, 1666 (2009).
 - [60] J. Simo and C. R. McInnes, Displaced solar sail orbits: dynamics and applications, in *20th AAS/AIAA Space Flight Mechanics Meeting, San Diego, California*, 2010.
 - [61] J. Simo and C. R. McInnes, Designing displaced lunar orbits using low-thrust propulsion, *J. of Guidance, Control, and Dynamics* **33**, 259 (2010).
 - [62] J. Simo and C. R. McInnes, Feedback stabilization of displaced periodic orbits: application to binary asteroids, *Acta Astronautica* **96**, 106 (2014).
 - [63] P. L. Bender et al., The lunar experiment, *Science* **182**, 229 (1973).
 - [64] I. I. Shapiro, R. D. Reasenberg, J. F. Chandler, and R. W. Babcock, Measurement of the de Sitter precession of the moon: A relativistic three-body effect, *Phys. Rev. Lett.* **61**, 2643 (1988).
 - [65] M. R. Pearlman, J. J. Degnan, and J. M. Bosworth, The international laser ranging service, *Adv. Space Res.* **30**, 135 (2002).
 - [66] J. G. Williams, S. G. Turyshev, and D. H. Boggs, Progress in lunar laser ranging tests of relativistic gravity, *Phys. Rev. Lett.* **93**, 261101 (2004).
 - [67] Z. Altamimi, X. Collilieux, J. Legrand et al., ITRF2005: A new release of the international terrestrial reference frame based on time series of station positions and earth orientation parameters, *J. Geophys. Res.* **112**, B09401 (2007).
 - [68] R. March, G. Bellettini, R. Tauraso, and S. Dell'Agnello, Constraining spacetime torsion with the Moon and Mercury, *Phys. Rev. D* **83**, 104008 (2011).
 - [69] R. March, G. Bellettini, R. Tauraso, and S. Dell'Agnello, Constraining spacetime torsion with LAGEOS, *Gen. Relativ. Gravit.* **43**, 3099 (2011).
 - [70] S. Dell'Agnello et al., Creation of the new industry-standard space test of laser retroreflectors for the GNSS and LAGEOS, *J. Adv. Space Res.* **47**, 822 (2011).
 - [71] S. Dell'Agnello et al., Fundamental physics and absolute positioning metrology with the MAGIA lunar orbiter, *Exp. Astron.* **32**, 19 (2011).
 - [72] S. Dell'Agnello et al., Probing general relativity and new physics with lunar laser ranging, *Nucl. Instr. Methods Phys. Res. A* **692**, 275 (2012).
 - [73] M. Martini, S. Dell'Agnello et al., MoonLIGHT: A USA-Italy lunar laser ranging retroreflector array for the 21st century, *Planet. Space Sci.* **74**, 276 (2012).
 - [74] D. Currie, S. Dell'Agnello, G. O. Delle Monache, B. Behr, and J. G. Williams, A lunar laser ranging retroreflector array for the 21st century, *Nucl. Phys. B Proc. Suppl.* **243**, 218 (2013).
 - [75] S. Dell'Agnello et al., Next-generation laser retroreflectors for GNSS, solar system exploration, geodesy, gravitational physics and earth-observation, in *ESA Proc. Int. Conf. on Space Optics (Tenerife, Spain, Oct. 2014)*.

- [76] D. Vokrouhlicky, A note on the solar radiation perturbations of lunar motion, *Icarus* **126**, 293 (1997).
- [77] T. Levi Civita, *The Absolute Differential Calculus* (Blackie & Son, London, 1926; Dover, New York, 1977).
- [78] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, (W. H. Freeman and Company, San Francisco, 1973)
- [79] E. Battista, Extreme regimes in quantum gravity, PhD thesis (Naples University, 2016); arXiv:1606.04259.
- [80] A. Buonanno and T. Damour, Effective one-body approach to general relativistic two-body dynamics, *Phys. Rev. D* **59**, 084006 (1999).
- [81] A. Buonanno and T. Damour, Transition from inspiral to plunge in binary black hole coalescences, *Phys. Rev. D* **62**, 064015 (2000).
- [82] A. Buonanno, Y. Chen, and T. Damour, Transition from inspiral to plunge in precessing binaries of spinning black holes, *Phys. Rev. D* **74**, 104005 (2006).
- [83] T. Damour, A. Nagar, M. Hannam, S. Husa, and B. Bruegmann, Accurate effective-one-body waveforms of inspiralling and coalescing black-hole binaries, *Phys. Rev. D* **78**, 044039 (2008).
- [84] T. Damour and A. Nagar, An improved analytical description of inspiralling and coalescing black-hole binaries, arXiv:0902.0136.
- [85] T. Damour, B. R. Iyer, and A. Nagar, Improved resummation of post-Newtonian multipolar waveforms from circularized compact binaries, *Phys. Rev. D* **79**, 064004 (2009).
- [86] D. Bini, T. Damour, and A. Geralico, Confirming and improving post-Newtonian and effective-one-body results from self-force computations along eccentric orbits around a Schwarzschild black hole, *Phys. Rev. D* **93**, 064023 (2016).
- [87] D. Bini, T. Damour, and A. Geralico, New gravitational self-force analytical results for eccentric orbits around a Schwarzschild black hole, *Phys. Rev. D* **93**, 104017 (2016).
- [88] T. Damour and D. Bini, Conservative second-order gravitational self-force on circular orbits and the effective one-body formalism, *Phys. Rev. D* **93**, 104040 (2016).
- [89] D. Bini, T. Damour, and A. Geralico, High post-Newtonian order gravitational self-force analytical results for eccentric equatorial orbits around a Kerr black hole, *Phys. Rev. D* **93**, 124058 (2016).
- [90] Y. Fourès-Bruhat, Théorème d'existence pour certains systèmes d'équations aux dérivées partielles non linéaires, *Acta Mathematica* **88**, 141 (1952); English translation in the Max Planck Institute for History of Science preprint series, document 480 (2016).
- [91] H. Poincaré, *Collected Works*, Vol. 1 (Gauthier-Villars, Paris, 1928; Jacques Gabais, Paris, 2011).
- [92] G. Valiron, *Functional Equations. Applications* (Masson, Paris, 1945).