




Publication Year	2017
Acceptance in OA	2020-08-21T11:55:14Z
Title	Activity and rotation of the X-ray emitting Kepler stars
Authors	STELZER, BEATE, Pizzocaro, D., PORETTI, Ennio, MICELA, Giuseppina, Belfiore, A., Marelli, M., Salvetti, D., DE LUCA, Andrea
Handle	http://hdl.handle.net/20.500.12386/26755



Activity and rotation of the X-ray emitting *Kepler* stars

Daniele Pizzocar^{*1,2}, Beate Stelzer^{3,4}, Ennio Poretti⁵, Giuseppina Micela⁴, Andrea Belfiore¹,

David Salvetti¹, Martino Marelli¹, Andrea De Luca¹

^{*}pizzocar@lambrate.inaf.it

¹INAF-IASF Milano, Italy

²Università degli Studi dell'Insubria, Varese, Italy

³Eberhard-Karls Universität Tübingen, Tübingen, Germany

⁴Osservatorio Astronomico di Palermo, Palermo, Italy

⁵INAF - Osservatorio Astronomico di Brera, via E. Bianchi 46, 23807 Merate (LC), Italy, Italy




ABSTRACT

The relation between magnetic activity and rotation in late-type stars provides fundamental information on stellar dynamos and spin evolution. Previous studies suffer from bias (spectroscopic rotation periods) and inhomogeneity (different instruments/techniques). **We characterize the activity/rotation relation of the X-ray emitting main-sequence late-type stars observed by XMM-Newton and Kepler.** We measure photometric rotation periods from the high-precision Kepler light curves. As activity indicators, we adopt the X-ray luminosity from XMM-Newton, the number/frequency of white-light and X-ray flares (from the light curves produced in the EXTrAS Project [1]).

This is the first analysis of the activity/relation link across the whole late-type spectral range performed using homogeneous photometric rotation periods from high precision Kepler light curves and homogeneous X-ray data. Our results reinforce the current picture of the rotation/activity relation and give important insights into the relation between X-ray activity and rotation and between X-ray and white-light flaring activity.

KEPLER

High photometric precision optical light curves
High sampling rate (29.4 min)
Large sample of stars (~170,000)
Kepler Input Catalogue: collection of photometry



SCIENTIFIC RATIONALE

ANALYSIS OF THE ACTIVITY/ROTATION RELATION IN
LATE-TYPE MAIN-SEQUENCE STARS

PHOTOMETRIC ROTATION PERIODS FROM KEPLER

MAIN ACTIVITY INDICATORS:

- X-RAYS FROM 3XMM-DR5 CATALOGUE;

- WHITE-LIGHT AND X-RAY FLARING

-> HOMOGENEITY IN ROTATION PERIOD & ACTIVITY MEASURE

DATA ANALYSIS

Sample: 107 late-type main-sequence stars detected in 3XMM-DR5 and with light curve in Kepler database


Kepler light curves: brightness modulation due to spots on the rotating stellar surface. Automated algorithm based on Autocorrelation function (ACF) & Lomb-Scargle Periodogram;

Rotation period, Rossby number, white-light flares

X-ray activity: X-ray luminosity from 3XMM-DR5 count rate 0.2-2.0 keV; thermal model $kT=0.82$ keV (from spectral fit of brightest sources). Search for flares in EXTrAS[1] light curves

XMM-NEWTON

3XMM-DR5 catalogue [2]: largest X-ray source catalogue (~390,000)
16 XMM-Newton observations in the Kepler field of view.



RESULTS

First characterization of the activity/rotation link on the whole late-type main-sequence spectral range with rotation periods measured from high-precision Kepler light curves and homogeneous measure of activity from XMM-Newton data and Kepler light curves.

Rotation periods: 76 stars (72%) present rotational variability -> rotation period in the range ~0.3-70 d. Very high fraction of rotators, due to selection bias towards very active stars. Twenty-nine new rotators.

X-ray activity/rotation: X-rays flux-limited sample + large average distance of Kepler stars (~270 pc) -> bias towards very active stars (see comparison with the distribution by [3], 90% completeness).

X-ray activity decreases with period; hints of two regimes: correlation for slow rotators, saturation for fast rotators. Good agreement with previous results [4] (K2 sample of M stars within 10 pc), [5], [6].

Flaring activity: white-light flaring rate increases towards later spectral types; possible decrease towards long rotation periods. Six X-ray flares: all with chromospheric long-duration counterpart. Consistent X-ray/white-light flaring rate. Good agreement with [4]

