



|                                  |                                                                                                 |
|----------------------------------|-------------------------------------------------------------------------------------------------|
| <b>Publication Year</b>          | 2015                                                                                            |
| <b>Acceptance in OA</b>          | 2020-04-16T17:18:49Z                                                                            |
| <b>Title</b>                     | Are jet ubiquitous in ULXs?                                                                     |
| <b>Authors</b>                   | WOLTER, Anna Luisa Maria, GHISELLINI, Gabriele, Costamante, L., Pintore, Fabio                  |
| <b>Publisher's version (DOI)</b> | 10.1017/S1743921315002549                                                                       |
| <b>Handle</b>                    | <a href="http://hdl.handle.net/20.500.12386/24082">http://hdl.handle.net/20.500.12386/24082</a> |
| <b>Serie</b>                     | PROCEEDINGS OF THE INTERNATIONAL ASTRONOMICAL UNION                                             |
| <b>Volume</b>                    | vol. 10, Issue S313                                                                             |

# Are jet ubiquitous in ULXs?

A. Wolter<sup>1</sup>, G. Ghisellini<sup>1</sup>, L. Costamante<sup>2</sup> and F. Pintore<sup>3</sup>

<sup>1</sup>INAF-Osservatorio Astronomico di Brera, Milano, Italy; email: [anna.wolter@brera.inaf.it](mailto:anna.wolter@brera.inaf.it)

<sup>2</sup>Università di Perugia, Italy, <sup>3</sup>Università di Cagliari, Italy

**Abstract.** A jet model represents well different X-ray states of the bright ULX IC342 X-1.

**Keywords.** galaxies: individual (IC 342), X-ray: galaxies, acceleration of particles

## 1. Introduction


Ultra Luminous X-ray sources (ULXs) are point-like, off-nuclear X-ray sources with inferred luminosity (if assumed to be isotropic) in excess of the Eddington limit for spherical accretion onto a  $10 M_{\odot}$  compact object (see e.g., Fabbiano 2006, for a review). IC 342 is a nearby starburst galaxy (3.9 Mpc; Tikhonov & Galazutdinova 2010) that contains two ULXs: X-1 and X-2, the first ULXs reported to transition between distinct spectral states (Kubota *et al.* 2001): X-1 went from an apparently disc-dominated state to a power-law-dominated state while the spectrum of X-2 coincidentally carried out the reverse transition. The source X-1 is surrounded by a large optical and radio nebula that may be an X-ray ionized bubble driven by strong outflows from the ULX (Pakull & Mirioni 2002; Roberts *et al.* 2003; Feng & Kaaret 2008). Radio observations of the bubble have been performed with the JVLA at 5 GHz (Cseh *et al.* 2012): an extended source is detected, with about the same extent of the optical bubble, and a possible unresolved source, albeit not confirmed by Marlowe *et al.* (2014). The upper limit of the point source is 14 mJy at 5.5 GHz.

Could the properties of the ULX X-1 be explained by the presence of a jet?

## 2. Results

We have collected spectral measures at all the available wavelengths in order to construct an overall spectral energy distribution (SED). This SED of IC 342 X-1 is plotted in Figure 1: *left*. In order to test the jet hypothesis we first compare the SED with a known jet source, the Galactic microquasar GRS 1915+105, known to be a superluminal (Mirabel & Rodriguez 1994) and complexly variable (Fender & Belloni 2004 for a review) bright source. To mimic this source as pointing towards our line of sight, at an angle of  $3^{\circ}$ , we boosted the X-ray and radio power by a factor  $\delta^4$  with  $\delta = 2.8$ . Given the strong agreement we deem that a jet is a viable description of the X-ray emission of this ULX.

We use a model (see Ghisellini & Tavecchio 2009, for details) that has  $M_{\text{BH}} = 50 M_{\odot}$  and  $L_{\text{Edd}} = 9 \times 10^{38}$  erg/s. The disk luminosity is  $L_{\text{disk}} = 12\%$  of  $L_{\text{Edd}}$ , while the jet, at  $100 R_{\text{S}}$ , has an angle to the line-of-sight  $\theta_{\text{v}} = 3^{\circ}$  and  $\Gamma_{\text{bulk}} = 4$ . The magnetic field is  $B = 2 \times 10^6$  G. We include also a corona at 30% of the disk luminosity and compare it with the SED and in particular with the different X-ray observations from 2004 (see Figure 1: *right*). The overall X-ray emission is well described by this model in the optical and X-ray energy range; the model never exceeds the upper limit from Fermi in the TeV band. Different X-ray states of IC 342 X-1 can be explained by changing only the injected power (lowest = 30% of highest) and slightly the electron distribution.



**Figure 1.** *Left:* The SED of IC342 X-1 in radio, optical, and 95% upper limit in the gamma-ray band (red). Fermi data are the upper limit front Aug 2008-Apr 2014 data version (*P7.REP*; SOURCE class events; Galactic diffuse model *v05\_rev1*;  $\Gamma=2$  in each of the 4 bands (Atwood *et al.* 2009)). The various datasets for XMM-Newton and Swift are represented in pink, blue, pale blue and gray. NuStar data in black. See Pintore *et al.* (2014); Rana *et al.* (2014) and references therein. The comparison SED of GRS 1915+105 is shown in green. *Right:* Zoom of the X-ray band with the models describing different states.

Another Galactic analogue of such a system is the W50 nebula, thought to be inflated by the relativistic jets of the microquasar SS433 (Dubner *et al.* 1998), which has similar energetic requirements to the IC342 X-1 nebula. Other indications of the presence of jets in ULXs: NGC 2276 3c/S6 (Mezcua *et al.* 2013; Wolter *et al.* 2014) with two lobes of radio emission; Holmberg II X-1 (Cseh *et al.* 2014) with an apparent triple radio structure and attributed to a BH with  $M_{\text{BH}} \geq 25M_{\odot}$ . The interpretation with a jet of the X-ray emission is worth of further investigation both with new multi-wavelength observations of this source and the analysis of other ULXs with radio detections.

## Acknowledgements

We acknowledge partial funding from ASI - INAF I/037/12/0.

## References

- Atwood, W. B. *et al.* 2009 *ApJ* 697, 1071  
 Cseh, D., Corbel, S., Kaaret, P., *et al.* 2012 *ApJ* 749, 17  
 Cseh, D., Kaaret, P., Corbel, S., *et al.* 2014 *MNRAS* 439, L1  
 Dubner, G. M., Holdaway, M., Goss, W. M., & Mirabel, I. F. 1998 *AJ* 116, 1842  
 Fabbiano, G. 2006, *ARAA* 44, 323  
 Feng, H., & Kaaret, P. 2008 *ApJ* 675, 1067  
 Ghisellini, G., & Tavecchio, F. 2009 *MNRAS* 397, 985  
 Kubota, A., Mizuno, T., Makishima, K., *et al.* 2001 *ApJ* 547, L119  
 Marlowe, H., Kaaret, P., Lang, C., *et al.* 2014 *MNRAS* 444, 642  
 Mezcua, M., Roberts, T. P., Sutton, A. D., & Lobanov, A. P. 2013, *MNRAS* 436, 3128  
 Pakull, M. W. & Mirioni, L. 2002, arXiv:astro-ph/0202488  
 Pintore F., Zampieri L, Wolter, A., & Belloni, T. 2014 *MNRAS* 439, 3461  
 Rana, V., *et al.* 2014 *ApJ* in press, arXiv:1401.4637  
 Roberts, T. P., Goad, M. R., Ward, M. J., & Warwick, R. S. 2003 *MNRAS* 342, 709  
 Tikhonov, N. A. & Galazutdinova, O. A. 2010 *Astron. Lett.* 36, 167  
 Wolter, A., Mapelli, M., Esposito, P., Pizzolato, F., & Ripamonti, E. 2014 *MNRAS* submitted