



|                                  |                                                                                                                           |
|----------------------------------|---------------------------------------------------------------------------------------------------------------------------|
| <b>Publication Year</b>          | 2015                                                                                                                      |
| <b>Acceptance in OA</b>          | 2020-05-11T12:09:17Z                                                                                                      |
| <b>Title</b>                     | Search for hydrogen peroxide in the Martian atmosphere by the Planetary Fourier Spectrometer onboard Mars Express         |
| <b>Authors</b>                   | Aoki, Shohei, GIURANNA, MARCO, Kasaba, Yasumasa, Nakagawa, Hiromu, SINDONI, Giuseppe, Geminale, Anna, Formisano, Vittorio |
| <b>Publisher's version (DOI)</b> | 10.1016/j.icarus.2014.09.034                                                                                              |
| <b>Handle</b>                    | <a href="http://hdl.handle.net/20.500.12386/24686">http://hdl.handle.net/20.500.12386/24686</a>                           |
| <b>Journal</b>                   | ICARUS                                                                                                                    |
| <b>Volume</b>                    | 245                                                                                                                       |

**Search for hydrogen peroxide in the Martian atmosphere by the Planetary  
Fourier Spectrometer onboard Mars Express**

Shohei Aoki <sup>a,b,\*</sup>, Marco Giuranna <sup>b</sup>, Yasumasa Kasaba <sup>a</sup>, Hiromu Nakagawa <sup>a</sup>, Giuseppe  
Sindoni <sup>b</sup>, Anna Geminale <sup>b</sup>, and Vittorio Formisano <sup>b</sup>

<sup>a</sup>Department of Geophysics, Graduate School of Science, Tohoku University, Sendai, Miyagi  
980-8578, Japan

<sup>b</sup> Istituto di Astrofisica e Planetologia Spaziali (IAPS), Istituto Nazionale di Astrofisica  
(INAF), Via del Fosso del Cavaliere 100, 00133 Roma, Italy

## 1    **Abstract**

2    We searched for hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) in the Martian atmosphere using data measured by  
3    the Planetary Fourier Spectrometer (PFS) onboard Mars Express during five Martian years  
4    (MY27–31). It is well known that  $\text{H}_2\text{O}_2$  plays a key role in the oxidizing capacity of the  
5    Martian atmosphere. However, only a few studies based on ground-based observations can be  
6    found in the literature. Here, we performed the first analysis of  $\text{H}_2\text{O}_2$  using long-term  
7    measurements by a spacecraft-borne instrument. We used the  $\nu_4$  band of  $\text{H}_2\text{O}_2$  in the spectral  
8    range between  $359\text{ cm}^{-1}$  and  $382\text{ cm}^{-1}$  where strong features of  $\text{H}_2\text{O}_2$  are present around  $362$   
9     $\text{cm}^{-1}$  and  $379\text{ cm}^{-1}$ . Since the features were expected to be very weak even at the strong band,  
10    sensitive data calibrations were performed and a large number of spectra were selected and  
11    averaged. We made three averaged spectra for different seasons over relatively low latitudes  
12    ( $50^\circ\text{S}$ – $50^\circ\text{N}$ ). We found features of  $\text{H}_2\text{O}_2$  at  $379\text{ cm}^{-1}$ , whereas no clear features were detected  
13    at  $362\text{ cm}^{-1}$  due to large amounts of uncertainty in the data. The derived mixing ratios of  $\text{H}_2\text{O}_2$   
14    were close to the detection limits:  $16 \pm 19$  ppb at  $L_s = 0$ – $120^\circ$ ,  $35 \pm 32$  ppb at  $L_s = 120$ – $240^\circ$ ,  
15    and  $41 \pm 28$  ppb at  $L_s = 240$ – $360^\circ$ . The retrieved value showed the detection of  $\text{H}_2\text{O}_2$  only for  
16    the third seasonal period, and the values in the other periods provided the upper limits. These  
17    long-term averaged abundances derived by the PFS generally agreed with the ones reported by  
18    ground-based observations. From our derived mixing ratio of  $\text{H}_2\text{O}_2$ , the lifetime of  $\text{CH}_4$  in the  
19    Martian atmosphere is estimated to be several decades in the shortest case. Our results and

sporadic detections of CH<sub>4</sub> suggest the presence of strong CH<sub>4</sub> sinks not subject to atmospheric oxidation.

## 1. Introduction

The atmosphere of Mars is strongly oxidized. Hydrogen radicals (H, OH, HO<sub>2</sub>) play a key role in regulating photochemistry in the Martian atmosphere, especially the oxidizing capacity. Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) represents a temporal reservoir of hydrogen radicals, and it is the best tracer of oxidizers in the atmosphere because the abundance is relatively high. Presently, only a few ground-based observations of H<sub>2</sub>O<sub>2</sub> can be found in the literature (e.g., Encrenaz et al., 2012). **Table 1** lists a summary of the H<sub>2</sub>O<sub>2</sub> mixing ratios derived from these observations. The observed values are distributed in the range from 0 to 50 ppb, and the maximum abundance is observed during the northern autumn ( $L_s = 206^\circ$ ).

Understanding of oxidizing capacity in the Martian atmosphere is important to constrain sinks of CH<sub>4</sub> on Mars. The spatial and temporal variation of CH<sub>4</sub> observed by ground-based and space-borne instruments has led to much discussion on its sources and sinks (Formisano et al., 2004; Mumma et al., 2009; Fonti and Marzo, 2010; Geminale et al., 2011). The observations to date suggest that the lifetime of CH<sub>4</sub> in the Martian atmosphere is on the order of weeks or months. In contrast, photochemical models have predicted that the photochemical lifetime of CH<sub>4</sub> is 670 years (Wong et al., 2003), and the required loss rate of CH<sub>4</sub> is estimated

1 to be about 600 times faster than that predicted by the standard photochemical model (Lefèvre  
2 and Forget, 2009). Since the main process that destroys CH<sub>4</sub> is oxidation by OH, long-term  
3 averaged characteristics of H<sub>2</sub>O<sub>2</sub> from Martian orbiters can provide further constraints on  
4 oxidant loss of CH<sub>4</sub> in the Martian atmosphere.

5 The reason why H<sub>2</sub>O<sub>2</sub> has not been detected by any instruments onboard Martian orbiters  
6 is that the spectral resolutions were insufficient to distinguish the weak absorption lines of  
7 H<sub>2</sub>O<sub>2</sub> from the stronger ones of CO<sub>2</sub> and H<sub>2</sub>O. The Planetary Fourier Spectrometer (PFS)  
8 onboard Mars Express (MEX) has relatively high spectral resolution ( $\sim 1.3 \text{ cm}^{-1}$  without  
9 apodization) in the spectral range between 250 and 8300  $\text{cm}^{-1}$ . Mars Express has been  
10 operating for ten years in Martian orbit since its launch in 2003, and the PFS has successfully  
11 obtained a large amount of observational spectra. This dataset enables us to perform a  
12 sensitive search for H<sub>2</sub>O<sub>2</sub>. Indeed, Encrenaz et al. (2012) pointed out that the PFS has the  
13 potential to distinguish the weak absorption lines of H<sub>2</sub>O<sub>2</sub> in the far-infrared range (350–400  
14  $\text{cm}^{-1}$ ). In this paper, we attempted to perform detections of H<sub>2</sub>O<sub>2</sub> from spacecraft-borne  
15 measurements for the first time with the PFS data.

## 16 17 **2. Dataset and reduction**

18 The PFS is a Fourier transform spectrometer that was mainly designed for atmospheric  
19 analyses such as measurements of the 3-D temperature field in the lower atmosphere,

variations of minor constituents, and optical properties of aerosols (Formisano et al., 2005). It has two spectral channels: one is the Short Wavelength Channel (SWC) and the other is the Long Wavelength Channel (LWC). The SWC covers the spectral range between 1750 and 8200  $\text{cm}^{-1}$ , and the LWC covers the spectral range between 250 and 1700  $\text{cm}^{-1}$ ; both channels have a sampling step of 1.0  $\text{cm}^{-1}$ . The spectral resolution is 1.3  $\text{cm}^{-1}$  without apodization and 2  $\text{cm}^{-1}$  when a Hamming function is applied to the interferograms. One of the advantages of the PFS is its wide spectral coverage with relatively high spectral resolution. An interferogram can be obtained in about 4 s, with a repetition interval of 8.5 s. Its field of view is  $1.6^\circ$  for the SWC and  $2.8^\circ$  for the LWC. In the period from January 2004 to August 2013, the PFS has obtained about two million spectra of Mars. These cover almost all the regions of Mars for different seasons and local times.

This study used the LWC data in the spectral range between 350 and 400  $\text{cm}^{-1}$  (25.0–28.5  $\mu\text{m}$ ), which includes the strong absorption lines of  $\text{H}_2\text{O}_2$  and  $\text{H}_2\text{O}$ . This spectral range does not have strong lines of  $\text{CO}_2$ , and dust and water ice features are expected to be very weak in this range (Smith et al., 2000). The previous observations have never used this spectral range for studies of  $\text{H}_2\text{O}_2$  because of the strong  $\text{H}_2\text{O}$  absorptions in the terrestrial atmosphere. **Figure 1** shows the line strengths of  $\text{H}_2\text{O}_2$  and  $\text{H}_2\text{O}$  obtained from the HITRAN08 database (Rothman et al., 2009), and the synthetic spectra for the spectral resolution of the PFS (2.0  $\text{cm}^{-1}$ ) with various mixing ratios of  $\text{H}_2\text{O}$  and  $\text{H}_2\text{O}_2$  (details of the model are described in Section 3). As

1 pointed out in Encrenaz et al. (2012), the  $\nu_4$  band of  $\text{H}_2\text{O}_2$  exhibits the strongest transition in  
2 the considered spectral range and the PFS can detect  $\text{H}_2\text{O}_2$  absorptions, although these are  
3 very weak, at 362 and 379  $\text{cm}^{-1}$ . In order to increase the signal to noise ratio (SNR), we  
4 averaged thousands of spectra. The noise equivalent radiance (NER) of the PFS/LWC in the  
5 considered spectral range (360–385  $\text{cm}^{-1}$ ) is around 1  $\text{erg}/(\text{s sr cm}^2 \text{ cm}^{-1})$  (Giuranna et al.,  
6 2005a). The measured radiance in this spectral range is typically  $\sim 90 \text{ erg}/(\text{s sr cm}^2 \text{ cm}^{-1})$  at the  
7 daytime low-latitude regions used in this study (i.e.,  $\text{SNR} = 90$  on a single spectrum). Thus,  
8 the SNR reaches the order of 1000 by averaging thousands of spectra. In addition, we  
9 performed sensitive calibrations in order to increase confidence in the tiny features of  $\text{H}_2\text{O}_2$ .  
10 The PFS/LWC calibration has been presented in Giuranna et al. (2005a). Based on that, we  
11 further improved the radiometric calibration. In-flight radiometric calibration uses the  
12 measurements of an internal blackbody and deep space. The temperatures inside the  
13 interferometer are very stable (Giuranna et al., 2005a), and they have remained stable during  
14 the 10 years of PFS operations. We thus collected all the available calibration measurements  
15 to derive the spectral responsivity and thermal emission of the instrument with a high degree  
16 of confidence, and then we performed a re-calibration of the whole dataset using the new  
17 accurate curves. Moreover, correction of asymmetric interferograms (phase correction) was  
18 also improved. As shown in Giuranna et al. (2005a), the PFS produces “double-sided”  
19 interferograms; owing to phase errors, all original interferograms have an asymmetric form.

1 So far, the asymmetry of the central parts of the PFS interferograms were corrected by means  
2 of a specially developed algorithm based on the method of phase correction suggested by  
3 Forman et al. (1966). Recently, a new and improved algorithm has been developed and  
4 applied to the PFS interferograms (Saggin et al., 2011). The implemented method is based on  
5 the identification of an instrumental phase that is dependent on the interferometer temperature  
6 and on the extraction of a linear phase component through a least-square approach. The use of  
7 an instrumental phase parameterized with the interferometer temperature eases the  
8 determination of the linear phase that can be extracted using only a narrow spectral region  
9 selected to be immune from disturbances (Saggin et al., 2011). The procedure has been  
10 intensively tested and successfully applied to whole datasets of the PFS. As a final remark, we  
11 applied Hamming apodization to the interferograms used in this analysis. Although this  
12 procedure slightly reduces spectral resolution, it significantly reduces side-lobes in the  
13 instrumental line shape (Giuranna et al., 2005b).

14 In order to search for  $\text{H}_2\text{O}_2$ , we collected three averaged spectra for different seasonal  
15 ranges:  $L_s = 0\text{--}120^\circ$ ,  $L_s = 120\text{--}240^\circ$ , and  $L_s = 240\text{--}360^\circ$ . We used the data collected during  
16 the period from January 2004 to October 2011, which corresponds to five Martian years,  
17 namely MY27 to MY31. Only nadir observations (emission angle less than  $10^\circ$ ) were selected  
18 for the analysis. In order to obtain a sufficient background radiance and reduce the uncertainty  
19 due to averaging various atmospheric conditions, we built a reduced dataset where the latitude

is between  $-50^\circ$  and  $+50^\circ$ , local time is between 10 h and 16 h (dayside), and surface temperature is between 250 K and 270 K. Finally, to further increase the homogeneity of selected data, we only considered spectra acquired for fixed instrumental parameters such as the spectral step, detector temperature, and position of the peak in the interferogram. In total, the numbers of co-added spectra in the three averages are 20,260, 7477, and 13,777 for  $L_s = 0-120^\circ$ ,  $L_s = 120-240^\circ$ , and  $L_s = 240-360^\circ$ , respectively. In principle, the SNRs for the three datasets reach to about 10,500, 6500, and 8600. **Figure 2** shows the distributions of solar longitude  $L_s$  in the three datasets. Note that about 90% of the spectra in the first average were acquired at  $L_s = 0-60^\circ$ , and about 80% of the spectra in the third average were acquired at  $L_s = 330-360^\circ$ .

### 3. Method of analysis

We retrieved the mixing ratio of  $H_2O_2$  from comparisons between synthetic spectra and the averaged PFS data in the spectral range between 359 and  $382\text{ cm}^{-1}$ . For calculation of the synthetic spectra, we developed a radiative transfer model for a plane-parallel and clear-sky atmosphere by the line-by-line method. Radiance  $I(\nu)$  is calculated by

$$I(\nu) = \varepsilon(\nu)B(T_{surf}, \nu)e^{-\tau_n} + \sum_{i=1}^n \left( (1 - e^{-(\tau_i - \tau_{i-1})}) B(T_{i-1/2}, \nu) e^{-\tau_{i-1}} \right),$$

where  $\nu$  is the wavenumber,  $\varepsilon(\nu)$  is the surface emissivity, and  $\tau_i$  is the optical depth at the  $i$ -th layer (Hanel et al., 2003).  $B(T_{surf}, \nu)$  and  $B(T_{i-1/2}, \nu)$  are the blackbody radiations at the Martian

1 surface temperature  $T_{surf}$  and the averaged temperature between the  $i$ -th and  $i-1$ -th atmospheric  
2 layer, respectively. In this study,  $H_2O$  and  $H_2O_2$  were considered as gaseous absorptions. The  
3 surface emissivity  $\varepsilon(\nu)$  was modeled as  $\varepsilon(\nu) = \alpha\nu + \beta$ , where  $\alpha$  and  $\beta$  were constant parameters  
4 (Smith et al., 2000; Fouchet et al., 2007). The spectroscopic parameters of the  $H_2O_2$  lines were  
5 obtained from the HITRAN08 database (Rothman et al., 2009), and those of  $H_2O$  lines were  
6 obtained from the water vapor line list for a  $CO_2$ -rich atmosphere  
7 (<http://astrobiology.gsfc.nasa.gov/Villanueva/spec.html>). This database has the line shape  
8 parameters (e.g., air-broadening coefficients) that were calculated for a  $CO_2$ -rich atmosphere  
9 following the complex Robert Bonamy formalism based on the HITRAN08 database  
10 (Villanueva et al., 2012). The terrestrial air-broadening coefficients of  $H_2O_2$  are multiplied by  
11 a factor of 1.5 uniformly in order to correct these values for the  $CO_2$ -rich atmosphere of Mars.  
12 Temperature correction of the line intensity and temperature/pressure correction of the line  
13 half-width were performed according to Rothman et al. (1998). The ratio of the total internal  
14 partition function, which is required for the corrections mentioned above, was calculated using  
15 the “TIPS” routine provided in the HITRAN08 database. The line shape function is assumed  
16 to be the Voigt function (Drayson, 1976). The vertical profiles of temperature, pressure, and  
17  $H_2O$  were obtained from the Mars Climate Database (MCD) version 4.3  
18 (<http://www-mars.lmd.jussieu.fr>). We extracted the profiles for individual observed  
19 geometries and averaged them. The vertical profile of the  $H_2O_2$  mixing ratio was assumed to

be uniform from 0 to 20 km (Encrenaz et al., 2012). The surface temperature was determined from the averaged PFS spectrum between 300 and 500  $\text{cm}^{-1}$ .

The synthetic spectrum to be compared with the PFS spectrum was obtained by a convolution between the radiance  $I(\nu)$  and the Instrumental Line Shape (ILS) of the PFS. The ILS was obtained from the calibration measurements in the laboratory (Giuranna et al., 2005b). As shown in Figure 9 of Giuranna et al. (2005b), the ILS has a weak side-lobe around  $-3 \text{ cm}^{-1}$ , and its amplitude has a certain uncertainty. Since the effect of the side-lobe cannot be neglected during comparisons to the absorption depth of  $\text{H}_2\text{O}_2$ , we set the amplitude as a variable parameter in the retrieval process. In addition, we also set the spectral grid of the ILS (i.e., spectral resolution) as a variable parameter because the peak position of the interferogram varies slightly for different data, i.e., the spectral resolution should be slightly modified.

In the retrieval process, we derived eight parameters: surface emissivity ( $\alpha$  and  $\beta$ ), the scaling factor for the side-lobe of the ILS, the scaling factor for the spectral grid of the ILS, the scaling factor for spectral grid of PFS, the scaling factor for the temperature profile, and the mixing ratios of  $\text{H}_2\text{O}$  and  $\text{H}_2\text{O}_2$ . The scaling factor for the spectral grid of the PFS was introduced because it is slightly shifted due to small temporal variations of the laser diode temperature (Sindoni et al., 2011). The scaling factor for the temperature profile was introduced because the water vapor lines are saturated and sensitive to the temperature profile.

Our retrieval was performed as follows: we prepared a parameter domain for the mixing ratio of H<sub>2</sub>O<sub>2</sub> from 0 to 100 ppb with intervals of 1 ppb. Then, we retrieved the other seven parameters using the Levenberg–Marquardt non-linear minimization algorithm. Finally, we derived the mixing ratio of H<sub>2</sub>O<sub>2</sub> that provides the minimum values of the cost function  $\chi^2$ .

The cost function  $\chi^2$  is defined as

$$\chi^2 = \frac{1}{N - M} \sum_{i=1}^N \left( \frac{y(v_i) - I(v_i)}{\sigma(v_i)} \right)^2,$$

where  $v_i$  is  $i$ -th spectral point,  $N$  is the number of spectral points in the considered range (i.e.,  $N = 23$ ),  $M$  is the number of parameters (i.e.,  $M = 7$ ),  $\sigma(v_i)$  is the noise of the averaged PFS spectra,  $y(v_i)$  is the averaged PFS spectrum, and  $I(v_i)$  is the synthetic spectrum.

#### 4. Results and discussion

**Figure 3** shows the averaged PFS spectra for the three different seasons. In the upper panels of the figure, the averaged PFS spectra are compared to the synthetic spectra. The synthetic spectra successfully reproduce the features of water vapor around 370 cm<sup>-1</sup> and 375 cm<sup>-1</sup>. The best-fit synthetic spectra show that the column densities of water vapor are  $2.43 \times 10^{19}$  cm<sup>-2</sup> at  $L_s = 0$ –120°,  $3.52 \times 10^{19}$  cm<sup>-2</sup> at  $L_s = 120$ –240°, and  $2.50 \times 10^{19}$  cm<sup>-2</sup> at  $L_s = 240$ –360°, which correspond to 7.25 pr-μm, 10.51 pr-μm, and 7.46 pr-μm, respectively. The absolute values and the enhancement during the summer–autumn season are consistent with the previous reports by spacecraft-borne observations (e.g., Smith, 2002). In the bottom panels of **Figure 3**, we

show residual spectra, where special synthetic spectra with all contributions except H<sub>2</sub>O<sub>2</sub> have been subtracted from the measured and synthetic spectra. The best-fit synthetic spectra show that the mixing ratios of H<sub>2</sub>O<sub>2</sub> are 16 ppb at  $L_s = 0\text{--}120^\circ$ , 35 ppb at  $L_s = 120\text{--}240^\circ$ , and 41 ppb at  $L_s = 240\text{--}360^\circ$ . As shown in the figures, H<sub>2</sub>O<sub>2</sub> band-like features are clearly visible around 379 cm<sup>-1</sup>. However, they were not detected around 362 cm<sup>-1</sup>.

The black vertical bars shown in **Figure 3** represent the errors estimated from the noise equivalent radiance divided by the square root of the number of measurements. However, discrepancies between the synthetic spectra and the measured ones imply that the error bars were underestimated. We think that this underestimation is due to the average of a number of spectra that were taken under a wide range of different conditions (i.e., temperatures, water vapor abundances, etc.). In order to evaluate realistic instrumental noises, we calculated the standard deviation (1-sigma) of the residuals between the PFS data and best-fit synthetic spectra. The red vertical bars shown in **Figure 3** represent the standard deviation. The SNRs based on the standard deviation were about 3770 for the dataset averaged over  $L_s = 0\text{--}120^\circ$ , 2630 for  $L_s = 120\text{--}240^\circ$ , and 4330 for  $L_s = 240\text{--}360^\circ$ . The uncertainties in the retrieved H<sub>2</sub>O<sub>2</sub> mixing ratio owing to the instrumental noise ( $\sigma_n$ ) were derived by comparisons between the absorption depths of H<sub>2</sub>O<sub>2</sub> and the retrieved standard deviation. The values are  $\sigma_n = 19$  ppb at  $L_s = 0\text{--}120^\circ$ ,  $\sigma_n = 31$  ppb at  $L_s = 120\text{--}240^\circ$ , and  $\sigma_n = 27$  ppb at  $L_s = 240\text{--}360^\circ$ . In addition to the instrumental noise, uncertainties in the synthetic spectrum are the other sources of errors.

The dominant one among them is uncertainty in the line strength of H<sub>2</sub>O<sub>2</sub> provided by the HITRAN08 database, which is 10% at maximum (Rothman et al., 2009). The other sources of model errors are due to the use of averaged temperatures, pressures, and water vapor, and due to the effect of dust and water ice. The uncertainty due to the use of the “averaged atmosphere” is estimated to be small and amounts to less than 1 ppb in the retrieved mixing ratio of H<sub>2</sub>O<sub>2</sub>. The uncertainty due to the aerosols is also estimated to be small and amounts to less than 5 ppb in the retrieved mixing ratio of H<sub>2</sub>O<sub>2</sub>. Finally, the error value in the retrieved H<sub>2</sub>O<sub>2</sub> mixing ratio  $\sigma$  was given by

$$\sigma = \sqrt{\sigma_n^2 + \sigma_{Int}^2 + \sigma_{avr}^2 + \sigma_{aer}^2},$$

where  $\sigma_n$  is error due to instrumental noise,  $\sigma_{Int}$  is that due to uncertainty in the line strength,  $\sigma_{avr}$  is that due to using the “averaged atmosphere,” and  $\sigma_{aer}$  is that due to ignoring the effect of aerosols. The derived mixing ratios of H<sub>2</sub>O<sub>2</sub> are  $16 \pm 19$  ppb at  $Ls = 0\text{--}120^\circ$  (i.e., 35 ppb upper limit),  $35 \pm 32$  ppb at  $Ls = 120\text{--}240^\circ$ , and  $41 \pm 28$  ppb at  $Ls = 240\text{--}360^\circ$ .

The derived mixing ratios of H<sub>2</sub>O<sub>2</sub> were close to the detection limits or upper limit because of the large uncertainty owing to instrumental noise. However, the values generally agreed with the ones reported by ground-based observations (see **Table 1**). Moreover, although the error values are large, our results imply that the mixing ratio of H<sub>2</sub>O<sub>2</sub> at  $Ls = 0\text{--}120^\circ$  is relatively low. The previous observations found similar aspects: IRTF/TEXES observations showed that the mean mixing ratio is  $10 \pm 5$  ppb at  $Ls = 80^\circ$  and the upper limit is 10 ppb at  $Ls$

1 = 112° (Encrenaz et al., 2012), and Herschel/HIFI observations revealed the upper limit is 2  
 2 ppb at  $L_s = 77^\circ$  (Hartogh et al., 2010).

3 From the retrieved mixing ratio of  $H_2O_2$ , we can restrict the oxidizing capacity in the  
 4 Martian atmosphere and constrain the  $CH_4$  sink on Mars. We calculated the  $CH_4$  lifetime near  
 5 the surface assuming photochemical steady state. First, the  $HO_2$  number density was estimated  
 6 from the observed  $H_2O_2$  density:

$$7 \quad [HO_2] = \sqrt{\frac{k_2[H_2O_2]}{k_1}},$$

8 where  $[HO_2]$  and  $[H_2O_2]$  represent the number densities, and  $k_1$  and  $k_2$  are the rate coefficients  
 9 of photochemical reactions shown in **Table 2**. Then, the OH number density was estimated  
 10 from the calculated  $HO_2$  density, the observed  $H_2O_2$  density, and the assumed densities of  $H_2O$ ,  
 11  $O_3$ , H,  $O(^1D)$ ,  $H_2$ , and CO listed in **Table 3**:

$$12 \quad [OH] = \frac{k_3[H_2O] + k_4[O_3][H] + k_5[HO_2][O] + 2k_6[H][HO_2] + 2k_2[H_2O_2] + k_7[HO_2] + 2k_8[O(^1D)][H_2O] + k_9[O(^1D)][H_2] + k_{10}[HO_2][O_3]}{k_{11}[O] + k_{12}[O_3] + k_{13}[HO_2] + k_{14}[H_2] + k_{15}[H_2O_2] + k_{16}[CO]},$$

13 where  $k_i$  is the rate coefficient of photochemical reactions shown in **Table 2**. In this equation,  
 14 the dominant terms are  $2k_6[HO_2][O]$ ,  $2k_2[H_2O_2]$ , and  $k_{16}[CO]$ . This means that the OH  
 15 amount depends on those of  $HO_2$  (which can be estimated from the  $H_2O_2$  amount),  $H_2O_2$ , and  
 16 CO. **Figure 4a** shows the OH number density estimated from the  $H_2O_2$  density. Finally, we  
 17 calculated the lifetime of  $CH_4$  as follows:

$$18 \quad \tau_{CH_4} = \frac{1}{k_{17}[OH] + k_{18}[O(^1D)] + k_{19}[O]},$$

19 where  $k_{17}$ ,  $k_{18}$ , and  $k_{19}$  are the rate coefficients of photochemical reactions shown in **Table 4**.

1 In this equation,  $k_{17}[OH]$  is the dominant term.

2 **Figure 4b** shows the calculated lifetime of  $CH_4$  associated with different  $H_2O_2$  number  
3 densities. Our observations showed that the averaged mixing ratio of  $H_2O_2$  is between 0 and  
4 70 ppb. The maximum value, i.e. 70 ppb, corresponds to  $1.1 \times 10^{10} \text{ cm}^{-3}$  near the surface. The  
5 calculated lifetime of  $CH_4$  with this  $H_2O_2$  abundance is about 50 years (solid line shown in  
6 **Figure 4b**), which is much longer than the observed lifetime on the order of weeks or months.  
7 The short observed lifetime of  $CH_4$  requires unrealistic large amounts of  $H_2O_2$  that correspond  
8 to  $10^{12}$ – $10^{13} \text{ cm}^{-3}$  (and this corresponds to more than  $10^6 \text{ cm}^{-3}$  of OH (see **Figure 4a**)). We  
9 also considered the CO abundance because OH density also depends on CO density. The  
10 observed minimum abundance of CO determined by the MEX/PFS was about 400 ppm  
11 (Sindoni et al., 2011), which corresponds to about  $5 \times 10^{13} \text{ cm}^{-3}$  near the surface. We adopted  
12 the minimum CO abundance and derived the lifetime of  $CH_4$  (dotted line in **Figure 4**).  
13 Although it became slightly shorter ( $\sim 30$  years), it is still much longer than weeks or months.  
14  $O(^1D)$  and O are the other minor candidates as  $CH_4$  sinks in the atmosphere (Wong et al.,  
15 2003). The dashed lines in **Figure 4b** represent the lifetime of  $CH_4$  with 10 times larger  
16 abundances of  $O(^1D)$  and O than the values predicted by Krasnopolsky (2006). In this case,  
17 the lifetime of  $CH_4$  becomes even shorter ( $\sim 10$  years), but it is still longer than the expected  
18 short lifetime of  $CH_4$ . We conclude that according to such a long-term averaged view, it is  
19 difficult to explain the observed short lifetime of  $CH_4$  by atmospheric oxidation processes.

Our result suggests the presence of strong CH<sub>4</sub> sinks not subject to atmospheric oxidation, e.g., an oxidant on the surface (Atreya et al., 2007) or wind eroded grains (Jansen et al., 2014).

## 5. Summary

We reported on the first attempt to search for H<sub>2</sub>O<sub>2</sub> in the Martian atmosphere from long-term measurements by a spacecraft-borne instrument using relatively high-dispersion spectroscopic observations with MEX/PFS. We used the strongest  $\nu_4$  band of H<sub>2</sub>O<sub>2</sub> around 362 cm<sup>-1</sup> and 379 cm<sup>-1</sup>. Although we performed sensitive data calibrations and averaged a large number of spectra in order to search for the tiny absorptions of H<sub>2</sub>O<sub>2</sub>, the derived mixing ratios of H<sub>2</sub>O<sub>2</sub> were close to the detection limits or upper limit. The derived mixing ratios were  $16 \pm 19$  ppb (i.e., 35 ppb upper limit) at  $L_s = 0\text{--}120^\circ$ ,  $35 \pm 32$  ppb at  $L_s = 120\text{--}240^\circ$ , and  $41 \pm 28$  ppb at  $L_s = 240\text{--}360^\circ$ . These results restrict the oxidizing capacity in the Martian atmosphere. Regarding the issue of CH<sub>4</sub> sinks on Mars, we conclude that atmospheric oxidation alone cannot explain the observed short lifetime of CH<sub>4</sub>.

## Acknowledgements

This work was supported by the Grant-in-Aid Fellows program of the Japan Society for the Promotion of Science (JSPS) (#233113). This work was also supported by a Grant-in-Aid for Scientific Research from the JSPS (#22340142) and by the Tohoku University Global Center

of Excellence (COE) program titled “Global Education and Research Center for Earth and Planetary Dynamics.” The PFS activities were funded by the Agenzia Spaziale Italiana (ASI) in the context of Italian participation in the European Space Agency (ESA) Mars Express mission. We would like to thank Dr. M. Smith and an anonymous reviewer for reviewing the manuscript and providing useful comments.

## References

- Atreya, S.K., Mahaffy, P.R., Wong, A., 2007. Methane and related trace species on Mars: Origin, loss, implications for life, and habitability. *Planet. Space Sci.* 55, 358–369.
- Clancy, R.T., Sandor, B.J., Moriarty-Schieven, G.H., 2004. A measurement of the 362 GHz absorption line of Mars atmospheric H<sub>2</sub>O<sub>2</sub>. *Icarus* 168, 116–121.
- Drayson, S.R., 1976. Rapid computation of the Voigt profile. *J. Quant. Spectrosc. Radiat. Transfer* 16, 611–614.
- Encrenaz, T., Greathouse, T.K., Bezard, B., Atreya, S.K., Wong, A.S., Richter, M.J., Lacy, J.H., 2002. A stringent upper limit of the H<sub>2</sub>O<sub>2</sub> abundance in the Martian atmosphere. *Astron. Astrophys.* 396, 1037–1044.
- Encrenaz, T., Bézard, B., Greathouse, T.K., Richter, M.J., Lacy, J.H., Atreya, S.K., Wong, A.S., Lebonnois, S., Lefèvre, F., Forget, F., 2004. Hydrogen peroxide on Mars: evidence for spatial and seasonal variations. *Icarus* 170, 424–429.

- 1 Encrenaz, T., Greathouse, T.K., Richter, M.J., Bézard, B., Fouchet, T., Lefèvre, F.,  
2 Montmessin, F., Forget, F., Lebonnois, S., Atreya, S.K., 2008. Simultaneous mapping of  
3 H<sub>2</sub>O and H<sub>2</sub>O<sub>2</sub> on Mars from infrared high-resolution imaging spectroscopy. *Icarus* 195,  
4 547–556.
- 5 Encrenaz, T., Greathouse, T.K., Lefèvre, F., Atreya, S.K., 2012. Hydrogen peroxide on Mars:  
6 Observations, interpretation and future plans. *Planet. Space Sci.* 68, 3–17.
- 7 Fonti, S., Marzo, G.A., 2010. Mapping of methane on Mars. *Astron. Astrophys.* 512, A51.
- 8 Forman, M.L., Steel, W.H., Vanasse, G.A., 1966. Correction of asymmetric interferograms  
9 obtained in Fourier spectroscopy. *J. Opt. Soc. Amer.* 56 (1), 59–61.
- 10 Formisano, V., Atreya, S.K., Encrenaz, T., Ignatiev, N., Giuranna, M., 2004. Detection of  
11 methane in the atmosphere of Mars. *Science* 306, 1758–1761.
- 12 Formisano, V., Angrilli, F., Arnold, G., Atreya, S., Bianchini, G., Biondi, D., Blanco, A.,  
13 Blecka, M.I., Coradini, A., Colangeli, L., Ekonomov, A., Esposito, F., Fonti, S., Giuranna,  
14 M., Grassi, D., Gnedykh, V., Grigoriev, A., Hansen, G., Hirsh, H., Khatuntsev, I., Kiselev,  
15 A., Ignatiev, N., Jurewicz, A., Lellouch, E., Lopez Moreno, J., Marten, A., Mattana, A.,  
16 Maturilli, A., Mencarelli, E., Michalska, M., Moroz, V., Moshkin, B., Nespoli, F.,  
17 Nikolsky, Y., Orfei, R., Orleanski, P., Orofino, V., Palomba, E., Patsaev, D., Piccioni, G.,  
18 Rataj, M., Rodrigo, R., Rodriguez, J., Rossi, M., Saggini, B., Titov, D., Zasova, L., 2005.  
19 The Planetary Fourier Spectrometer (PFS) onboard the European Mars Express mission.

- 1 Planet. Space Sci. 53 (10), 963–974.
- 2 Fouchet, T., Lellouch, E., Ignatiev, N.I., Forget, F., Titov, D.V., Tschimmel, M., Montmessin,  
3 F., Formisano, V., Giuranna, M., Maturilli, A., Encrenaz, T., 2007. Martian water vapor:  
4 Mars Express PFS/LW observations. *Icarus* 190, 32–49.
- 5 Geminale, A., Formisano, V., Sindoni, G., 2011. Mapping methane in Martian atmosphere  
6 with PFS-MEX data. *Planet. Space Sci.* 59, 137–148.
- 7 Giuranna, M., Formisano, V., Biondi, D., Ekonomov, A., Fonti, S., Grassi, D., Hirsch, H.,  
8 Khatuntsev, I., Ignatiev, N., Michalska, M., Mattana, A., Maturilli, A., Mencarelli, E.,  
9 Nespoli, F., Orfei, R., Orleanski, P., Piccioni, G., Rataj, M., Saggin, B., Zasova, L., 2005a.  
10 Calibration of the Planetary Fourier Spectrometer long wavelength channel. *Planet. Space*  
11 *Sci.* 53 (10), 993–1007.
- 12 Giuranna, M., Formisano, V., Biondi, D., Ekonomov, A., Fonti, S., Grassi, D., Hirsch, H.,  
13 Khatuntsev, I., Ignatiev, N., Michalska, M., Mattana, A., Maturilli, A., Moshkin, B.E.,  
14 Mencarelli, E., Nespoli, F., Orfei, R., Orleanski, P., Piccioni, G., Rataj, M., Saggin, B.,  
15 Zasova, L., 2005b. Calibration of the Planetary Fourier Spectrometer short wavelength  
16 channel. *Planet. Space Sci.* 53 (10), 975–991.
- 17 Hanel, R.A., Conrath, B.J., Jennings, D.E., Samuelson, R.E., 2003. Exploration of the Solar  
18 System by Infrared Remote Sensing, second edition, Cambridge University Press, U.K.

1 Hartogh, P., Jarchow, C., Lellouch, E., de Val-Borro, M., Rengel, M., Moreno, R., Medvedev,  
 2 A.S., Sagawa, H., Swinyard, B.M., Cavalié, T., Lis, D.C., Błęcka, M.I., Banaszkiewicz,  
 3 M., Bockelée-Morvan, D., Crovisier, J., Encrenaz, T., Küppers, M., Lara, L.-M.,  
 4 Szutowicz, S., Vandenbussche, B., Bensch, F., Bergin, E.A., Billebaud, F., Biver, N.,  
 5 Blake, G.A., Blommaert, J.A.D.L., Cernicharo, J., Decin, L., Encrenaz, P., Feuchtgruber,  
 6 H., Fulton, T., de Graauw, T., Jehin, E., Kidger, M., Lorente, R., Naylor, D.A.,  
 7 Portyankina, G., Sánchez-Portal, M., Schieder, R., Sidher, S., Thomas, N., Verdugo, E.,  
 8 Waelkens, C., Whyborn, N., Teyssier, D., Helmich, F., Roelfsema, P., Stutzki, J., LeDuc,  
 9 H.G., Stern, J.A., 2010. Herschel/HIFI observations of Mars: First detection of O<sub>2</sub> at  
 10 submillimetre wavelengths and upper limits on HCl and H<sub>2</sub>O<sub>2</sub>. *Astron. Astrophys.* 521,  
 11 L49.

12 Jansen, S.J.K., Skibsted, J., Jakobsen, H.J., Kate, I.L.T., Gunnlaugsson, H.P., Merrison, J.P.,  
 13 Finster, K., Bak, E., Iversen, J.J., Kondrup, J.C., Nørnberg, P., 2014. A sink of methane on  
 14 Mars? The answer is blowing in the wind. *Icarus* 236, 24–27.

15 Kong, T.Y., McElroy, M.B., 1977. Photochemistry of the Martian atmosphere. *Icarus* 32,  
 16 168–189.

17 Krasnopolsky, V.A., Bjoraker, G.L., Mumma, M.J., Jennings, D.E., 1997. High-resolution  
 18 spectroscopy of Mars at 3.7 and 8  $\mu$ m: A sensitive search of H<sub>2</sub>O<sub>2</sub>, H<sub>2</sub>CO, HCl, and CH<sub>4</sub>,  
 19 and detection of HDO. *J. Geophys. Res.* 102, 6525–6534.

- 1 Krasnopolsky, V.A., 2006. Photochemistry of the Martian atmosphere: Seasonal, latitudinal,  
2 and diurnal variations. *Icarus* 185, 153-170.
- 3 Lefèvre, F., Forget, F., 2009. Observed variations of methane on Mars unexplained by known  
4 atmospheric chemistry and physics. *Nature* 460 (7256), 720–723.
- 5 Mumma, M.J., Villanueva, G.L., Novak, R.E., Hewagama, T., Bonev, B.P., DiSanti, M.A.,  
6 Mandell, A.M., Smith, M.D., 2009. Strong release of methane on Mars in northern  
7 summer 2003. *Science* 323 (5917), 1041–1045.
- 8 Nair, H., Allen, M., Anbar, A.D., Yung, Y.L., Clancy, R.T., 1994. A photochemical model of  
9 the Martian atmosphere. *Icarus* 111, 124–150.
- 10 Rothman, L.S., Rinsland, C.P., Goldman, A., Massie, S.T., Edwards, D.P., Flaud, J.-M., Perrin,  
11 A., Camy-Peyret, C., Dana, V., Mandin, J.-Y., Schroeder, J., McCann, A., Gamache, R.R.,  
12 Wattson, R.B., Yoshino, K., Chance, K.V., Jucks, K.W., Brown, L.R., Nemtchinov, V.,  
13 Varanasi, P., 1998. The HITRAN molecular spectroscopic database and HAWKS  
14 (HITRAN Atmospheric Workstation): 1996 Edition. *J. Quant. Spectrosc. Radiat. Transfer*  
15 60, 665–710.
- 16 Rothman, L.S., Gordon, I.E., Barbe, A., Chris Benner, D., Bernath, P.F., Birk, M., Boudon, V.,  
17 Brown, L.R., Campargue, A., Champion, J.-P., Chance, K., Coudert, L.H., Dana, V., Devi,  
18 V.M., Fally, S., Flaud, J.-M., Gamache, R.R., Goldman, A., Jacquemart, D., Kleiner, I.,  
19 Lacome, N., Lafferty, W.J., Mandin, J.-Y., Massie, S.T., Mikhailenko, S.N., Miller, C.E.,

1 Moazzen-Ahmadi, N., Naumenko, O.V., Nikitin, A.V., Orphal, J., Perevalov, V.I., Perrin,  
2 A., Predoi-Cross, A., Rinsland, C.P., Rotger, M., Simeckova, M., Smith, M.A.H., Sung, K.,  
3 Tashkun, S.A., Tennyson, J., Toth, R.A., Vandaele, A.C., Vander Auwera, J., 2009. The  
4 HITRAN 2008 molecular spectroscopic database. *J. Quant. Spectrosc. Radiat. Transfer*  
5 110, 533–572.

6 Saggin, B., Scaccabarozzi, D., Tarabini, M., 2011. Instrumental phase-based method for  
7 Fourier transform spectrometer measurements processing. *Appl. Opt.* 50, 1717–1725.

8 Sander, S.P., Golden, D.M., Kurylo, M.J., Moortgat, G.K., Wine, P.H., Ravishankara, A.R.,  
9 Kolb, C.E., Molina, M.J., Finlayson-Pitts, B.J., Huie, R.E., Orkin, V.L., 2006. Chemical  
10 kinetics and photochemical data for use in atmospheric studies evaluation number 15. JPL  
11 Publ. 06-3, Jet Propul. Lab., Pasadena, Calif.

12 Sindoni, G., Formisano, V., Geminale, A., 2011. Observations of water vapour and carbon  
13 monoxide in the Martian atmosphere with the SWC of PFS/MEX. *Planet. Space Sci.* 59,  
14 149–162.

15 Smith, M.D., Bandfield, J.L., Christensen, P.R., 2000. Separation of atmospheric and surface  
16 spectral features in Mars Global Surveyor Thermal Emission Spectrometer (TES) spectra.  
17 *J. Geophys. Res.* 105 (E4), 9589–9607.

18 Smith, M.D., 2002. The annual cycle of water vapor on Mars as observed by the Thermal  
19 Emission Spectrometer. *J. Geophys. Res.* 107, E11.

1 Villanueva, G.L., Mumma, M.J., Bonev, B.P., Noval, R.E., Barber, R.J., Disanti, M.A., 2012.

2 Water in planetary and cometary atmospheres: H<sub>2</sub>O/HDO transmittance and fluorescence

3 models. J. Quant. Spectrosc. Radiat. Transfer 113, 202–220.

4 Wong, A.S., Atreya, S.K., Encrenaz, T., 2003. Chemical markers of possible hot spots on

5 Mars. J. Geophys. Res. 108 (E4), 5026.

6

7

8

9

10

11

12

13

14

15


16

17

18


19

# 1 Figures




**Figure 1.** (Top) Line strength of H<sub>2</sub>O and H<sub>2</sub>O<sub>2</sub> in the spectral range between 350 and 390 cm<sup>-1</sup> obtained from HITRAN08. (Middle) Synthetic spectra in the spectral range between 350 and 390 cm<sup>-1</sup> with the spectral resolution of the apodized PFS spectrum. The black curves show the spectra with various mixing ratios of H<sub>2</sub>O (dash: 100 ppm, dot: 200 ppm, solid: 300 ppm). The color curves show the spectra with various mixing ratios of H<sub>2</sub>O<sub>2</sub> (blue: 50 ppb,


light blue: 100 ppb, green: 500 ppb, red: 1 ppm). The arrows show the spectral features used for the search of H<sub>2</sub>O<sub>2</sub>. The gray box represents the spectral ranges used in this analysis as shown in **Figure 3**. (Bottom) The same as in the middle figure but with a different Y-axis scale. Note that an enhanced feature around 361 cm<sup>-1</sup> in the spectra of H<sub>2</sub>O is due to the side-lobes in the instrumental line shape of the PFS (Giuranna et al., 2005b).




**Figure 2.** The distributions of solar longitude  $L_s$  in the three datasets that were used. The histogram shows the numbers of measurements as a function of the solar longitude. The interval of the histogram is  $10^\circ$ .




(b) Ls: 120-240





**Figure 3.** The PFS averaged spectra over (a)  $L_s = 0-120^\circ$ , (b)  $L_s = 120-240^\circ$ , and (c)  $L_s = 240-360^\circ$  in the spectral range between 359 and 382  $\text{cm}^{-1}$ . (Top) Comparison between the synthetic spectra and the averaged PFS spectra. The red curves show the best-fit synthetic spectra. The other color curves represent synthetic spectra with various mixing ratios of  $\text{H}_2\text{O}_2$  (blue: 20 ppb, light blue: 40 ppb, green: 60 ppb, orange: 100 ppb). The black vertical bars show instrumental noise calculated by the noise equivalent radiance divided by the square root of the number of measurements. (Bottom) Residual spectra, where special synthetic spectra with all contributions except  $\text{H}_2\text{O}_2$  have been subtracted from the measured and synthetic (including  $\text{H}_2\text{O}_2$  contribution) spectra. The retrieved  $\text{H}_2\text{O}_2$  values are  $16 \pm 19$  ppb at  $L_s =$

0–120° (i.e., 35 ppb upper limit),  $35 \pm 32$  ppb at  $L_s = 120\text{--}240^\circ$ , and  $41 \pm 28$  ppb at  $L_s =$   
240–360°; the derived H<sub>2</sub>O column densities in the three averages were  $2.43 \times 10^{19} \text{ cm}^{-2}$ ,  $3.52$   
 $\times 10^{19} \text{ cm}^{-2}$ , and  $2.50 \times 10^{19} \text{ cm}^{-2}$ , respectively. The red vertical bars show instrumental noise  
estimated from the standard deviation of the residuals between the PFS data and the best-fit  
synthetic spectra.



**Figure 4.** (a) OH number density estimated from the H<sub>2</sub>O<sub>2</sub> number density. (b) Estimated photochemical lifetime of CH<sub>4</sub> with the abundances of H<sub>2</sub>O<sub>2</sub> near the surface. Our observations showed that the maximum mixing ratio of H<sub>2</sub>O<sub>2</sub> is  $1.1 \times 10^{10} \text{ cm}^{-3}$  near the surface. The solid line represents the lifetime with the normal input parameters listed in **Table 2**. The dotted line represents the lifetime with rich CO conditions ( $5 \times 10^8 \text{ cm}^{-3}$ ). The dashed lines represent the lifetime with rich O(<sup>1</sup>D) and O conditions, and these were 10 times larger than the values predicted in Krasnopolsky et al. (2006).