



Publication Year	2017
Acceptance in OA	2020-08-27T09:07:58Z
Title	Characterization of the white ovals on Jupiter's southern hemisphere using the first data by the Juno/JIRAM instrument
Authors	Sindoni, G., GRASSI, Davide, ADRIANI, Alberto, MURA, Alessandro, Moriconi, M. L., Dinelli, B. M., FILACCHIONE, GIANRICO, TOSI, Federico, PICCIONI, GIUSEPPE, MIGLIORINI, Alessandra, ALTIERI, FRANCESCA, Fabiano, F., TURRINI, Diego, NOSCHESE, RAFFAELLA, CICCETTI, ANDREA, STEFANI, STEFANIA, Bolton, S. J., Connerney, J. E. P., Atreya, S. K., Bagenal, F., Hansen, C., Ingersoll, A., Janssen, M., Levin, S. M., Lunine, J. I., Orton, G., Olivieri, A., Amoroso, M.
Publisher's version (DOI)	10.1002/2017GL072940
Handle	http://hdl.handle.net/20.500.12386/26859
Journal	GEOPHYSICAL RESEARCH LETTERS
Volume	44

Characterization of the white ovals on Jupiter's southern hemisphere using the first data by the Juno/JIRAM instrument

G. Sindoni^{1*}, D. Grassi¹, A. Adriani¹, A. Mura¹, M. L. Moriconi^{2,1}, B. M. Dinelli², G. Filacchione¹, F. Tosi¹, G. Piccioni¹, A. Migliorini¹, F. Altieri¹, F. Fabiano^{13,2}, D. Turrini^{1,14}, R. Noschese¹, A. Cicchetti¹, S. Stefani¹, S. J. Bolton³, J. E. P. Connerney⁴, S. K. Atreya⁵, F. Bagenal⁶, C. Hansen⁷, A. Ingersoll⁸, M. Jansen⁹, S. M. Levin⁹, J. I. Lunine¹¹, G. Orton⁹, A. Olivieri¹² and M. Amoroso¹²

¹Institute for Space Astrophysics and Planetology (IAPS-INAF), Rome, Italy

²Institute of Atmospheric Sciences and Climate (ISAC-CNR), Bologna, Italy

³Southwest Research Institute, San Antonio, Texas, USA

⁴NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

⁵University of Michigan, Ann Arbor, Michigan, USA

⁶University of Colorado, Boulder, Colorado, USA

⁷Planetary Science Institute, Tucson, Arizona, USA

⁸California Institute of Technology, Pasadena, California, USA

⁹Jet Propulsion Laboratory, Pasadena, California, USA

¹⁰University of Iowa, Iowa City, Iowa, USA

¹¹Cornell University, Ithaca, New York, USA

¹²Agenzia Spaziale Italiana, Roma, Italy

¹³Dipartimento di Fisica e Astronomia, Università di Bologna, Italy

¹⁴Departamento de Física, Universidad de Atacama, Copayapu 485, Copiapó, Chile

Corresponding author: Giuseppe Sindoni (giuseppe.sindoni@iaps.inaf.it)

*Address: Via del Fosso del Cavaliere 100, 00133 Roma, ITALY. Phone: 00390645488511

Key Points:

- JIRAM data from first Juno orbit highlighted the presence of several ovals in Jupiter's southern hemisphere.
- Column densities and altitudes for an NH₃ cloud and a N₂H₄ haze have been retrieved for three ovals in the region between 60-120°W and 30-45°S.
- Evidences of cyclonic and anticyclonic structures inside the ovals have been found.

Abstract

During the first perijove passage of the Juno mission, the Jovian InfraRed Auroral Mapper (JIRAM) observed a line of closely spaced oval features in Jupiter's southern hemisphere, between 30°S and 45°S. In this work, we focused on the longitudinal region covering the three ovals having higher contrast at 5 μm , i.e. between 120°W and 60°W in System III coordinates. We used the JIRAM's full spectral capability in the range 2.4-3 μm together with a Bayesian data inversion approach to retrieve maps of column densities and altitudes for an NH_3 cloud and a N_2H_4 haze. The deep (under the saturation level) volume mixing ratio and the relative humidity for gaseous ammonia were also retrieved. Our results suggest different vortex activity for the three ovals. Updraft and downdraft together with considerations about the ammonia condensation could explain our maps providing evidences of cyclonic and anticyclonic structures.

1 Introduction

The white ovals, together with the Great Red Spot (GRS), are the most prominent features in the Jupiter's atmosphere. They were first observed by ground-based measurements [Peek, 1958], then in more detail during the flybys of Jupiter carried out by the NASA Voyager probes in 1979 and during the extensive in-system tour performed by the NASA Galileo spacecraft in 1995-2003. These ovals are in a stable configuration known as a "Karman vortex street", where anticyclones are staggered with cyclones [Youssef and Marcus, 2003]. These vortices appear very bright in the visible range and for this reason they are usually called "white". Their visible brightness is due to the scattering of the sunlight by upper clouds and hazes condensed there. On the other hand, they appear very dark in the spectral range where the thermal radiation becomes dominant since thick clouds block photons coming from the deeper and hotter atmospheric layers. Color changes in the visible range, from white to red, have been observed in ovals [Go *et al.*, 2006]. The reddish color is usually attributed to red "chromophores" [West *et al.*, 1986], which are products of a series of complex chemical reactions, such as the UV photolization of ammonia with acetylene [Carlson *et al.*, 2016]. These chromophores can act as coating material for the ammonia particles.

The current understanding of the clouds structure in the Jupiter's atmosphere is mainly related to theoretical models based on the thermochemical equilibrium theory [Weidenschilling and Lewis, 1973; Atreya *et al.*, 1999]. They expect an ammonia ice layer at about 700 mbar, an ammonium hydrosulfide cloud layer at about 2 bars and a water cloud layer with a base near 6 bars. Moreover, observations from the ground and from space in the visible, near and thermal infrared provided important information about the aerosol composition of the planet [West *et al.*, 1986; West *et al.*, 2004]. In particular, the equatorial region, the GRS and other small-scale cloud formations (as the ovals) were investigated using the Galileo/SSI (Solid-State Imaging) instrument by Banfield *et al.* [1998]. They modeled the atmosphere with a stratospheric haze layer, a denser upper tropospheric haze and a cloud like layer at pressures between 600 and 950 mbar. Their results suggested an increase of the altitude and of the aerosol optical depth inside the GRS and the white ovals with a two-cell structure with subsidence in the upper troposphere and upwelling in the lower troposphere. The three layered structure for the cloudy atmosphere of Jupiter was also confirmed by Irwin *et al.* [1998]. Their analysis of Galileo/NIMS (Near Infrared Mapping Spectrometer) spectra suggested the presence of a tropospheric haze at 200 mbar, an ammonia cloud at about 700 mbar, and an ammonium hydrosulfide cloud at pressures between

1.4 and 5 bars. Using the Galileo/NIMS and SSI data, *Dyudina et al.* [2001] investigated the cloud structure over the GRS and a white oval. They again suggested an increase in the optical thickness and in the elevation of the cloud inside both the atmospheric structures. Radiative transfer modeling of the infrared spectra in the 1-2 μm range acquired using ground-based measurements with Keck telescope [*de Pater et al.*, 2010] showed an overall similar vertical structure of all the studied features (GRS, Oval BA and two white ovals at 41°S). An atmospheric model based on three layer hazes (2-200 mbar, 200-650 mbar and 650-700 mbar) composed of ammonia ice particles revealed that the particle densities of the tropospheric-stratospheric hazes in the two white ovals are about 5-8 times smaller than in the GRS and Oval BA. The center of the vortexes shows a cloud at a pressure below of 600 mbar [*Dyudina et al.*, 2001], which is approximately the ammonia condensation level considering the expected mixing ratio in Jupiter's atmosphere. An optically thick cloud near the NH_3 condensation level should contain large NH_3 -ice particles, which drop down to the equilibrium condensation level where they sublime [*West et al.*, 1986]. Thus, gaseous NH_3 depletion is associated with cloud formation. Both large-scale updrafts and turbulence can bring NH_3 back into the cloud raising in altitude its upper boundary.

Simultaneous observations in the solar (below $\sim 2 \mu\text{m}$) and thermal ($\sim 5 \mu\text{m}$) spectral ranges highlighted the presence of hot rings (225-250 K) surrounding the periphery of ovals [*de Pater et al.*, 2010]. Since these thermal bright regions should be cloud-free down to about 5 bars, they suggested that anticyclones could extend vertically from the water cloud to the tropopause. Moreover, the observation of an anticorrelation in the opacities of the high tropospheric haze and of the low $5\text{-}\mu\text{m}$ absorbing cloud could be explained by the descent of air at high altitude and the air ascent at low altitude, and vice versa [*Dyudina et al.*, 2001].

The cloud structure of the Jupiter's atmosphere, and in particular the nature of vortex features, as the GRS and the white ovals, is still puzzling. Indeed, only a very limited number of ovals has been investigated in details using the visible and infrared spectroscopy. Space-borne new instruments will cast light on this issue thanks to their improved spatial resolution and the possibility to have a relatively long-time monitoring of the atmosphere.

Here we report the preliminary results about the cloud and haze structures inside and outside three ovals in the southern hemisphere of Jupiter obtained by the analysis of the Jovian InfraRed Auroral Mapper (JIRAM) observations acquired during the first perijove passage of the NASA spacecraft Juno after orbit insertion (PJ1). Spectral observations with a typical spatial resolution of around 250 km at the 1 bar level, the highest achieved so far from spacecraft, allow for an accurate characterization of these peculiar atmospheric features.

2 Dataset

The JIRAM instrument, aboard the NASA/Juno spacecraft, is composed of an infrared imager (IMG) and a spectrometer (SPE) [*Adriani et al.*, 2014], sharing a common optical head. The imager is further split in two spectral channels: L-filter, centered at $3.45 \mu\text{m}$ with a 290 nm bandwidth, and M-filter, centered at $4.78 \mu\text{m}$ with a 480 nm bandwidth. Each channel has a Field of View (FOV) of $1.75^\circ \times 5.94^\circ$ (128 x 432 pixels corresponding to the along and across track directions) and an Instantaneous Field of View (IFOV) of about $250 \times 250 \mu\text{rad}$. The spectrometer, based on grating diffraction of a pixel size slit, covers the spectral interval 2.0-5.0 μm with a spectral resolution around 12 nm and has a FOV of 3.52° (across track) sampled by 256 pixels with a square IFOV of $250 \times 250 \mu\text{rad}$. The radiometric calibration is described in

Adriani et al. [2014], whereas in-flight performances were first evaluated during the Earth-Moon flyby occurred in October 2013 [*Adriani et al.*, 2016]. In this work we used the M-filter of the IMG for the context and the SPE for the characterization analysis.

Juno orbit insertion successfully occurred on July 4, 2016, and during its first perijove passage (August 27-28, 2016) JIRAM acquired measurements covering almost the entire planet. In particular, JIRAM highlighted the presence of the white ovals belt in the southern hemisphere, between 30°S and 45°S. In this work, we focused on the longitudinal region covering the three ovals having higher contrast at 5 μm , i.e. between 60°W and 120°W (Fig. 1, in System III coordinates). Hereafter we refer to the region in this planetocentric rectangle as our Region of Interest (ROI). The identified ovals are all centered at a latitude of about 37°S and at longitudes of about 115°W, 88°W, and 67°W for Oval#1, Oval#2 and Oval#3, respectively. In the M band they show approximately the same diameter of less than 5° on both axes. At that latitude, this angular diameter results in a size of about 5000×6000 km.

In order to achieve the best spatial coverage, we divided the ROI in 3600 square bins of 0.5° in both latitude and longitude. The JIRAM spectrometer dayside acquisitions in PJ1 covered about 85% of these bins with a variable number of spectra spanning from 1 to 19, with an average of about 3. Since the variability in the observational (incidence, emission and phase) angles of spectra falling into each bin is mostly of the order of few degrees or less, measurements were then averaged to build a dataset composed of 3068 mean spectra (Fig. 1b-c).

We estimated the instrumental noise affecting the measurements by the standard deviation of a selected set of 522 spectra acquired when the JIRAM instrument pointed to deep space in the same science session. Therefore, our dataset has a mean signal-to-noise ratio (SNR) of 632 with a maximum of 1703, a minimum of 6 and a standard deviation of 269 at 2.75 μm .

3 The Analysis Procedure

3.1 Method

Clouds and hazes strongly affect the JIRAM spectra. We limited our analysis to the spectral range between 2.4 and 3 μm since it is sensitive to reflected sunlight and it can be used to sound the higher troposphere and the stratosphere. Moreover, it appears to be free from any calibration or instrumental issue. This range covers the spectral features of both gasses and particles of Jupiter's atmosphere and for this reason their interaction with radiation should be appropriately treated by using multiple scattering radiative transfer techniques.

The retrieval of the atmospheric composition exploits a data inversion technique with the Bayesian approach [*Rodgers*, 2000]. It uses the Gauss-Newton iterative procedure to minimize the χ^2 function. The solution at the n^{th} iterative step is:

$$x_{n+1} = x_n + (S_a^{-1} + K_n^T S_e^{-1} K_n)^{-1} (K_n^T S_e^{-1} [y - F(x_n)] - S_a^{-1} [x_n - x_a]) \quad \text{Eq. 1}$$

where x is the state vector of the free parameters, y is the vector with the measured radiances, F is the simulated spectrum, K is the Jacobian matrix, x_a is the a-priori knowledge of the state vector and S_a and S_e are the variance-covariance matrices for x_a and y , respectively. The synthetic spectrum F takes into account the multiple scatterings, both by molecules and particles, and it is computed using a line-by-line code in plane-parallel approximation [*Ignatiev et al.*, 2005]. This approach is suitable for simulating nadir or slant observations (incidence and emission angles < 70°), as is our case. Finally, the convergence criterion takes into account the χ^2

minimization. Convergence is reached when the minimum χ^2 does not decrease within 10 subsequent iterations.

3.2 Atmospheric Model

The *a priori* knowledge of Jupiter's atmosphere is based on the model suggested by *Grassi et al.* [2010] and takes into account the gaseous opacities by CH_4 , H_2O , NH_3 , and PH_3 and the collision-induced absorptions (CIA) by $\text{H}_2\text{--H}_2$, $\text{H}_2\text{--He}$ and $\text{H}_2\text{--CH}_4$. We used a temperature-pressure profile for Jupiter's atmosphere as derived from the Galileo Probe data [*Seiff et al.*, 1998], which can be considered representative also for the conditions in ovals [*de Pater et al.*, 2010]. Two tropospheric clouds and a stratospheric haze describe the atmospheric particulate component. The properties of the deepest NH_4SH cloud we considered are the ones suggested by *Giles et al.* [2015]. They are assumed to have single scattering albedo $\omega=0.9$ and asymmetry factor $g=0.7$ for the Henyey-Greenstein phase function in the entire spectral range of our interest. Moreover, the NH_4SH cloud is considered to extend between 1.3 and 0.73 bars with a constant volume density of 2.16 particles/ cm^3 . Methane photochemistry models produced a haze of complex hydrocarbons in the stratosphere of Jupiter that was postulated to coat the ammonia clouds below [*Atreya et al.*, 2005], thus obscuring them except for fresh clouds that alone could be spectrally identified [*Baines et al.*, 2002]. Laboratory experiments [*Kalogerakis et al.*, 2008] suggested tholin as a possible coating substance for NH_3 ice particles. In our work, we computed the single scattering parameters for the tholin-coated NH_3 cloud using the BART code [*Quirantes and Delgado*, 1997], which is based on the Aden-Kerker theory, starting from the refractive index by *Howett et al.* [2007] and *Khare et al.* [1984] for ammonia ice and tholin, respectively. We tested different tholin refractive indexes covering the spectral region of interest as suggested in the work by *Brassé et al.* [2015]. The ones from *Khare et al.* [1984] provided the best-fit results on a sample of our dataset. For this cloud we assumed a Gaussian vertical distribution between 1 bar and 200 mbar with the center at 500 mbar and a volume density of about 19 particles/ cm^3 . Finally, we considered a hydrazine (N_2H_4) stratospheric haze as suggested by *Moreno* [1996] with a Gaussian vertical distribution between 40 and 20 mbar centered at 30 mbar, and a particle volume density of about 4 particles/ cm^3 . The optical properties for the haze were computed using the refractive index by *Clapp and Miller* [1996] with a code based on the Mie theory. Finally, the solar spectrum was modeled following *Drummond and Thekaekara* [1973] and rescaled according to the actual heliocentric distance at the time of the acquisition.

The analysis procedure retrieves for each bin, starting from the reported *apriori* values (apv), the following parameters:

- the column number density for the NH_3 -tholin cloud ($\text{NH}_3\text{-Tho_cnd}$): $\text{apv} = 2.8 \times 10^7$ #/ cm^2 ;
- the peak altitude of the NH_3 -tholin cloud ($\text{NH}_3\text{-Tho_alt}$): $\text{apv} = 17.45$ km;
- the column number density for the N_2H_4 haze ($\text{N}_2\text{H}_4\text_cnd}$): $\text{apv} = 2.0 \times 10^6$ #/ cm^2 ;
- the peak altitude of the N_2H_4 haze ($\text{N}_2\text{H}_4\text_alt}$): $\text{apv} = 70.10$ km;
- the deep mixing ratio of gaseous ammonia ($\text{NH}_3\text_vmr_deep}$): $\text{apv} = 6.77 \times 10^{-4}$;
- the relative “humidity” of gaseous ammonia ($\text{NH}_3\text_rh}$): $\text{apv} = 2.02$ %.

We fixed the radii ratio to 0.5 and the total radius to 0.8 μm for the tholin-coated NH_3 particles and the radius of the N_2H_4 haze to 0.3 μm as the best result of several tests obtained from a sample of our dataset. Among the gaseous atmospheric components, only ammonia showed variability in the analyzed spectral range. The other atmospheric quantities were fixed to their *a priori* values. A typical example of spectral best fit is displayed in Fig. 2.

The quality of the retrieval is assessed using the χ^2 . Results showing a reduced χ^2 greater than 50 are discarded. Then, for each parameter we further filtered the retrieved values, discarding those out of the 99.9% confidence level for the statistical distribution of the parameter population on the entire retrieved dataset. This statistical filter in fact rules out retrieved values with a high probability to be non-physical, even though providing a good fit (low χ^2). Following this approach, we finally discarded about 16.5% of our results and retained retrievals obtained at around 2560 out of 3068 bins.

3.3 Uncertainties

We estimated the uncertainties for the retrieved parameters by applying the retrieval procedure to a simulated synthetic dataset. The statistics on our retrieved dataset naturally provide the range of variability of the considered parameters. We created a set of 100 synthetic spectra for each of the six parameters with randomly variable values in that range, obtaining a total of 600 synthetic spectra. After the addition of the instrumental noise we analyzed them. The mean differences between the known random parameters used for the spectra generation and the retrieved parameters provide the uncertainties on each of them. Thus, we obtained 24.76%, 29.77%, 69.59%, 63.36%, 49.23% and 32.05% for the relative maxima uncertainties on NH_3 -Tho_cnd, NH_3 -Tho_alt, N_2H_4 _cnd, N_2H_4 _alt, NH_3 _vmr_deep, NH_3 _rh, respectively, over our dataset. The two parameters related to the upper haze are those with higher uncertainty, whereas the retrieval of other parameters has errors of about 30%.

4 Results and Discussion

In Fig. 3 we report the main results of the analysis of JIRAM's spectroscopic data in the ROI. Maps of retrieved parameters are smoothed over two neighboring points to improve legibility.

The entire latitudinal belt between 41° and 37°S is characterized for the most part by very low NH_3 -tholin cloud densities at altitudes relatively higher than the surrounding atmosphere. The three ovals are generally identifiable in all the retrieved maps. However, Oval#1 and Oval#3 show more prominent features. They are mainly characterized by higher column densities for the NH_3 -tholin cloud. The first cloud (Fig. 3, upper panels) has values around 8.0×10^8 particles/ cm^2 in Oval#1 and Oval#3, whereas Oval#2 shows lower values by a factor of two. In Ovals#2 and Oval#3, the thickness of their NH_3 -tholin cloud increases from the border to the center of the vortex. On the contrary, the cloud thickness at the center of Oval#1 is 1.5 times lower than in the surrounding ring, which delimits the boundary of the vortex. In all the ovals we find an increasing altitude (measured from the pressure level of 1 bar) of the cloud from the boundary rings, with a peak at about -1 km (1.04 bar), toward the vortex center, where the peak is at about 10 km (660 mbar). This result is in agreement with previous observations by Galileo/NIMS and Galileo/SSI data [Dyudina *et al.*, 2001], where the altitude of the NH_3 ice cloud in the center of a white oval is at pressures less than 600 mbar. These instruments also observed an increase of the cloud thickness from the boundary ring toward the center of the vortex. Our analysis shows a

similar behavior for Oval#3 and partially for Oval#2, whereas Oval#1 exhibits the opposite configuration. A possible explanation for these differences can be found in the updrafts and downdrafts in the centers of vortices that carry gaseous ammonia and condensed particles [West *et al.*, 1986]. In Oval#2 and Oval#3 an updraft transports the NH_3 gas to higher altitudes (above 10 km, 660 mbar), where it condenses into ice particles. Then they subside in the ring boundary and sublimate at lower altitude. This is consistent with the behavior of an anticyclonic structure in a giant planet [Irwin, 2009; de Pater *et al.*, 2011]. On the contrary, in Oval#1, NH_3 condenses at lower altitudes (around 940 mbar) in the ring and then ice particles move with increasing altitudes (up to the level around 660 mbar) toward the center, where they subside. The central downdrafts and turbulence push the particles down to atmospheric layers where the temperature is above the condensation level, so they can sublimate and decrease the thickness of the cloud. This behavior is instead typical of cyclones.

In Fig. 3 (middle panel) we observe a different behavior for the stratospheric N_2H_4 haze. Only Oval#3 shows a notably high column density of about 6.0×10^7 particles/cm². The latitudinal belt 40-35°S, which contains the three ovals, shows a diffuse and relatively thick haze (about 4.0×10^7 particles/cm²). This haze stripe hides the presence of Oval#1 and Oval#2. These ovals cannot be identified also in the haze altitude map. Oval#3, the only one identifiable, corresponds to the minimum value of the entire map. It is characterized by increasing densities and decreasing peak altitude of the N_2H_4 haze, suggesting that hydrazine is condensing at lower altitudes (around 69 km, 29 mbar). Another atmospheric structure on the east side of Oval#3 dominates the N_2H_4 maps. It has both high densities (5.5×10^7 particles/cm²) and altitudes (71.9 km, 25.27 mbar). A possible explanation could involve an updraft centered around 63°W-38°S that, carrying tropospheric molecules to higher altitude, triggers the chemical production and condensation of hydrazine. However, this peculiar atmospheric feature will be investigated in detail in future works.

The map for the deep mixing ratio of gaseous ammonia is generally anticorrelated with the density map for the NH_3 -tholin map. This suggests that, as we can expect, ammonia condenses in the deep atmosphere and then atmospheric dynamics transport the NH_3 ice particles to the upper layers. While moving, they can act as condensation nuclei for hydrocarbons or other chemical products.

A peculiarity shows up in the map for the ammonia relative humidity. Oval#2 and Oval#3 show high values up to about 10.0% and 19.5%, respectively, increasing from the boundary ring toward the vortex center, whereas Oval#1 has lower values (~4-7%) with a flatter spatial distribution. This result can be explained by invoking condensation at different altitudes. Indeed, higher values for relative humidity imply lower atmospheric pressures (higher altitudes) to condense particles, and vice versa. Our finding suggests a correlation between the condensation altitude for ammonia ice particles and the relative humidity map, highlighting again the high differentiation in the dynamics of ovals in the same latitudinal belt.

An interesting atmospheric structure appears in all of our maps. It is located at latitudes slightly higher with respect to the ovals and it has an irregular shape. It extends from 92° to 113°W longitude and from 33° to 37°S latitude (see panels in Fig. 3). It is characterized by a very high column density for the NH_3 -tholin cloud growing from the boundaries to the center, where it reaches values around 1.2×10^9 particles/cm². Consistently with the gaseous ammonia condensation at low altitude, this structure shows a very low NH_3 deep mixing ratio ($< 10^{-6}$ below the saturation level). Similarly, it results to be strongly depleted in N_2H_4 haze particles.

Here we can sound altitudes down to about 1.3 bars, where we find the peak of the first significant cloud.

Several other atmospheric features can be recognized in our maps, whose detailed analysis is beyond the scope of this paper and will be the subject of forthcoming papers.

5 Conclusions

The JIRAM measurements acquired during the first orbit of Juno around Jupiter highlighted the presence of several ovals in the southern hemisphere. We selected the three ovals (Oval#1, Oval#2 and Oval#3) showing the higher contrast in the M band (around 5 μm) on which we performed a preliminary analysis using the JIRAM's full spectral capability in the range 2.4-3 μm . This spectral range is sensitive to changes in high tropospheric clouds and in stratospheric hazes, as well as to gaseous ammonia. The application of a data inversion algorithm based on a Bayesian approach provided information about the atmospheric structure inside and outside the oval vortices. As a result of our retrieval we produced maps of the region between 60-120°W and 30-45°S for the column densities and the altitudes of a tholin-coated NH_3 cloud and of a N_2H_4 haze. Moreover, we mapped also the gaseous ammonia deep mixing ratio (below the saturation level) and its relative "humidity".

Our results show a different behavior for the three ovals we studied. We observe evidences of a strong anticyclone producing Oval#3, where both the NH_3 -tholin cloud thickness and peak altitude increase toward the center. This is consistent with the central updraft that carries gaseous ammonia above the saturation level. The same structure is observed for the Oval#2 but with much less intense features. On the contrary, Oval#1, having higher cloud thickness in the boundary ring and higher cloud peak altitude in the vortex center, implies the presence of a central downdraft and thus shows evidence of a typical cyclonic structure in a giant planet. JIRAM data from further orbits will be used to monitor these behaviors.

Very high densities in the hydrazine haze with respect to other vortices and the surrounding environment also suggest the stronger activity of Oval#3. Moreover, the relative humidity map for the gaseous NH_3 reveals higher values in Oval#3, which implies condensation for ice particles at higher altitudes. Condensation is also observed in Oval#2 and Oval#1, though with weaker effects and at lower altitudes.

The results reported in this paper are the first obtained from JIRAM observations and therefore should be regarded as preliminary. Simultaneous analysis of solar and thermal dominated spectral range, as well as the extension to other interesting atmospheric features, could provide important insights into the clouds structure and the gaseous content in Jupiter's atmosphere.

Acknowledgments

The JIRAM project is founded by the Italian Space Agency (ASI). In particular this work has been developed under the agreement n. 2016-23-H.0. JIL and SKA acknowledge support from NASA through the Juno Project. GSO acknowledges support from NASA through funds that were distributed to the Jet Propulsion Laboratory, California Institute of Technology.

The JIRAM-Juno data are not currently publicly available. They will be made publicly available through the NASA Planetary Data System (<https://pds.nasa.gov>) at the end of proprietary period (currently under definition).

The radiative transfer code is based on the ARS software, developed by Nikolay Ignatiev, Space Research Institute of the Russian Academy of Sciences.

We thank the two anonymous reviewers, whose comments facilitated the improvement of this paper.

Emiliano D'Aversa and Fabrizio Oliva (IAPS-INAF) are acknowledged for the support in software development and extensive discussion.

The JIRAM instrument has been developed by Leonardo at the Officine Galileo - Campi Bisenzio site.

The computational resources used in this research have been supplied by INAF-IAPS through the projects "HPP - High Performance Planetology" and "DataWell".

References

- Adriani A., G. Filacchione, T. Di Iorio, D. Turrini, R. Noschese, A. Cicchetti, D. Grassi, A. Mura, G. Sindoni, M. Zambelli, G. Piccioni, M.T. Capria, F. Tosi, R. Orosei, B.M. Dinelli, M.L. Moriconi, E. Roncon, J.I. Lunine, H.N. Becker, A. Bini, A. Barbis, L. Calamai, C. Pasqui, S. Nencioni, M. Rossi, M. Lastri, R. Formaro, A. Olivieri (2014), JIRAM, the Jovian Infrared Auroral Mapper. *Space Sci. Rev.*, DOI 10.1007/s11214-014-0094-y.
- Adriani, A., Moriconi, M. L., Mura, A., Tosi, F., Sindoni, G., Noschese, R., Cicchetti, A., Filacchione, G. (2016), Juno's Earth flyby: the Jovian infrared Auroral Mapper preliminary results. *Astrophysics and Space Science*, Volume 361, Issue 8, article id.272. DOI: 10.1007/s10509-016-2842-9.
- Atreya S.K., Wong M.H., Owen T.C., Mahaffy P.R., Niemann H.B., de Pater I., Drossart P., Encrenaz Th. (1999), Comparison of the atmospheres of Jupiter and Saturn: deep atmospheric composition, cloud structure, vertical mixing, and origin. *Planet. Space Sci.*, 47, 1243-62. doi:10.1016/S0032-0633(99)00047-1.
- Atreya S. K., Wong A.S., Baines K.H., Wong M.H., Owen T.C. (2005). Jupiter's ammonia clouds—localized or ubiquitous? *Planetary and Space Science* 53, 498–507.
- Baines K.H., Carlson R.W., Kamp L.W. (2002), Fresh ammonia ice clouds in Jupiter. I. Spectroscopic identification, spatial distribution, and dynamical implications. *Icarus* 159:74-94.
- Banfield, D., Gierasch, P.J., Bell, M., Ustinov, E., Ingersoll, A.P., Vasavada, A.R., West, R.A., Belton, M.J.S., (1998). Jupiter's cloud structure from Galileo imaging data. *Icarus* 135, 230–250.
- Brassé C., Muñoz O., Coll P., Raulin F., (2015). Optical constants of Titan aerosols and their tholins analogs: Experimental results and modeling/observational data. *Planetary and Space Science* 109-110 : 159–174.
- Carlson, R. W.; Baines, K. H.; Anderson, M. S.; Filacchione, G.; Simon, A. A., (2016), Chromophores from photolyzed ammonia reacting with acetylene: Application to Jupiter's Great Red Spot. *Icarus*, Volume 274, p. 106-115. DOI: 10.1016/j.icarus.2016.03.008.

- 374 Clapp M.L. and Miller R.E. (1996), Complex Refractive Indices of Crystalline Hydrazine from
375 Aerosol Extinction Spectra. *Icarus* 123:396–403.
- 376 de Pater I., Wong M. H., Marcus P., Luszcz-Cook S., Ádámkovics M., Conrad A., Asay-Davis
377 X.; Go C. (2010). Persistent rings in and around Jupiter's anticyclones - Observations and
378 theory. *Icarus* 210, 742–762.
- 379 de Pater I., Wong M. H., de Kleer K., Hammel H. B., Ádámkovics M., Conrad A., (2011). Keck
380 adaptive optics images of Jupiter's north polar cap and Northern Red Oval. *Icarus* 213,
381 559–563.
- 382 Drummond A.J. and Thekaekara M. P. (1973). The Extraterrestrial Solar Spectrum. Institute of
383 Environmental Science, Mount Prospect, Illinois.
- 384 Dyudina U.A., Ingersoll A.P., Danielson G.E., Baines K.H., Carlson R.W. and The Galileo
385 NIMS and SSI Teams (2001). Interpretation of NIMS and SSI Images on the Jovian
386 Cloud Structure. *Icarus* 150, 219–233.
- 387 Giles R.S., Fletcher L. N. and Irwin P. G. J. (2015), Cloud structure and composition of Jupiter's
388 troposphere from 5-m Cassini VIMS spectroscopy, *Icarus*, 257, 457-470,
389 doi:10.1016/j.icarus.2015.05.030.
- 390 Grassi, D., Adriani, A., Moriconi, M. L., Ignatiev, N. I., D'Aversa, E., Colosimo, F., Negrão, A.,
391 Brower, L., Dinelli, B. M., Coradini, A., Piccioni, G. (2010) Jupiter's hot spots:
392 Quantitative assessment of the retrieval capabilities of future IR spectro-imagers,
393 *Planetary and Space Science*, 58, 1265-1278, doi:10.1016/j.pss.2010.05.003.
- 394 Kalogerakis K.S., Marschall J., Oza A.U., Engel P.A., Meharchand R.T., Wong M.H., (2008),
395 The coating hypothesis for ammonia ice particles in Jupiter: Laboratory experiments and
396 optical modeling. *Icarus* 196:202–215. Doi: doi:10.1016/j.icarus.2008.03.001
- 397 Khare, B.N., Sagan, C., Arakawa, E.T., Suits, F., Callcott, T.A., Williams, M.W., (1984). Optical
398 constants of organic tholins produced in a simulated titanian atmosphere: From soft X-ray
399 to microwave frequencies. *Icarus* 60, 127–137.
- 400 Go, C., de Pater, I., Wong, M., Lockwood, S., Marcus, P., Asay-Davis, X., Shetty, S., (2006).
401 Evolution of the Oval BA during 2004–2005. *Bull. Am. Astron. Soc.* 38, 495.
- 402 Howett, C.J.A., Carlson, R.W., Irwin, P.G.J., et al., (2007). Optical constants of ammonium
403 hydrosulfide ice and ammonia ice. *J. Opt. Soc. Am. B Opt. Phys.* 24 (1), 126–136.
- 404 Ignatiev, N.I., Grassi, D., Zasova, L.V., (2005). Planetary Fourier spectrometer data analysis: fast
405 radiative transfer models. *Planet. Space Sci.* 53, 1035–1042.
406 doi:10.1016/j.pss.2004.12.009.
- 407 Irwin, P.G.J., Weir, A.L., Smith, S.E., Taylor, F.W., Lambert, A.L., Calcutt, S.B.,
408 Cameron-Smith, P.J., (1998). Cloud structure and atmospheric composition of Jupiter
409 retrieved from Galileo near-infrared mapping spectrometer real-time spectra. *J.*
410 *Geophys. Res.* 103, 22001–23021. Irwin, P.G.J., (2009). *Giant Planets of Our Solar*
411 *System. Atmospheres, Composition, and Structure*, second ed. Springer, Praxis
412 Publishing Ltd, Chichester, UK.
- 413 Moreno F. (1996). The Structure of the Stratospheric Aerosol Layer in the Equatorial and South
414 Polar Regions of Jupiter *ICARUS* 124, 632–644.

- Quirantes, A, and Delgado, A.V., (1997). The scattering of light by a suspension of coated spherical particles: effects of polydispersity on cross sections, *J. Phys. D: Appl. Phys.* 30 2123–2131.
- Peek, B.M., (1958). *The Planet Jupiter*. Faber and Faber, London.
- Rodgers, C.D., (2000). *Inverse Methods for Atmospheric Sounding: Theory and Practice*. World Scientific Publishing Co. Pte. Ltd., Singapore.
- Seiff, A., Kirk, D.B., Knight, T.C.D., Young, R.E., Mihalov, J.D., Young, L.A., Milos, F.S., Schubert, G., Blanchard, R.C., Atkinson, D., (1998). Thermal structure of Jupiter’s atmosphere near the edge of a 5-m hot spot in the north equatorial belt. *J. Geophys. Res.* 103, 22857–22890.
- Youssef, A., Marcus, P.S., (2003). The dynamics of jovian white ovals from formation to merger. *Icarus* 162, 74–93.
- Weidenschilling, S.J., Lewis, J.S., (1973). Atmospheric and cloud structures of the jovian planets. *Icarus* 20, 465–476.
- West, R.A., Strobel, D.F., Tomasko, M.G., (1986). Clouds, aerosols and photochemistry in the Jovian atmosphere. *Icarus* 65, 161–217.
- West, R.A., Baines, K.H., Friedson, A.J., Banfield, D., Ragent, B., Taylor, F.W., (2004). Jovian clouds and haze. In: Bagenal, F., Dowling, T., McKinnon, W. (Eds.), *Jupiter: The Planet, Satellites, and Magnetosphere*. Cambridge Univ. Press, Cambridge, pp. 79–104.

Figure Captions

Figure 1: M-filter single image acquired by JIRAM Imager on 28th August 2016 (a), where the three main ovals are clearly visible in the southern hemisphere of Jupiter. Integrated radiance in the 4.30-5.26 μm (b) and 2.40-3.00 μm (c) spectral ranges as acquired by the JIRAM Spectrometer during the first orbit. In panels b and c spectra are averaged in square bins of size 0.5° in planetocentric longitude and latitude. Data are projected in System III West coordinates and blank bins represent no coverage.

Figure 2: Example of two spectral best fits as a result of the data inversion procedure. Black dots and the red line represent the JIRAM measured mean spectrum and the best-fit synthetic spectrum, respectively. The spectra refer to a bin inside the Oval#1 around longitude 115.70°W and latitude 36.54°S (a) and a bin outside the ovals around longitude 75.75°W and latitude 36.68°S (b). The horizontal colored bars highlight the ranges where the gaseous features affect the spectrum.

Figure 3: Maps of the retrieved parameters over the ovals ROI. Blank data represents no coverage or discarded results. Top: Column densities (left) and peak altitudes (right) for the NH_3 -tholin clouds. Middle: Column densities (left) and peak altitudes (right) for the N_2H_4 haze. Bottom: Deep volume mixing ratio (left) and relative humidity (right) for the gaseous ammonia. Column densities and altitudes are expressed in $\text{particles}/\text{cm}^2$ and km above the 1 bar level, respectively, whereas other quantities are dimensionless.

Figure 1.

a

JIRAM IMG M filter – August 28, 2016




**b**JIRAM SPE – Integrated Radiance 4.30–5.26 μm **c**JIRAM SPE – Integrated Radiance 2.40–3.00 μm 

Figure 2.





Figure 3.

