



Publication Year	2019
Acceptance in OA	2020-12-21T10:20:17Z
Title	Mineralogical mapping of Coniraya quadrangle of the dwarf planet Ceres
Authors	RAPONI, Andrea, CARROZZO, FILIPPO GIACOMO, ZAMBON, Francesca, DE SANCTIS, MARIA CRISTINA, CIARNIELLO, Mauro, FRIGERI, ALESSANDRO, Ammannito, E., TOSI, Federico, Combe, J. -Ph., Longobardo, A., PALOMBA, Ernesto, Pieters, C. M., Raymond, C. A., Russell, C. T.
Publisher's version (DOI)	10.1016/j.icarus.2017.10.023
Handle	http://hdl.handle.net/20.500.12386/29037
Journal	ICARUS
Volume	318

Mineralogical mapping of Coniraya Quadrangle of the Dwarf Planet Ceres

A. Raponi^a, F.G. Carrozzo^a, F. Zambon^a, M.C. De Sanctis^a, M. Ciarniello^a, A. Frigeri^a, E. Ammannito^b, F. Tosi^a, J.-Ph. Combe^c, A. Longobardo^a, E. Palomba^a, C.M. Pieters^d, C.A. Raymond^e, C.T. Russell^f.

^a INAF-IAPS Istituto di Astrofisica e Planetologia Spaziali, Via del Fosso del Cavaliere, 100, I-00133 Rome, Italy;

^b Italian Space Agency (ASI), Via del Politecnico snc, I-00133 Rome, Italy;

^c Bear Fight Institute, 22, Fiddler's Road, P.O. Box 667, Winthrop, WA, 98862, USA;

^d Department of Earth, Environmental and Planetary Sciences, Brown University, 324 Brook Street, Providence, RI 02912, USA;

^e NASA/Jet Propulsion Laboratory and California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA;

^f Institute of Geophysics and Planetary Physics, University of California at Los Angeles, 3845 Slichter Hall, 603 Charles E. Young Drive, East, Los Angeles, CA 90095-1567, USA.

Proposed Running Head: Mineralogical analysis of the Ac-H-2 Coniraya quadrangle.

*** Corresponding author:** Dr. Andrea Raponi

INAF-IAPS Istituto di Astrofisica e Planetologia Spaziali

Via del Fosso del Cavaliere, 100, I-00133 Rome, Italy

Email : andrea.raponi@iaps.inaf.it

Abstract

Ceres has been explored by NASA/Dawn spacecraft, which allowed for the discovery of the main mineralogical and compositional characteristics of Ceres' surface. Here, we use mainly data from the Visible and InfraRed imaging spectrometer (VIR) in order to investigate the main spectral characteristics of the quadrangle Ac-H-2 Coniraya, one of the 15 quads in which Ceres' surface has been divided. Coniraya quadrangle is characterized by the presence of mostly highly degraded impact craters of diameters between 50 and 200 km and clusters of small to midsize impact craters. Although the composition over the quadrangle appears to be quite uniform, significant differences have been detected between different craters by spectral parameters analysis and spectral modeling. Ernutet crater presents two regions with very peculiar band at 3.4 μm , typical of organics aliphatic material. One region result to be correlated with larger amount of carbonates, the other region does not present such correlation. Ikapati crater shows strong absorption bands at 4.0 μm , indicating the presence of Na-carbonates in the floor and ejecta. Ikapati, Gaue and other craters present smaller spectral features of NH_4 and/or OH stretching, suggesting a volatile depletion process induced by the heating of the impact event.

1. Introduction

NASA's Dawn spacecraft arrived at dwarf planet 1-Ceres on March 6, 2015, with its scientific payload: the Visible and near-InfraRed imaging spectrometer (VIR) (De Sanctis et al., 2011), the Gamma Ray and Neutron Detector (GRaND) (Prettyman et al., 2011), and the Framing Camera (FC) (Sierks et al., 2011), along with the radio science package (Konopliv et al., 2011).

Here, we use data returned by VIR and FC instruments in order to study the mineralogical composition of the quadrangle Ac-H-2 'Coniraya' (21-66 °N and 0-90 °E), which is one of four quadrangles in Ceres' northern hemisphere.

Ceres' surface shows ubiquitous absorption bands at 2.7 μm (OH stretching in Mg-phyllsilicates) and 3.1 μm related NH₄-phyllsilicates, respectively (De Sanctis et al., 2015; Ammannito et al., 2016).

The thermally-corrected reflectance spectrum of Ceres shows several distinct absorption bands at 3.3-3.5, and 3.95 μm , related to the presence of Mg-carbonates and NH₄-phyllsilicates (De Sanctis et al., 2015).

Although spectral properties of Ceres' surface are quite uniform, there are some peculiar areas where significant differences have been detected in spectral parameters such as slopes, albedo, band depths and band centers. The case that stands out above the surrounding terrains are the bright areas ("Ceralia and Vinalia faculae") in the Occator crater (De Sanctis et al., 2016; Raponi et al., 2017; Longobardo et al., this issue), where the albedo is 5-10 times higher than the average surface. The OH feature in these faculae is shifted from 2.72 to 2.76 μm , indicating the presence of Al-phyllsilicates, while the carbonate features are shifted from 3.95 to 4.0 μm with increasing band depth, suggesting a large abundance of Na-carbonates (De Sanctis et al., 2016). Additionally to the Occator crater, other areas show a signature of Na-carbonates (Carrozzo et al., 2017), e.g., Ikapati crater, located in the 'Coniraya' quadrangle, which is part of the analysis done in the present work.

Knowledge of the mineralogy of Ceres has been enriched by the discovery of organic material mainly in the form of aliphatic material (De Sanctis et al., 2017) near the Ernutet crater (latitude $\sim 53^\circ\text{N}$, longitude 45.5°E , diameter 53.4 km), contained in Coniraya quadrangle.

Oxo crater is another interesting region, which the Coniraya quadrangle shares with the Ac-H-5 'Fejokoo' crater, but which is mostly located in the latter. It is a fresh crater where a large amount of water ice has been detected (Combe et al., 2016; Raponi et al., 2016; Combe et al., this issue). Moreover, it is the second brightest feature on Ceres after the bright material of the Occator crater. Water ice was detected near the southern rim and corresponds to the high albedo materials on the floor of the crater (Combe et al., 2016). The water ice location is entirely in the 'Fejokoo' quadrangle. We refer to Combe et al. (this issue), and Singh et al. (this issue) for further details about this crater.

In section 2, we present the dataset and the methodology used to perform the spectral analysis. In section 3, the thermal emission removal, relevant for the discussion of the spectra in the thermal emission range, is described. In section 4, we give a general view of the quadrangle, discussing the spectral parameters, such as band centers, band depth, spectral slope and albedo. In section 5 we present a quantitative analysis of abundances of the main minerals identified as component of surface materials, by means of the Hapke radiative transfer model. The main findings are discussed in section 6, under the general context of Ceres' surface.




Figure 1. Upper panel: figure adapted from Roatsch et al. (2016), showing the 15-quadrangle scheme in which Ceres' surface has been divided. The Coniraya quadrangle, highlighted in red, is the main focus of this study and is located at 0° - 90°, 21° - 66° N. Bottom: Dawn Framing Camera clear filter LAMO base map mosaic in the Lambert Conical projection, with 35 m/pixel resolution. The names of the main craters are indicated.

2. Data analysis description

The present work uses the dataset acquired by VIR, the mapping spectrometer of the Dawn mission (Russell and Raymond, 2011; De Sanctis et al., 2011). Images provided by the Dawn Framing Camera (Sierks et al., 2011) are also used for context and morphological analysis.

VIR is an imaging spectrometer operating in two channels: the visible channel, ranging between 0.25-1.05 μm , and the infrared channel, between 1.0-5.1 μm . VIR is capable of high spatial (IFOV= 250 $\mu\text{rad/pixel}$, FOV= 64×64 mrad) and spectral ($\Delta\lambda_{\text{VIS}} = 1.8$ nm/band; $\Delta\lambda_{\text{IR}} = 9.8$ nm/band) performances, allowing for the identification of spectral features in order to derive the composition, structure of the surface, and thermal emission.

VIR acquired data of Ceres during all of the mission phases: Survey (spacecraft altitude 4350 km), High Altitude Mapping Orbit (HAMO) (spacecraft altitude 1450 km) and Low Altitude Mapping Orbit (LAMO) (spacecraft altitude 370 km) (Russell and Raymond, 2011). Here, we used the HAMO dataset with nominal spatial resolution of 360-400 m/pix.

The calibrated data (Filacchione et al., 2011, Ammannito et al., 2011) are cleaned for artifacts with the procedure described in Carrozzo et al. (2016). In order to analyze the spectral signature in the range of wavelengths affected by thermal emission (usually from 3.2 μm longward), we use the method described in section 3 to remove the thermal emission. In this paper photometric effects are corrected for both topographic variations and physical characteristics of the regolith using the Hapke approach (Ciarniello et al., 2016). However, other methods for photometric correction can be applied (Longobardo et al., in preparation), yielding similar results, but they are not considered in this work.

To analyze a large dataset like the one considered in this work, we developed an automatic data process able to return different spectral indicators from Ceres observations. This algorithm allows us to map the spatial distribution of each spectral indicator across the surface (for more details about this algorithm, see Frigeri et al., this issue). The main spectral parameters selected are:

- photometrically corrected reflectance at 1.2 μm ,
- spectral slope in the range 1.891-1.163 μm ,
- band depths of phyllosilicate bands at 2.7,
- band depths of NH_4 -phyllosilicate bands at 3.1 μm ,
- band depths at 3.4 μm ,

In the present work we also consider the band depths of carbonates at 3.3-3.5 μm , and 3.9 μm (not computed in the automatic algorithm).

3. Thermal removal

The observed spectra of Ceres' surface are affected by thermal emission from ~ 3.2 μm longward. The radiance of the thermal emission hides the absorption bands and prevents a comparison with laboratory data. We implemented a proper algorithm in order to remove the thermal emission while preserving the spectral features present in the spectra. This is made by modeling the total radiance as the sum of the solar radiance reflected by the surface and the thermal emission of the surface itself, and then performing the removal of the latter, as follows:

- The model of the solar reflected radiance is produced by a model of reflectance of the surface, multiplied by the solar irradiance at Ceres' heliocentric distance. To be consistent with the spectral modeling discussed in Section 5 we used the same reflectance model to estimate the reflectance level in the thermal emission range.
- The Planck function is summed up to this model in order to fit the total radiance level. Free parameters of the Planck function are temperature and emissivity. Their retrieved values are not discussed in the present work.

- Once the Planck function has been derived, we subtract it from the total measured radiance, and we divide the result by the solar irradiance to obtain the reflectance in the whole range.

The reflectance continuum in the thermal emission range, after removal of the Planck function, could be over or underestimated depending on the reflectance model used. However, the measured spectral features are not affected by this uncertainty. Typical result of thermal removal is shown in Fig. 2. The spectral range 3.3 – 4.2 μm , after thermal removal, present the absorption bands of the carbonates at 3.5 μm and 3.95 μm (see also Fig. 8).




Figure 2. Thermal emission removal of a typical spectrum of the average surface of Ceres. Red and black lines are respectively the reflectance spectra before and after thermal emission removal.

4. Spectral analysis

The quadrangle is dominated by mostly highly degraded impact craters of diameters between 50 and 200 km and clusters of small to midsize impact craters. The seven most prominent impact craters have been named Coniraya, Gaue, Omonga, Ikapati, Achita, Ernutet, and Liber (see Fig. 1). Coniraya is the largest impact crater in the quadrangle, while Gaue and Ikapati craters are the freshest large impact craters. We refer to Pasckert et al. (2016) for a complete geological mapping of the Coniraya quadrangle.




Figure 3. The band depth at 2.7 μm shows the abundance of Mg-phyllosilicates.




Figure 4. The band depth at 3.1 μm is indicative of the NH₄-phyllosilicates abundance.




Figure 5. The band depth at 3.4 μm is linked with the abundance of organic materials.




Figure 6. Coniraya photometrically corrected reflectance map at 1.2 μm .




Figure 7. Coniraya 1.891-1.163 μm spectral slope map. To calculate the spectral slope, we use the definition used by Cuzzi et al. (2009), and Filacchione et al. (2012): $\text{Slope} = (R_{\lambda_2} - R_{\lambda_1}) / (\lambda_2 - \lambda_1) / R_{\lambda_1}$, where $\lambda_1 = 1.163 \mu\text{m}$, $\lambda_2 = 1.891 \mu\text{m}$, and R_{λ} is the reflectance at λ .

Maps showing band parameters (Figs.3-7) have been computed from VIR data acquired during the survey and HAMO phase.

We consider thermally corrected spectra as discussed in section 3. Before producing the maps, the Hapke photometric correction has been applied to the VIR data (Hapke et al., 1993; Ciarniello et al., 2016). The maps have been corrected for roughness, single scattering phase function, and opposition effects using parameters derived by Ciarniello et al. (2016). The result of the correction procedure is the reflectance as observed in standard viewing geometry (incidence angle = 30° , emission angle = 0°). The methods used to produce the VIR data mosaic is fully described in Frigeri et al. (this issue). The distribution of the 2.7 μm , 3.1 μm and 3.4 μm band depths are shown in Figs. 3, 4, and 5. The corrected reflectance map at 1.2 μm is shown in Fig. 6, and the spectral slope is shown in Fig. 7.

According to De Sanctis et al. (2015); Ammannito et al. (2016); and De Sanctis et al. (2017), the band depth at 2.7 μm is correlated with Mg-phyllosilicate abundance, the 3.1 μm band depth is correlated with NH_4 -phyllosilicate, and the 3.4- μm band depth in Ernutet is correlated with organic material abundance.

A global trend for the band depth at 2.7 μm (Fig. 3) shows a lower amount of phyllosilicates in crater floors and the crater's ejecta (see also Pasckert et al. (2016)), especially for the three central craters Ernutet, Liber, and Ikapati and for Gaue crater in the Eastern side of the quadrangle. The highest values for these endmembers are found in the western, and southwestern sides of the map, where significant craters are not present.

The band depth at 3.1 μm has almost the same behavior of the 2.7- μm band depth (Fig. 4), apart from two considerations:

- the high latitudes present a systematic deeper band depth than the region at lower latitude;

- the Ernutet crater floor and surrounding terrains show an increase in band depth, differently from the 2.7- μm band depth and all the other large-sized craters.

The deep band depth at 3.4 μm in the Ernutet crater terrain (Fig. 5 and 7C) is an indication of a large abundance of organic aliphatic materials, as discussed in De Sanctis et al. (2017). It is the only case on the whole Ceres' surface where organics features are so evident.

The photometrically-corrected reflectance level at 1.2 μm returns a good correlation with the band depth at 3.4 μm (see Fig. 6). Moreover, the spectral slope appears redder in the Ernutet crater organic-rich areas (Fig. 7 and 8A). This would point to brighter albedo, and reddish spectra of the organic material than the average surface of Ceres.

Conversely, bluer spectra are detected in Ikapati floor and its crater ejecta (see Fig. 7, and 8A) pointing to differences in the composition as discussed in sections 5 and 6.

We extract average spectra from three regions which appear to be the most peculiar within the quadrangle because of their differences with respect the average Ceres spectrum: Ernutet, Ikapati, and Gaue craters. In particular, carbonates absorptions of Ikapati are broader and shifted toward longer wavelengths with respect the other craters (Fig. 8). This is similar to the bright region in the Occator crater, where this shift has been attributed to a change in the carbonates composition, from Mg-Carbonate to Na-carbonates (De Sanctis et al., 2016). The Ikapati region shows Na-carbonates and could thus have been involved in processes similar to those suggested for the Occator crater, which include a recent aqueous alteration (De Sanctis et al., 2016). We refer to Carrozzo et al. (2017) for a complete discussion on the carbonate distribution on Ceres' surface.




Figure 8. Panel A: collected spectra normalized at 1.89 μm , from three different crater of Coniraya quadrangle, and the average Ceres spectrum (Ciarniello et al. 2017) for comparison. Panel B-D: spectral range centered on the main absorption bands, after smoothing and continuum removed. Each spectrum is an average of tens of pixel taken from the region indicated in Panel E.

5. Spectral Modeling

To obtain an information on the abundances of the minerals making up the surface, we used a quantitative spectral analysis of the composition using Hapke's radiative transfer model (Hapke 1993, 2012) as described by Ciarniello et al. (2011) and Raponi et al. (2016). The whole formulation of the bidirectional reflectance (r) is:

$$\text{Eq. 1} \quad r = \frac{SSA}{4\pi} \frac{\mu_0}{\mu + \mu_0} [B_{SH}p(g) + H(SSA, \mu)H(SSA, \mu_0) - 1] \times S(i, e, g, \theta) B_{CB}$$

Where i , e , g are the incidence, emission, and phase angles, respectively, and μ_0 , μ are the cosines of the incidence and emission angles. These parameters come from the shape model and position of the spacecraft at the time of observation. The parameters that contain most of the spectral information are the single scattering albedo (SSA), and the related Ambartsumian–Chandrasekhar functions $H(SSA, \mu)$ describing the multiple scattering components.

Other parameters describe the photometric behavior as a function of the viewing geometry, especially the phase function; they are:

- the single particle phase function $p(g)$,
- the shadow hiding opposition effect $B_{SH}(g)$,
- the coherent back-scattering opposition effect $B_{CB}(g)$,
- the shadow function modelling large-scale roughness $S(i, e, g, \theta)$, with θ being the average surface slope.

These photometric parameters are fixed after Ciarniello et al. (2015), who defined the average scattering properties of Ceres' regolith. The spectral properties are mainly affected by the SSA parameters. The latter have been modeled for an intimate mixing between different minerals, which implies that the particles of the end-member materials are in contact with each other and all are involved in the scattering of a single photon. The SSA of each mineral is defined starting from their grain size and their optical constants as described in Hapke (2012). The optical constants are derived from laboratory measurements (Table 1) with the method described by Carli et al. (2014).

The average SSA of the regolith is defined through the weights p , which represent the relative abundances of the minerals. The weight p_i is defined as the cross section of the grains of the i th mineral as a fraction of the area. p is also a volume fraction, assuming grain sizes are equal for all minerals. Hereafter we refer to the weights p as “abundances”.

$$\text{Eq. 2} \quad SSA = SSA_1p_1 + SSA_2p_2 + SSA_3p_3 + .. \quad \text{with: } p_1 + p_2 + p_3 + .. = 1$$

The best-fitting result is obtained by comparison of the model with the measured spectra, applying the Levenberg–Marquardt method for non-linear least-squares multiple regression (Marquardt 1963).

Free model parameters to be retrieved are:

- abundances of the end-members;
- grain size of the regolith (assumed equal for all end-members);
- a multiplicative constant of the absolute level of reflectance of the model in order to account for uncertainties in the radiometric and photometric accuracies, as well as errors on the local geometry information due to unresolved shadows and roughness;
- a slope added to the model in order to better fit the measured spectrum: in some cases, the measured spectra present an artificial slope where high signal contrast is measured between adjacent pixels, like regions near shadows. This is due to a varying spatial point spread function towards longer wavelengths (Filacchione 2006);

(v) temperature and effective emissivity (Davidsson et al., 2009). The latter is the product of the directional emissivity (Hapke 2012) and a free parameter used to account for unresolved shadow and the structure of the surface (Davidsson et al. 2009). Its interpretation is outside the scope of this work.

The total radiance is modeled by accounting for both the contributions of the reflected sunlight, and the thermal emission:

$$\text{Eq. 3} \quad \text{Rad} = r \times \frac{F_{\odot}}{D^2} + \varepsilon_{eff} \times B(\lambda, T)$$

where r is the Hapke bidirectional reflectance (Eq.1), $F_{\odot}$ is the solar irradiance at 1 AU, D is the heliocentric distance (in AU), ε_{eff} is the effective emissivity, $B(\lambda, T)$ is the Planck function. Thus, the estimation of the thermal emission discussed in Section 3 is done simultaneously with the reflectance modeling in order to yield a consistent result between these two contributions to the total signal measured.

The SSA is modeled starting from minerals already discussed in De Sanctis et al. (2015, 2016) (see Table 1) which are related to the average Ceres surface.

Mineral	Type	Sample ID
Antigorite	Mg-phylosilicate	AT-TXH-007
Dolomite	Mg-Ca-carbonate	CB-EAC-003
NH ₄ -montmorillonite	NH ₄ -phylosilicate	JB-JLB-189
Magnetite	Dark material	MG-EAC-002
Heated Natrite	Na-carbonate	CB-EAC-034-C
Kerite	Organic-aliphatics	MA-ATB-043


Table 1. End-members used in calculating optical constants of the mineral types. Spectra are taken from the Relab spectral database.

On Ceres' surface the predominant component is a dark material, whose identification is challenging, because its spectrum is featureless, except for the tentative absorption band centered at 1 μm , which can be attributed to iron (Fe). We found a good fit with magnetite (Fe_3O_4), however, a large amount of Fe is not consistent with GRAND measurements (Prettyman et al. 2017). More likely dark surface of Ceres should be composed by a large amount of carbon bearing material, being carbonaceous chondrite its close meteoritic analogue (Chapman, C. R. & Salisbury, J. W., 1973, McSween et al., 2017). Moreover, we emphasize that the model used in this work is only based on spectral features, being the absolute signal level of the model adjusted with the multiplicative constant and the additional slope.

We apply the model to the average spectra of Ernutet, Ikapati and Gaue shown in Fig. 8, which turned out to be peculiar from the analysis of the quadrangle (previous section). The best fit are shown in Fig. 9, and the retrieved parameters in table 2.

We also apply the spectral modeling to each pixel of the observations covering the whole area of the three craters. The abundances of the endmembers have been retrieved and projected (Fig. 10-12). The grain size retrieved was in all cases of the order of 100 μm , without clear variations on the maps. Thus we decided to fix this value in the fitting procedure.

353
354



355 **Figure 9.** Measured data with error bars of calibration uncertainties (black), and models (red), for the 4 average
356 spectra shown in Fig. 8A. Retrieved parameters are shown in table 2.


357
358
359

	Ernutet	Ikapati	Gaue	Ceres Average
Mg-phyllsilicates	6.6%	5.8%	6.4%	6.2%
NH ₄ -phyllsilicates	7.5%	3.3%	3.8%	7.0%
Mg-Ca-carbonate	3.6%	2.3%	2.2%	2.2%
Na-carbonate	0.0%	3.3%	0.0%	0.0%
Organics	5.6%	0.0%	0.0%	0.0%


360 **Table 2.** Retrieved abundances from the best fits shown in Fig. 9. In all cases the rest of the composition is a
361 dark neutral material as described in the text.

362
363
364
365
366
367
368
369


370
371
372



373 **Figure 10.** Distributions of the abundances of the modeled endmembers of the Ernutet crater. Panel A: Mg-
374 phyllosilicates, panel B: NH₄-phyllosilicate, panel C: organics, panel D: Mg-Ca-carbonates.
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390



391 **Figure 11.** Distributions of the abundances of the modeled endmembers of the Ikapati crater. Panel A: Mg-
 392 phyllsilicates, panel B: NH₄-phyllsilicate, panel C: Na-carbonates. Other end-members abundances are not
 393 shown because they do not present significant differences with respect the average surface.
 394



395 **Figure 12.** Gaue crater. Mg-phyllsilicates and NH₄-phyllsilicate abundances. Other parameters are not shown
 396 because they do not present significant differences with respect the average surface.
 397
 398

Ernutet

Maps of Fig. 10C shows that abundance distribution of organics in Ernutet crater clearly define two regions with higher concentration:

- a larger region in the south-western rim with the higher concentration on the crater floor, but extending outside the crater
- a smaller region in the northern-western rim, located outside the crater floor.

With the resolution of the map (~400 m/px) we obtain a maximum concentration of 6%. However, we cannot exclude that non-resolved areas can contain more organics. The same consideration is valid for all the other abundances retrieved, and discussed hereafter. The rest of the surface appear to be devoid of organics. However, from the modeling we cannot exclude an amount < 1%.

Larger amount of Mg-phylosilicates (Fig 10A) is mapped inside the floor of Ernutet, up to 8-9%. The surrounding region of the rim are slightly depleted with respect the average Ceres surface. In particular the region with higher concentration of organics present a lower amount of Mg-phylosilicates (5-6%). Differently from the Mg-phylosilicates, organic-rich regions contain larger amount of NH₄-phylosilicates (up to 8-9% as shown in Fig. 10B). Eastern part of the map presents lower amount of NH₄-phylosilicates (4-5%), while the floor of the crater has an heterogeneous amount.

Map of carbonates (Fig. 10D) shows an interesting correlation with the larger/southern organic-rich region. From modeling, carbonates are up to 4.5% in the southern part of the floor. The smaller/northern organic-rich region, does not present such a good correlation with carbonates.

Ikapati

Phyllosilicates maps of Ikapati crater shown in figure 11A and 11B, indicate a clear correlation between OH and NH₄ stretching in this crater. Lower amount has been found in the crater floor and in the western ejecta. In the same areas larger concentration of Na-carbonates has been derived from the model (up to 3%). Those carbonates are absent in the average Ceres surface, and only present in small specific areas (Palomba et al., submitted to Icarus), such as the faculae of the Occator crater.

Gaue

Differently from Ikapati crater, phyllosilicates maps of Gaue (Figs. 12A and 12B) are not well correlated each other: the map of Mg-phylosilicates has an heterogeneous distribution with values ranging from 5 to 10%, while the map of NH₄-phylosilicate clearly presents a lower amount on the crater floor (2-3 %), and about 6% outside it. However, the two end-member are well correlated in a small region in the north/western part of the map, where larger amount of both Mg and NH₄-phylosilicates have been derived: respectively 10% and 7%.

6. Discussion and Conclusions

From the analysis of the Coniraya quadrangle, some relevant characteristics emerge. Here we discuss the major findings in the general context of Ceres' surface.

Phyllosilicates distribution

The band depth at 2.7 and 3.1 μm mapped in this work are both indicative of phyllosilicate abundance, and relative differences in their distributions are detectable (e.g., between Ikapati and Gaue craters). This was also highlighted by Ammannito et al. (2016) as a general result for Ceres' surface.

The floor of the freshest and largest craters present smaller band depth at 2.7 μm and 3.1 μm .

The modeling performed for Ernutet, Ikapati, and Gaue craters reveals that the smaller band depth can be interpreted as a lower amount of Mg/NH₄-phylosilicates. Smaller phyllosilicates band associates with relatively young crater seems quite common on Ceres' surface (De Sanctis et al., this issue),

which may be the result of a volatile depletion due to impact heating. A noticeable exception is represented by the large-scale depression of Vendimia Planitia, located in the eastern adjacent quadrangles. The measured phyllosilicate absorptions at 2.7 and 3.1 μm , in two main craters located in this area, Dantu (Stephan et al., this issue) and Kerwan (Palomba et al., this issue), are stronger than its surrounding areas, and they actually are the deepest phyllosilicate absorptions measured on Ceres (Stephan et al., this issue). This observed trend may reveal a change in the composition of Ceres' crust with increasing depth below the topographic mean (Stephan et al., this issue).

The Mg and NH_4 -phyllosilicates modeled present different depletion in the three craters mapped in Fig. 10-12. The cases that stand out are the high contrast for Ikapati crater floor and ejecta (Fig. 11A, 11B), and the NH_4 -phyllosilicates on the Gaue crater floor (Fig. 12B).

This would point to a complex picture in which different processes can concur to determine the volatiles abundances during the impact (e.g. energy released by the impactor, depth of excavation), after the impact (e.g. aqueous alteration, hydrothermal activity, lateral mixing by micrometeoritic impacts, space weather), and could also reflect heterogeneities on Ceres' surface preexistent to the impact event.

Ikapati crater and ejecta

As discussed in the previous section, Ikapati fresh crater shows slightly smaller band depth at 2.7 μm and 3.1 μm , which would mean a lower concentration of Mg/ NH_4 -phyllosilicates. By taking into account the average spectrum shown in Fig. 10, we notice other relevant differences: larger absorption band of carbonates, and bluer spectral slope. Moreover, the band center of the carbonates absorption band shows a shift from 3.95 to 4.0 μm (see Fig. 8D), similar to the bright region in the Occator crater, where this shift has been attributed to a change in the carbonates composition, from Mg-Carbonate to Na-Carbonates (De Sanctis et al., 2016; Longobardo et al., this issue). The Ikapati region could thus have been involved in similar processes, as suggested for the Occator crater, which include a recent aqueous alteration (De Sanctis et al., 2016). However, in Ikapati the Na-carbonate is spread in the floor and ejecta, rather than being concentrated in specific regions inside the crater floor. This can be the result of the impact event of Ikapati that spread bright material previously concentrated in faculae like in Occator crater. This hypothesis would point to a unique formation process for all bright material present on the surface as suggested by (Stein et al., submitted), being their differences due to the following story of impact events.

Contrarily to the slow process occurring to form the bright material, the volatile depletion due to the impact induced heating could be fast, and it can explain the correlation of the lower amount of hydrated (OH- stretching) and ammoniated phyllosilicates (NH_4 -stretching) in correspondence of the ejecta.

Ernutet crater and organic-rich regions

The Ernutet crater is a unique feature of Ceres' surface, revealing a large amount of organic material that is mapped in this work by the 3.4- μm absorption band, and the abundance map of Fig. 10C. This absorptions is characteristic of the symmetric and antisymmetric stretching frequencies of methyl (CH_3) and methylene (CH_2) functional groups, typical of aliphatic hydrocarbons (F.-R. Orthous-Daunay et al., 2013). The model performed in the present work (Fig 9A) and in (De Sanctis et al. 2016) shows that Ceres organics signatures are very similar with the organic bands of terrestrial hydrocarbons such as kerite.

An evident correlation with organics distribution has been found with the absolute level of reflectance at 1.2 μm (Fig. 6), and the reddish slope (Fig. 7), revealing a possible constraint on the optical properties of the organic material: brighter and more reddish than the average surface of Ceres.

The abundance map of organics shown in Fig. 10C clearly define two regions with higher concentration of organics. In the work of De Sanctis et al. (2016) these two regions were labeled as region A (southern) and B (northern). In the same two regions we derived:

- 1) smaller abundance of Mg-phyllsilicates (Fig. 10A);
- 2) slightly larger abundance of NH₄-phyllsilicates (Fig. 10B);
- 3) only in the southern region larger abundance of Mg-Ca-carbonates (Fig. 10D);

De Sanctis et al. 2016 suggested an endogenous origin of the organics, on the base of the larger presence of carbonates, (and slight larger presence of NH₄-phyllsilicates), both being products of an hydrothermal activity together with the organics. An exogenous origin would require further explanations about the correlation with minerals (carbonates, NH₄-phyllsilicates) extensively present on Ceres surface, but unseen in the asteroid belt with the same amount.

The uniqueness of such organic-rich region would point to an uncommon condition over Ceres' surface. However, the asymmetry of the retrieved parameters in the two regions (point 3), would point to formation processes under different thermophysical and/or chemical conditions, occurring very close each other, but both favorable for the organics formation.

Further efforts are needed to assess the mechanism responsible for the formation and transport to the surface of minerals which should have a different origin with respect the common Ceres minerals, like organics and Na-carbonates.

The uniqueness of specific places on Ceres surface, together with the diversity of mineralogy is an indication of a complex environment typical of an active body.

Acknowledgments

VIR is funded by the Italian Space Agency-ASI and was developed under the leadership of INAF-Istituto di Astrofisica e Planetologia Spaziali, Rome-Italy. The instrument was built by Selex-Galileo, Florence-Italy. The authors acknowledge the support of the Dawn Science, Instrument, and Operations Teams. This work was supported by ASI and NASA.

References

Ammannito, E., De Sanctis, M.C., Ciarniello, M., Frigeri, A., Carrozzo, F.G., Combe, J.-Ph., Ehlmann, B.L., Marchi, S., McSween, H.Y., Raponi, A., Toplis, M.J., Tosi, F., Castillo-Rogez, J.C., Capaccioni, F., Capria, M.T., Fonte, S., Giardino, M., Jaumann, R., Longobardo, A., Joy, S.P., Magni, G., McCord, T.B., McFadden, L.A., Palomba, E., Pieters, C.M., Polanskey, C.A., Rayman, M.D., Raymond, C.A., Schenk, P.M., Zambon, F., Russell, C.T., 2016. Distribution of phyllosilicates on the surface of Ceres. *Science* 353 (6303), id.aaf4279. Doi: 10.1126/science.aaf4279.

Carrozzo, F.G., Raponi, A., De Sanctis, M.C., Ammannito, E., Giardino, M., D'Aversa, E., Fonte, S., Tosi, F., 2016. Artifacts reduction in VIR/Dawn data. *Rev. Sci. Instrum.* 87 (12), id.124501. Doi: 10.1063/1.4972256.

539

540 Carrozzo, F.G., De Sanctis, M.C., Raponi, A., Ammannito, E., Castillo-Rogez, J.C., Ehlmann, B.L.,
541 Marchi, S., Stein, N., Ciarniello, M., Tosi, F., Capaccioni, F., Capria, M.T., Fonte, S., Formisano, M.,
542 Frigeri, A., Giardino, M., Longobardo, A., Magni, G., Palomba, E., Zambon, F., Raymond, C.A.,
543 Russell, C.T., 2017a. Nature, formation and distribution of Carbonates on Ceres. Submitted to Science.
544 Under review.

545

546 Chapman, C. R. & Salisbury, J. W. Comparisons of meteorite and asteroid spectral reflectivities. *Icarus*
547 19, 507–522 (1973).
548

549 Ciarniello, M., De Sanctis, M.C., Ammannito, E., Raponi, A., Longobardo, A., Palomba, E., Carrozzo,
550 F.G., Tosi, F., Li, J.-Y., Schröder, S.E., Zambon, F., Frigeri, A., Fonte, S., Giardino, M., Pieters, C.M.,
551 Raymond, C.A., Russell, C.T., 2017. Spectrophotometric properties of dwarf planet Ceres from the
552 VIR spectrometer on board the Dawn mission. *Astron. Astrophys.* 598, id.A130. Doi: 10.1051/0004-
553 6361/201629490.

554

555 Combe, J.-Ph., McCord, T.B., Tosi, F., Ammannito, E., Carrozzo, F.G., De Sanctis, M.C., Raponi, A.,
556 Byrne, S., Landis, M.E., Hughson, K.H.G., Raymond, C.A., Russell, C.T., 2016. Detection of local H₂O
557 exposed at the surface of Ceres. *Science* 353 (6303), id.aaf3010. Doi: 10.1126/science.aaf3010.

558

559 Combe, J.-Ph., Raponi, A., Tosi, F., De Sanctis, M.C., Carrozzo, F.G., Zambon, F., Ammannito, E.,
560 Hughson, K.H.G., Nathues, A., Hoffmann, M., Platz, T., Thangjam, G., Schorghofer, N., Schröder,
561 S.E., Byrne, S., Landis, M.E., Ruesch, O., McCord, T.B., Johnson, K.E., Singh, S.M., Raymond, C.A.,
562 Russell, C.T., 2017a. Exposed H₂O-rich areas detected on Ceres with the Dawn Visible and InfraRed
563 mapping spectrometer. Submitted to *Icarus* (this issue). Under review.

564

565 De Sanctis, M.C., Coradini, A., Ammannito, E., Filacchione, G., Capria, M.T., Fonte, S., Magni, G.,
566 Barbis, A., Bini, A., Dami, M., Fikai-Veltroni, I., Preti, G., and the VIR Team, 2011. The VIR
567 Spectrometer. *Space Sci Rev.* 163, Issue 1–4, 329-369. Doi: 10.1007/s11214-010-9668-5.

568

569 De Sanctis, M.C., Ammannito, E., Raponi, A., Marchi, S., McCord, T.B., McSween, H.Y., Capaccioni,
570 F., Capria, M.T., Carrozzo, F.G., Ciarniello, M., Longobardo, A., Tosi, F., Fonte, S., Formisano, M.,
571 Frigeri, A., Giardino, M., Magni, G., Palomba, E., Turrini, D., Zambon, F., Combe, J.-Ph., Feldman,
572 W., Jaumann, R., McFadden, L.A., Pieters, C.M., Prettyman, T., Toplis, M., Raymond, C.A., Russell,
573 C.T., 2015. Ammoniated phyllosilicates with a likely outer Solar System origin on (1) Ceres. *Nature*
574 528 (7581), 241-244. Doi: 10.1038/nature16172.

575

576 De Sanctis, M.C., Raponi, A., Ammannito, E., Ciarniello, M., Toplis, M.J., McSween, H.Y., Castillo-
577 Rogez, J.C., Ehlmann, B.L., Carrozzo, F.G., Marchi, S., Tosi, F., Zambon, F., Capaccioni, F., Capria,
578 M.T., Fonte, S., Formisano, M., Frigeri, A., Giardino, M., Longobardo, A., Magni, G., Palomba, E.,

579 McFadden, L.A., Pieters, C.M., Jaumann, R., Schenk, P., Mugnuolo, R., Raymond, C.A., Russell, C.T.,
580 2016. Bright carbonate deposits as evidence of aqueous alteration on (1) Ceres. *Nature* 536 (7614), 54-
581 57. Doi: 10.1038/nature18290.

582

583 De Sanctis, M.C., Ammannito, E., McSween, H.Y., Raponi, A., Marchi, S., Capaccioni, F., Capria,
584 M.T., Carrozzo, F.G., Ciarniello, M., Fonte, S., Formisano, M., Frigeri, A., Giardino, M., Longobardo,
585 A., Magni, G., McFadden, L.A., Palomba, E., Pieters, C.M., Tosi, F., Zambon, F., Raymond, C.A.,
586 Russell, C.T., 2017a. Localized aliphatic organic material on the surface of Ceres. *Science* 355 (6326),
587 719-722. Doi: 10.1126/science.aaj2305.

588

589 Frigeri, A. De Sanctis, M.C., Ammannito, E., Tosi, F., Ciarniello, M., Zambon, F., Carrozzo, F.G.,
590 Raponi, A., McCord, T.B., Raymond, C.A., Russell, C.T., 2017. The spectral parameter maps of Ceres
591 from NASA/DAWN VIR data. Submitted to *Icarus* (this issue). Under review.

592

593 Hapke B., Theory of reflectance and emittance spectroscopy, 1993.

594

595 Hapke B., 1993, Theory of reflectance and emittance spectroscopy Second Edition, 2012

596

597 Konopliv, A. S.; Asmar, S. W.; Bills, B. G.; Mastrodemos, N.; Park, R. S.; Raymond, C. A.;
598 Smith, D. E.; Zuber, M. T. The Dawn Gravity Investigation at Vesta and Ceres. *Space Science*
599 *Reviews*, 163, pp. 461-486

600

601 Longobardo, A., et al., 2017. Mineralogy of the Occator quadrangle. Submitted to *Icarus* (this issue).
602 Under review.

603

604 McSween, H. Y.; Castillo-Rogez, J.; Emery, J. P.; De Sanctis, M. C.; Dawn Science Team.
605 Rationalizing the Composition and Alteration of Ceres. LPI Contribution No. 1903, p.1258, 2016.

606

607 Orthous-Daunay, F.-R.; Quirico, E.; Beck, P.; Brissaud, O.; Dartois, E.; Pino, T.; Schmitt, B., Mid-
608 infrared study of the molecular structure variability of insoluble organic matter from primitive
609 chondrites. *Icarus* 223, 534–543, 2013.

610

611 Palomba, E., Longobardo, A., De Sanctis, M.C., Carrozzo, F.G., Galiano, A., Zambon, F., Raponi, A.,
612 Ciarniello, M., Stephan, K., Williams, D., Ammannito, E., Capria, M.T., Fonte, S., Giardino, M., Tosi,
613 F., Raymond, C.A., Russell, C.T., 2017. Mineralogical mapping of the Kerwan quadrangle on Ceres.
614 Submitted to *Icarus* (this issue). Under review.

615

616 Pasckert, J. H.; Hiesinger, H.; Williams, D. A.; Crown, D. A.; Mest, S. C.; Buczkowski, D. L.; Scully,
 617 J. E. C.; Schmedemann, N.; Jaumann, R.; Roatsch, T.; Preusker, F.; Nass, A.; Nathues, A.; Hoffmann,
 618 M.; Schäfer, M.; De Sanctis, M. C.; Raymond, C. A.; Russell, C. T. Geologic Mapping of the Ac-H-2
 619 Coniraya Quadrangle of Ceres from NASA's Dawn Mission. 47th Lunar and Planetary Science
 620 Conference, LPI Contribution No. 1903, p.1450, 2016.
 621
 622 N. Stein, B. L. Ehlmann, E. Palomba, M. C. De Sanctis, A. Nathues, H. Hiesinger, E. Ammannito, C.
 623 A. Raymond, R. Jaumann, A. Longobardo, C. T. Russell, The Formation and Evolution of Bright Spots
 624 on Ceres, submitted to Icarus.
 625
 626 Prettyman, Thomas H.; Feldman, William C.; McSween, Harry Y.; Dingler, Robert D.;
 627 Enemark, Donald C.; Patrick, Douglas E.; Storms, Steven A.; Hendricks, John S.;
 628 Morgenthaler, Jeffery P.; Pitman, Karly M.; Reedy, Robert C. Dawn's Gamma Ray and Neutron
 629 Detector. Space Science Reviews, 163, pp. 371-459
 630
 631 Raponi, A.; Ciarniello, M.; Capaccioni, F.; Filacchione, G.; Tosi, F.; De Sanctis, M. C.; Capria, M. T.;
 632 Barucci, M. A.; Longobardo, A.; Palomba, E.; Kappel, D.; Arnold, G.; Mottola, S.; Rousseau, B.;
 633 Rinaldi, G.; Erard, S.; Bockelee-Morvan, D.; Leyrat, C. The temporal evolution of exposed water ice-
 634 rich areas on the surface of 67P/Churyumov–Gerasimenko: spectral analysis. MNRAS 462, 2016.
 635
 636 Roatsch, T., Kersten, E., Matz, K.-D., Preusker, F., Scholten, F., Jaumann, R., Raymond, C.A., Russell,
 637 C.T., 2016a. Ceres survey atlas derived from Dawn Framing Camera images. Planet. Space Sci. 121,
 638 115-120. Doi: 10.1016/j.pss.2015.12.005.
 639
 640 Roatsch, T., Kersten, E., Matz, K.-D., Preusker, F., Scholten, F., Jaumann, R., Raymond, C.A., Russell,
 641 C.T., 2016b. High-resolution Ceres High Altitude Mapping Orbit atlas derived from Dawn Framing
 642 Camera images. Planet. Space Sci. 129, 103-107. Doi: 10.1016/j.pss.2016.05.011.
 643
 644 Russell, C. T.; Raymond, C. A. The Dawn Mission to Vesta and Ceres. Space Science Reviews, 163,
 645 pp. 3-23, 2011.
 646
 647 Sierks, H., Keller, H.U., Jaumann, R., Michalik, H., Behnke, T., Bubenhausen, F., Büttner, I., Carsenty,
 648 U., Christensen, U. Enge, R., Fiethe, B., Gutiérrez-Marqués, P., Hartwig, H., Krüger, H., Kühne, W.,
 649 Maue, T., Mottola, S., Nathues, A., Reiche, K.-U., Richards, M.L., Roatsch, T., Schröder, S.E.,
 650 Szemerey, I., Tschentscher, M., 2011. The Dawn Framing Camera. Space Sci Rev. 163, Issue 1–4, 263-
 651 327. Doi: 10.1007/s11214-011-9745-4.
 652
 653 Singh, S., Combe, J.-Ph., McFadden, L.A., McCord, T.B., Johnson, K.E., Hughson, K.H.G., Zambon,

654 F., Ciarniello, M., Carrozzo, F.G., Ammannito, E., De Sanctis, M.C., Ruesch, O., Stephan, K., Tosi, F.,
655 Longobardo, A., Raymond, C.A., Russell, C.T., 2017. Mineralogy mapping of the Ac-H-5 Fejokoo
656 quadrangle of Ceres. Submitted to Icarus (this issue). Under review.

657

658 Stephan, K., Jaumann, R., Zambon, F., Carrozzo, F.G., De Sanctis, M.C., Tosi, F., Ammannito, E.,
659 Longobardo, A., Palomba, E., McFadden, L.A., Krohn, K., Williams, D.A., Raponi, A., Ciarniello, M.,
660 Combe, J.-Ph., Frigeri, A., Roatsch, T., Matz, K.-D., Preusker, F., Raymond, C.A., Russell, C.T.,
661 2017b. Spectral investigation of quadrangle Ac-H-3 of the dwarf planet Ceres – The region of impact
662 crater Dantu. Submitted to Icarus (this issue). Under review.

663

664 Tosi, F., Carrozzo F.G., Zambon F., Ciarniello M., Frigeri A., Combe J.-Ph., De Sanctis M.C.,
665 Hoffmann M., Longobardo A., Nathues A., Raponi A., Ammannito E., Krohn K., McFadden L.A.,
666 Palomba E., Pieters C.M., Stephan K., Raymond C.A., Russell C.T., and the *Dawn* Science Team.
667 Mineralogical analysis of the Haulani quadrangle of the dwarf planet Ceres. Submitted to Icarus (this
668 issue). Under review.