



|                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|----------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Publication Year</b>          | 2021                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Acceptance in OA</b>          | 2023-01-26T14:28:06Z                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Title</b>                     | GAUSS – Genesis of Asteroids and evolUtion of the Solar System. A sample return mission to Ceres.                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Authors</b>                   | Shi, Xian, Castillo-Rogez, Julie, Hsieh, Henry, Hui, Hejiu, Ip, Wing Huen, Lei, Hanlun, Li, Jian Yang, TOSI, Federico, Zhou, Liyong, Agarwal, Jessica, Barucci, Antonella, Beck, Pierre, Bagatin, Adriano Campo, CAPACCIONI, FABRIZIO, Coates, Andrew J., CREMONESE, Gabriele, Duffard, Rene, Grande, Manuel, Jaumann, Ralf, Jones, Geraint H., Kallio, Esa, Lin, Yangting, Mousis, Olivier, Nathues, Andreas, Oberst, Jürgen, Sierks, Holger, Ulamec, Stephan, Wang, Mingyuan |
| <b>Publisher's version (DOI)</b> | 10.1007/s10686-021-09800-1                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Handle</b>                    | <a href="http://hdl.handle.net/20.500.12386/33082">http://hdl.handle.net/20.500.12386/33082</a>                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Journal</b>                   | EXPERIMENTAL ASTRONOMY                                                                                                                                                                                                                                                                                                                                                                                                                                                         |

# GAUSS - Genesis of Asteroids and Evolution of the Solar System

## A Sample Return Mission to Ceres

Xian Shi · Julie Castillo-Rogez · Henry Hsieh · Hejiu Hui · Wing-Huen Ip · Hanlun Lei · Jian-Yang Li · Federico Tosi · Liyong Zhou · Jessica Agarwal · Antonella Barucci · Pierre Beck · Adriano Campo Bagatin · Fabrizio Capaccioni · Andrew J. Coates · Gabriele Cremonese · Rene Duffard · Manuel Grande · Ralf Jaumann · Geraint H. Jones · Esa Kallio · Yangting Lin · Olivier Mousis · Andreas Nathues · Jürgen Oberst · Adam Showman · Holger Sierks · Stephan Ulamec · Mingyuan Wang · and The GAUSS Team

Received: date / Accepted: date

---

X. Shi · J. Agarwal · H. Sierks · A. Nathues  
Max Planck Institute for Solar System Research, Göttingen, Germany  
E-mail: shi@shao.ac.cn  
*Present address: Shanghai Astronomical Observatory, Shanghai, China* for X. Shi

J. Castillo-Rogez  
NASA Jet Propulsion Laboratory, CA, USA

H. Hsieh · J.-Y Li  
Planetary Science Institute, AZ, USA

H. Hui  
School of Earth Sciences and Engineering, Nanjing University, Nanjing, China

W.-H Ip  
Institute of Astronomy and Space Science, National Central University, Chung Li, Taiwan

H. Lei · L. Zhou  
School of Astronomy and Space Science, Nanjing University, Nanjing, China

F. Tosi · F. Capaccioni  
INAF-IAPS, Istituto di Astrofisica e Planetologia Spaziali, Rome, Italy

J. Agarwal  
Technical University Braunschweig, Institute for Geophysics and Extraterrestrial Physics, Braunschweig, Germany

A. Barucci  
LESIA-Observatoire de Paris, Université PSL, CNRS, Université de Paris, Sorbonne Uni-

**Abstract** The goal of Project GAUSS (Genesis of Asteroids and evolUtion of the Solar System) is to return samples from the dwarf planet Ceres. Ceres is the most accessible candidate of ocean worlds and the largest reservoir of water in the inner Solar System. It shows active volcanism and hydrothermal activities in recent history. Recent evidence for the existence of a subsurface ocean on Ceres and the complex geochemistry suggest past habitability and even the potential for ongoing habitability. GAUSS will return samples from Ceres with the aim of answering following top-level scientific questions:

- What is the origin of Ceres and what does this imply for the origin of water and other volatiles in the inner Solar System?
- What are the physical properties and internal structure of Ceres? What do they tell us about the evolutionary and aqueous alteration history of dwarf planets?
- What are the astrobiological implications of Ceres? Is it still habitable today?
- What are the mineralogical connections between Ceres and our current collections of carbonaceous meteorites?

---

versité, F-92195 Meudon Principal Cedex, France.

P. Beck

Univ. Grenoble Alpes, CNRS Institut de Planétologie et d'Astrophysique, Grenoble, France

A. C. Bagatin

Universidad de Alicante, Departamento de Física, Ingeniería de Sistemas y Teoría de la Señal. Alicante, Spain

A. J. Coates · G. H. Jones

Mullard Space Science Laboratory, University College London, Surrey, UK

G. Cremonese

INAF - Osservatorio Astronomico di Padova, Padova, Italy

Rene Duffard

Instituto de Astrofísica de Andalucía (CSIC), Granada, Spain

M. Grande

University of Aberystwyth, Aberystwyth, UK

Ralf Jaumann · J. Oberst

DLR Institute of Planetary Research, Berlin, Germany

E. Kallio

School of Electrical Engineering, Aalto University, Aalto, Finland

Y. Lin

Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, China

O. Mousis

Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France

A. Showman

Lunar and Planetary Laboratory, University of Arizona, AZ, USA

S. Ulamec

German Aerospace Centre (DLR), Cologne, Germany

M. Wang

National Astronomical Observatory, Chinese Academy of Science, Beijing, China

**Keywords** Ceres · Dwarf planet · Sample return · Ocean world · Habitability · Voyage 2050

## 1 Background

Though the Rosetta mission to comet 67P/Churyumov-Gerasimenko came to an end only recently, in 2016, it is important to recall that the planning activity eventually leading to its approval by ESA was initiated more than three decades earlier in 1983. An equally, if not more, ambitious project in the framework of “Voyage 2050” is proposed here. The target is the innermost dwarf planet, Ceres, which was discovered on New Year’s Day of 1801, by the Italian astronomer Giuseppe Piazzi in Palermo Observatory. At the time of its discovery, Ceres was considered to be the missing planet between the orbits of Mars and Jupiter as predicted by the Titius-Bode law (Grant, 1852). Ceres’ location was confirmed in December the same year using the orbital elements calculated by then 24-year-old Carl Fredrich Gauss (Teets and Whitehead, 1999). The name of the project with the acronym of GAUSS for “Genesis of Asteroids and evolUtion of the Solar System” is partly a tribute to this scientific episode of great importance in astronomy and planetary science.

## 2 Scientific Rationale

### 2.1 Ceres in the history of the Solar System

The formation of the oldest solids found in the Solar System – calcium-aluminium inclusions (CAIs) – dates back 4567.5 Myr (Connelly et al., 2008). After that, the formation of gas giants must finish before the dispersal of the protoplanetary gas disc, which lasts 2-10 Myr (Hillenbrand, 2008; Bell et al., 2013). Then, about 30-100 Myr later, the terrestrial planets formed in the planetesimal disc (Jacobson et al., 2014).


The planetesimal disc not only provided all the materials that constitute the terrestrial planets and cores of gas giants, but also exerts perturbing forces on the grown planets (Kokubo and Ida, 1995; Hahn and Malhotra, 1999). This perturbation may continuously modify the planetary orbits in a relatively gentle way (Fernandez and Ip, 1984). Before the dispersal of the gas disc, planets could already experience migration due to the interaction with the massive gas disc (Walsh et al., 2011). During the planetary migration afterwards, mechanisms such as low-order mean motion resonance between planets brought up abrupt variations in the orbital configuration of planets (Tsiganis et al., 2005). The interaction between the planetesimal disc and migrating planets/protoplanets reconfigured the structure of the Solar System (Morbidelli et al., 2005; Gomes et al., 2005; Walsh et al., 2011). However, the growth and evolution of terrestrial planets and asteroids (in a wide radial range from Mars orbit to the Kuiper belt) in the early stages of the Solar System are not fully understood.

Given that their properties depend on their formation circumstances and evolutionary processes experienced since their formation, small bodies (e.g., asteroids, comets, and trans-Neptunian objects) in the Solar System are of interest to researchers because they provide a way to probe the protoplanetary disc from which our Solar System formed (Scott et al., 1989; Jones et al., 1990; Bottke et al., 2002). By determining the chemical and physical properties of various small bodies in our Solar System, we can gain insight into chemical and thermal conditions in different areas of the protoplanetary disc and also investigate the chemical, thermal, collisional, and dynamical processes that have shaped those populations since their formation. A successful scenario of planet migration should provide the mechanisms that were able to efficiently deliver planetesimals from different zones to the main belt. In addition, the structure of the main belt, the mass depletion, the proper excitation of orbital eccentricities and inclinations, should all be reproduced.

One of the earliest attempts to map the compositional structure of our asteroid belt and use it to infer information about the origin and evolution of the Solar System was performed by Gradie and Tedesco (1982). They found a systematic distribution of compositional types of asteroids in the main asteroid belt that they suggested was consistent with chemical condensation models of the Solar System (Fig. 1). The authors concluded that it was unlikely that this distribution could be explained by the chaotic transport of objects from different regions of the Solar System into the asteroid belt, and proposed instead that it indicated that the asteroids formed at or near their present locations.

The work of Gradie and Tedesco (1982) has since undergone a major update in the form of a study by DeMeo and Carry (2013), who used the Sloan Digital Sky Survey (SDSS) Moving Object Catalog data to derive taxonomic classifications for about 35000 objects, including objects as small as 5 km in diameter. The inclusion of much smaller objects in this sample than were previously available, as well as the computation of compositional distributions by mass (instead of by number, see Fig. 2 in DeMeo and Carry, 2014) suggest that the asteroid belt preserves a history of Solar System evolution that is far more complex than previously thought (Gomes et al., 2005; Levison et al., 2009; Walsh et al., 2011; DeMeo and Carry, 2014), with asteroids of various taxonomic types scattered throughout the asteroid belt, including in regions where they are not expected based on dynamically static Solar System formation models. Instead, the history of the Solar System as recorded by main-belt asteroids likely includes relatively brief periods of dramatic mixing caused by giant planetary migration (e.g., those proposed as parts of the well-known Nice and Grand Tack models (Tsiganis et al., 2005; Morbidelli et al., 2005; Tsiganis et al., 2005; Levison et al., 2009, 2011; Walsh et al., 2011) as well as less dramatic but ongoing processes such as collisions and small body migration due to mean-motion resonances with the giant planets and the Yarkovsky effect (e.g., Bottke et al., 2005; Bottke Jr et al., 2006; Gladman et al., 1997; Farinella et al., 1998; Raymond and Izidoro, 2017).

The fraction and composition of ice in a body is of particular interest due to the strong temperature constraints they provide. The so-called “snow



**Fig. 1** Relative distribution of taxonomic types for a bias-corrected sample of 656 main-belt asteroids. Letters indicate different compositional types. Refer to Table 1 in Gradie and Tedesco (1982) for the definition of each type. Locations of major resonances with Jupiter are indicated at the top. (Figure reprinted from Gradie and Tedesco, 1982).

line” refers to the distance from the Sun at which the temperature dips below the condensation temperature of water, causing it to freeze into solid ice and then become incorporated into accreting planetesimals. The exact location of the snow line in our protoplanetary disc was dependent on a variety of poorly constrained environmental conditions including opacity, mass density, and accretion rate in the disc, and is also thought to have shifted with time as planetesimal accretion progressed and the aforementioned properties of the disc changed (Morbidelli et al., 2016).

Bodies formed in the outer Solar System such as between Jupiter and Neptune (the original accretion zone of current Oort Cloud objects; Hahn and Malhotra 1999, Weissman 1999) and beyond the orbit of Neptune (the Kuiper Belt) are well beyond the snow line. Closer to the Sun, the situation is less certain. Observations of asteroids suggest that the snow line probably existed as close as 2.5 au from the Sun (Gradie and Tedesco, 1982; Jones et al., 1990), but theoretical studies (Sasselov and Lecar, 2000; Ciesla and Cuzzi, 2006; Lecar et al., 2006) have placed it as close in as the orbit of Mars, or even closer. If this is true, it means that objects throughout the main asteroid belt could have incorporated some water ice at the time of their formation.


Evidence of past and present-day ice has in fact been found in main-belt asteroids. Studies of meteorites linked to the asteroid belt as well as remote spectroscopic observations of asteroids have revealed the presence of aqueously altered minerals, indications that liquid water was once present (Hiroi

et al., 1996; Keil, 2000; Rivkin et al., 2002). Meanwhile, spectroscopic evidence of water ice frost has been reported for various main-belt asteroids including (24) Themis (Rivkin and Emery, 2010; Takir and Emery, 2012; Hargrove et al., 2015), while some main-belt objects have even been observed to exhibit comet-like activity that has been attributed to the sublimation of present-day volatile ices, i.e., the so-called main-belt comets (Hsieh and Jewitt, 2006). The location, abundance, distribution, and inferred water content of currently and formerly icy main-belt objects provide valuable clues for discerning the primordial location and evolution of the snow line, although this of course is also complicated by the aforementioned likely transport of at least some small bodies from their original formation locations due to giant planet migration and other dynamical processes.

The bulk-rock isotope anomalies of meteorites reveal an isotopic dichotomy: two distinct trends defined by the non-carbonaceous (NC) bodies and the carbonaceous (CC) bodies respectively (Fig. 2). The CC bodies include carbonaceous chondrites, a few ungrouped achondrites, and IIC, IID, IIF, IIIF, IVB iron meteorites. The NC bodies include Earth, the Moon, Mars, ordinary chondrites, enstatite chondrites, and most of achondrites. This dichotomy has been further observed in Mo, W, Ru, and Ni isotopic systematics (Budde et al., 2016; Kruijer et al., 2017; Nanne et al., 2019; Poole et al., 2017). However, the origins of the isotopic anomalies and their correlations are unknown, although  $^{54}\text{Cr}$ -enrichment can be attributed to nano-sized Cr-oxide probably ejected from supernovas (Qin et al., 2011). Based on the isotopic dichotomy, Warren (2011) proposed that the CC bodies formed in the outer Solar System and the NC bodies formed in the inner Solar System. The formation of Jupiter blocked the material exchange between the two groups. As a consequence of radial migration and mass growth of the giant planets in the Solar System, some CC planetesimals and embryos moved inwards from the outer Solar System to the main belt (Walsh et al., 2011; Warren, 2011; Kruijer et al., 2017; Raymond and Izidoro, 2017).

Ceres, the largest object in the main belt, preserves many clues to the formation and evolution of the main belt, as well as of the Solar System. The surface material of Ceres is similar to carbonaceous chondrite, though no meteorite has so far been linked with Ceres (Rivkin et al., 2011). Interpretation of the data from the Dawn mission with geochemical simulations show that the surface mineralogy of Ceres is consistent with the aqueous alteration of CI and CM chondrite (Castillo-Rogez et al., 2018).

The ammonia detected on Ceres (De Sanctis et al., 2015) suggests that Ceres could have originated in the outer Solar System, and migrated to its current neighbourhood. However, this scenario still misses definitive evidence. It is not clear whether Ceres formed in the outer Solar System or formed in situ but accreted materials radially transported from the outer Solar System (McKinnon, 2012). The stable isotope anomalies of samples from Ceres could provide ground truth on where it formed (Fig. 2). Furthermore, geochemical information in samples could be used to understand the evolution path of Ceres.




**Fig. 2** Dichotomy of stable isotopes among planetary materials. (Figure reprinted from Warren, 2011).

## 2.2 Results from the Dawn mission

The Dawn mission performed a 3.5 years rendezvous with Ceres starting from early 2015, mapping its topography and geomorphology at resolutions down to  $\sim 35$  m globally, surface mineralogy at resolutions down to 90 m, the top layer ( $\sim 0.5$ -1 m) elemental abundance at effective resolutions of tens of km, as well as the gravity field to 18 degrees of spherical harmonics. During the last mission phase, Dawn entered a long elliptical orbit with an altitude of 35 km at periastris for a detailed study of the narrow longitudinal strip at about  $240^\circ\text{E}$  that stretches from north of Occator crater and goes up to Azacca crater on the other hemisphere. Images were acquired with resolutions as high as 3.5 m/pixel.

Dawn results confirmed that Ceres is volatile-rich, has a partially differentiated interior, and has experienced global aqueous alteration. Dawn revealed Ceres as a geologically active dwarf planet with brine-driven volcanism and geothermal activity as recently as of a few Myr and probably even at present. Ammoniated phyllosilicates are distributed all over the surface of Ceres (De Sanctis et al., 2015; Ammannito et al., 2016). The incorporation of ammonia in Ceres' surface mineralogy is an indication that Ceres may have accreted materials from the giant planet formation region.

The surface layer of Ceres is rich in water of hydration and water ice. The abundance of water ice is relatively low towards the equator and high towards the polar regions (Prettyman et al., 2017). The latitudinal variation is consistent with the evolution of subsurface water ice as controlled by the thermal condition on the surface and shallow subsurface of Ceres (Schorghofer, 2008, 2016) and the low obliquity of Ceres (Russell et al., 2016). Water ice has been directly identified in about ten specific places on the surface, near rim shadows in fresh craters at latitudes poleward of  $30^\circ$  (Combe et al., 2016, 2019), and inside the permanently shadowed craters in the polar regions (Platz




**Fig. 3** Comparison view of water abundances on Earth, Mars, and Ceres.

et al., 2016; Schorghofer et al., 2016). The existence of water ice in the first few km of the crust is also inferred by analyzing lobate morphologies (Schmidt et al., 2017).

The abundant water ice on Ceres (Fig. 3) led to differentiation and aqueous alteration that shaped the mineralogical composition of Ceres in its crust and mantle (Castillo-Rogez et al., 2018). Near-infrared data returned by the Visible and InfraRed mapping spectrometer (VIR) on board the Dawn spacecraft detected carbonates and ammonium salts that are previously found only on Earth and Enceladus (De Sanctis et al., 2016). The existence of brines at depth played a key role in driving the geological process on Ceres, including volcanism and activities in its geologically recent history (Ruesch et al., 2016, 2019; Scully et al., 2019a).

The most prominent geomorphological feature that is considered of volcanic origin is Ahuna Mons (Fig. 4) (Ruesch et al., 2016, 2019). Its distinct size, shape, and morphology are consistent with being a volcanic dome formed by extrusions of highly viscous melt-bearing material. At the summit is the buildup of a brittle carapace, which partially fractured and disintegrated to generate the slope debris. The bright streaks in the slope debris are rich in Na-carbonates (Zambon et al., 2017; Carrozzo et al., 2018; Zambon et al., 2019). The gravitational relaxation of the enclosed ductile core shapes the overall topographic profile of the summit, requiring an extruded material of high viscosity. The age of the most recent activity on Ahuna Mons is about  $210 \pm$




**Fig. 4** Perspective view of Ahuna Mons on Ceres from Dawn Framing Camera data (no vertical exaggeration). The mountain is 4 km high and 17 km wide in this south-looking view (image credit: NASA/JPL/Caltech/IAPS/MPS/DLR/INAF/ASI).

30 Myr (Ruesch et al., 2016). Other possible features involving the presence of melt include smaller domes (Sori et al., 2017, 2018), fractures in the crater floor (Buczkowski et al., 2016), post-impact modification by the deposition of extended plains material with pits and widely dispersed deposits that form a diffuse veneer on the preexisting surface (Krohn et al., 2016).

Shallow subsurface volatiles are also evident from the many other geomorphological features, such as the pitted terrain (Sizemore et al., 2017). On the other hand, the crater morphology and the simple-to-complex crater transition indicate that Ceres’ outer shell is likely neither pure ice nor pure rock, but a mixture of ice, rock, salts and/or clathrates that allows for limited and spatially variable viscous relaxation (Hiesinger et al., 2016; Bland et al., 2016).

The distinctive bright regions within Occator crater are one of the most remarkable features on Ceres observed by Dawn (Fig. 5). Occator crater is about 90 km in diameter, hosting the bright deposits covering the pit-dome complex named Cerealia Facula in the center, and a group of secondary bright deposits named Vinalia Faculae on the east side of the crater floor. While Ceres’ average surface contains Mg- and Ca-carbonates and Mg- and  $\text{NH}_4$ -phyllosilicates, the Occator’s faculae contain Na-carbonate, Al-phyllosilicates, and  $\text{NH}_4$ -chloride (Raponi et al., 2019). Theoretical modelling suggested the possibility of a brine reservoir beneath Occator crater, and the gradual freezing of this reservoir was the driver of briny lavas (Quick et al., 2019). The salts lower the eutectic point and extend the duration of volcanic activity on Ceres. Laboratory studies suggest that a slow freezing process ( $<30$  K/min) of the exposed ammonium-sodium-carbonate-chloride brines is most compatible with the observed composition of brines observed in Occator crater (Thomas et al., 2019). A variety of morphological features are observed in the crater. Linear and concentric fractures on the crater floor are associated with cryomagmatic intrusion, and degassing or desiccation processes for the volatile rich Occator ejecta (Buczkowski et al., 2019). The cross-cutting relationship



**Fig. 5** Top: Perspective view of Occator crater from the south and the bright deposits from Dawn Framing Camera data. The Cerealia Facula is saturated in this brightness stretch. Bottom left: Zoom-in view of the Cerealia Facula at the center of Occator crater shows the details of the bright deposit. Bottom right: Zoom-in view of the Vinalia Faculae (image credit: NASA/JPL/Caltech/IAPS/MPS/DLR/INAF/ASI).

between stratigraphic units indicate that the Cerealia Facula was emplaced in multiple episodes (Scully et al., 2019b). Crater counting in the Occator ejecta yields an age of about 20 Myr (Neesemann et al., 2019), while the age of faculae is only a few Myr (Nathues et al., 2017, 2019).

The presence of water ice deposits on the surface of Ceres, as well as the widespread distribution of shallow subsurface water has been associated with the active outgassing of Ceres previously observed from Earth orbit or the Earth-Sun L2 Lagrange point (A’Hearn and Feldman, 1992; Küppers et al., 2014). However, the observed surface and subsurface water ice on Ceres do not appear to be sufficient to supply the observed water production rate


(Schorghofer et al., 2017; Landis et al., 2017, 2019). Multiple attempts to detect water outgassing around Ceres failed, suggesting that water outgassing from Ceres is variable but also does not appear to depend on heliocentric distance. Variation in the amount of water ice was detected on the wall of the Juling crater, indicating the possible existence of a seasonal water cycle (Raponi et al., 2018). Another hypothesis proposes that the solar energetic particles might be a driver of outgassing (Villarreal et al., 2017). The water loss mechanisms on Ceres and the characteristics of its transient water exosphere are therefore still uncertain.

Gravitational data and geophysical modelling suggested that Ceres is partially differentiated to a  $\sim 40$  km thick and strong crust composed of rock, ice, salts, and/or clathrates with no more than 30% water ice (Ermakov et al., 2017b; Bland et al., 2016; Fu et al., 2017). Below the crust is a denser rocky mantle with a relatively weak upper layer with brine-filled pore space that controls the global shape of Ceres (Fu et al., 2017). The possibility of a dehydrated core below 100 km cannot be ruled out (King et al., 2018).

VIR first detected an organic absorption feature at  $3.4\text{-}\mu\text{m}$  on Ceres. This signature is diagnostic of organic matter and is mainly localized on a broad region of  $\sim 1000\text{ km}^2$  close to the  $\sim 50$  km Ernutet crater (De Sanctis et al., 2017). The shape of the  $3.4\text{-}\mu\text{m}$  band and the lack of an associated  $3.25\text{-}\mu\text{m}$  feature could exclude organics with a high content of aromatic carbon such as anthraxolites as main carriers of the features on Ceres, in favour of hydrocarbons rich in aliphatic carbon (like asphaltite and kerite; De Sanctis et al., 2017). However, a firm identification of the specific organic material responsible for the  $3.4\text{-}\mu\text{m}$  absorption band observed close to crater Ernutet is missing, and the assessment of its nature (endogenous vs. exogenic) is still debated, making this a compelling scientific question for any future space mission to Ceres. Furthermore, organic compounds possibly also exist on top of Cerealia Facula, the brightest spot located roughly in the middle of crater Occator (De Sanctis et al., 2020).

The volcanism and hydrothermal activity in the recent history of Ceres, the existence of brines on a global scale at the present suggest an active planetary body that could have strong astrobiological significance. The occurrence on Ceres of ammonia-bearing hydrated minerals, water ice, carbonates, salts, and organic material, revealed a complex chemical environment, potentially favourable to prebiotic chemistry in a subsurface aqueous environment (De Sanctis et al., 2017). The ability to detect, determine, and quantify any organics on Ceres is a clear step toward assessing habitability (Castillo-Rogez et al., 2020).

Dawn findings have placed Ceres among Solar System “ocean worlds”, bodies that host current liquid ocean (not necessarily global, according to Hendrix et al., 2019). Originally classified as a “candidate” ocean world in NASA’s Roadmap to Ocean Worlds (Hendrix et al., 2019), Ceres is now suggested as a “real” ocean world in the light of latest results from analyses of Dawn data acquired during its extended mission (Castillo-Rogez, 2020 and



**Fig. 6** Summary of Dawn's observations of Ceres addressed in the text. (a) Geophysical data confirmed the abundance of water ice and the need for gas and salt hydrates to explain the observed topography and crustal density. (b) Various types of carbonates and ammonium chloride have been found in many sites across Ceres' surface (e.g., salts exposed on the floor of Dantu crater). (c) Ernutet crater ( $\sim 52$  km, above) and its area present carbon species in three forms (reduced in  $CxHy$  form, oxidized in the form of carbonates and intermediate as graphitic compounds). (d) Ceres shows extensive evidence for water ice in the form of ground ice and exposure via mass wasting and impacts (Left: Juling crater,  $\sim 20$  km). (e) Recent expressions of volcanism point to the combined role of radiogenic heating and low-eutectic brines in preserving melt and driving activity (Left: Ahuna Mons,  $\sim 4.5$  km tall,  $\sim 20$  km diameter). (f) Impacts could create local chemical energy gradients in transient melt reservoirs throughout Ceres' history (Left: Cerealia Facula,  $\sim 14$  km diameter). (Figure reprinted from Castillo-Rogez et al., 2020; credit for individual images: NASA/JPL/Caltech/IAPS/MPS/DLR/INAF/ASI).

references therein). In summary, as illustrated in Fig. 6, Dawn revealed the great scientific significance of Ceres:

- Rich in water ice and other volatiles relevant to our understanding of the history of volatiles and organic matter in the inner Solar System
- Formation and evolutionary history representative of ice-rich objects in the outer Solar System
- Geologically active with minerals present only on Earth and Enceladus
- Closest and most accessible dwarf planet with volcanism and hydrothermal activity
- Ammonia-bearing hydrated minerals, water ice, carbonates, salts, and organic matter make a complex chemical environment that could favour pre-biotic chemistry

### 2.3 The need for Ceres sample return

Vesta has now joined Mars and the Moon as the best understood extraterrestrial bodies, due at least in part to the fact that we have samples of all of them to study in the laboratory. Ceres can join that group with sample return.

Some exploration and scientific arguments for Ceres sample return are:


- Although carbonaceous chondrites provide the best analog for Ceres (McSween et al., 2018), we have no meteorites from Ceres. The value of having samples for proper calibration of flight instruments and rigorous interpretation of remote sensing data is illustrated by studies of Vesta, Mars, and the Moon.
- Global spectral mapping of Ceres by Dawn’s VIR demonstrates that its surface is covered almost everywhere by the same assemblage (ammoniated clay, serpentine, carbonate, and a darkening agent), but in slightly differing proportions (Ammannito et al., 2016). Thus, a representative regolith sample can be collected from nearly any location on the surface.
- The ice table occurs below several meters depth near the equator, and approaches the surface at higher latitudes (Prettyman et al., 2017). Sampling the regolith in the equatorial region would thus not require cryogenic collection and return.
- Ceres is heavily altered by aqueous processes, and is differentiated with separation of ice and rock (McSween et al., 2018). We have never before sampled an ocean world, and our understanding of its alteration is based only on models. Remote sensing, at best, can only identify a few coexisting minerals. Other phases in minor proportions in a returned sample could constrain the conditions of alteration. It is also likely that Ceres’ regolith may contain some amorphous phases that could not be characterized by remote sensing.
- There is presently some controversy about the extent to which Ceres’ surface has been contaminated by exogenic chondrite impactors. These could be recognized and quantified by petrologic examination of a returned sample.
- Ceres’ chronology is based on crater size distribution analysis, which is dependent on the reference impactor flux (Marchi et al., 2010), leaving the absolute ages uncertain. Radiometric dating of a sample from a mapped geologic unit (Williams et al., 2018) could help define Ceres’ chronology and the impactor flux in the asteroid belt.
- At several locations on Ceres, notably Ahuna Mons and Occator Crater, recent brines have erupted and deposited salts (sodium-carbonate, ammonium chloride, plus more phases that have not been identified (De Sanctis et al., 2016; Zambon et al., 2017; Tosi et al., 2018, 2019; Carrozzo et al., 2018). Small quantities of these phases, which provide important constraints on the nature of subsurface fluids, may occur in regolith samples close to such outcrops.

- Organic matter discovered at one location on Ceres (De Sanctis et al., 2017) suggests that it should be widely distributed in lesser amounts, although in some areas it may have been converted to graphite through UV irradiation. The presence of carbon all across Ceres’ surface with abundances greater than in CI chondrites suggests a widespread water alteration process associated with organic chemistry (Prettyman et al., 2018; Marchi et al., 2019). Understanding the organic component on Ceres has important implications for prebiotic chemistry and astrobiology.
- We do not know whether Ceres formed near its present location in the asteroid belt or in the giant planet region and was later scattered into the main belt by giant planet migration and resonances. Measurement of isotopes of H, C, and N will place constraints on the origin of water and organics. Isotopes of oxygen, chromium, titanium, etc. (Scott et al., 2018) can place Ceres into its proper formation setting, as it has for other bodies for which we have samples.
- Comets from the outer Solar System have long been suspected as the source of Earth’s water. However, while a recent deuterium-to-hydrogen (D/H) ratio measurement for a comet is compatible with terrestrial ocean water (Hartogh et al., 2011), most comet D/H measurements are not (Altwegg et al., 2015). Meanwhile, dynamical studies (O’Brien et al., 2018) indicate that large quantities of Earth’s water could have been supplied by objects from the region of the Solar System coinciding with the present-day outer asteroid belt. If Ceres can be determined to have formed in situ, a D/H ratio (in addition to other isotopic ratios) measured for Ceres would be very valuable in helping to assess the plausibility of the main asteroid belt as a source of the terrestrial water that is so critical to the rise of life on Earth.

### 3 Mission scenarios

#### 3.1 Overview

If we follow the long-term trend in the development of scientific missions to the Moon and Mars, respectively, and that of asteroidal exploration, we would probably reach the conclusion that they might likely converge on a large-scale international space program of Ceres. NASA’s Dawn mission has yielded many exciting results, but the scientific observations were limited to just three remote-sensing instruments, namely the framing camera, the visible and near infrared imaging spectrometer, the gamma-ray spectrometer/neutron detector, as well as gravity science. A more comprehensive payload on future orbiter(s) is needed for a full characterization and understanding of the surface and atmospheric/exospheric environment of Ceres. In addition, just like in the case of lunar or Mars exploration, different types of platforms such as lander and rovers might be required to address astrobiology objectives. A sample return mission might be regarded as the final step. Furthermore, the water- and



**Fig. 7** The contribution of future Ceres in-situ and sample return mission to the exploration of the Solar System ocean worlds and their habitability.

organic-rich environment of Ceres makes it an attractive candidate location for future research stations in support of deep space exploration.

As implied by “Dawn”, the name of the first mission to Ceres, and the name of the present mission proposal “GAUSS”, the in-depth exploration of Ceres should be viewed in terms of detailed investigations of the genesis and evolution of the asteroid belt and the Solar System. The contribution of future in-situ and sample return explorations of Ceres in the context of studying the Solar System ocean worlds and their habitability is illustrated in Fig. 7. It can therefore be envisaged that besides orbiter observations, a set of lander(s) and rover(s) would be needed. It is also possible that this program can be composed of a series of missions over several decades to address different specific scientific questions.

The level of ESA engagement could vary between an M-class mission to an L-class mission. For example, even though our main focus is to initiate the planning of an L-class sample return mission, it could be streamlined to a Rosetta/Philae lander style or an ExoMars/Rosalind Franklin rover style mission for cost reason and expediency. The cost envelope might hence be fitted within the budget of an M-class project depending on the mission component to be chosen. Similar consideration would probably be pursued by CNSA also if it agrees to support a joint assessment study. It is important to emphasize

**Table 1** Summary of mission scenarios

| Mission type   | Mission component               | Mission class                                               |
|----------------|---------------------------------|-------------------------------------------------------------|
| Sample return  | Orbiter+lander+return capsule   | L-class, or M-class with significant contribution from CNSA |
| Landing/Roving | Orbiter+lander / hopper / rover | M-class with possible lander/rover contribution by CNSA     |

that a sample return mission might still be possible with the participation of additional national agencies, besides CNSA and ESA as postulated here. Possible mission scenarios are summarized in Table 1.

In the scenario of a cryogenic sample return mission, GAUSS will first perform a high-resolution global remote sensing investigation, characterizing the geophysical and geochemical properties of Ceres. Candidate sampling sites will then be identified, and observation campaigns will be run for an in-depth assessment of the candidate sites. Once the sampling site is selected, a lander will be deployed on the surface to collect samples and return them to Earth in cryogenic conditions that preserve the volatile and organic composition as well as the original physical state enabled by state of the art technologies.


The scientific goals and related measurement objectives of Project GAUSS are summarized in the traceability matrix (Table 2). The four scientific goals focus on understanding: the origin of Ceres, its evolution and current status, its habitability, and its connection to the carbonaceous meteorite collections. The results from these investigations will have direct implications on our understanding of the evolution of the Solar System and in particular of the icy satellites.

### 3.2 Candidate sites for in-situ investigation and sample return


The geomorphological features on the surface of Ceres are connected to its interior and formed through recent geological processes (Fig. 8). The mantle of Ceres is composed of hydrated minerals, and may be enriched of organics and high-density, low melting point organic matter in localized areas. The ~40 km thick crust above the mantle is composed of the original ocean materials, salts, carbonates, and brine. Liquid brine pockets existed in the recent past and drove volcanic activity. Ahuna Mons might be an extrusion feature of briny mud and organics from mantle. The surface layer of crust is covered by a mixture of infalls, salts, and organics. Cerealia Facula is covered by salts that are left behind after subsurface liquid brine was accessed by impact-produced fractures and reached the surface through the cracks and eventually evaporated. The Haulani crater is one of the youngest impact craters, associated with bright blue rays of ejecta that are freshly exposed crustal materials and could be associated with the original ocean materials (Tosi et al., 2018). Therefore, the Dawn observations of Ceres indicates these sites carry profound sci-

**Table 2** Traceability matrix for Project GAUSS

| Scientific goal                                                                                                                           | Measurement objective                                                             | Instrument                                                                                                                                                                 |
|-------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>The origin and transportation of water and other volatiles in the inner Solar System: Where does Ceres come from?</i>                  |                                                                                   |                                                                                                                                                                            |
| Ceres chronology                                                                                                                          | Crater counting, radiometric dating of samples                                    | Sample analysis, camera, topographic camera                                                                                                                                |
| Connection between Ceres composition and pre-solar materials                                                                              | Stable isotopes of oxygen, chromium, titanium                                     | Sampling mechanism, microscopic camera                                                                                                                                     |
| Volatile inventory on Ceres                                                                                                               | Surface and subsurface water, other volatile species including Na, K, S, Cl, etc. | Sampling mechanism, Near-Infrared spectrometer (NIR), Thermal Infrared spectrometer (TIR), Active Particle-induced X-ray Spectrometer (APXS), Gamma Ray Spectrometer (GRS) |
| Contamination from exogenous materials on Ceres and their roles in the evolution of Ceres                                                 | Surface mineralogy, petrological units in samples                                 | Sampling mechanism, camera, NIR, TIR, microscopic camera                                                                                                                   |
| <i>Physical properties and internal structure of Ceres: What do ice dwarf planets look like?</i>                                          |                                                                                   |                                                                                                                                                                            |
| Structure of the near-subsurface and deep interior, and the implications to the differentiation and aqueous alteration processes of Ceres | Surface mineralogy, elemental abundance, gravity                                  | Imaging, NIR, TIR, seismology, radar, subsurface science package                                                                                                           |
| Geological processes, in particular current and past cryovolcanism and geothermal activity                                                | Topography and morphology, gravity                                                | Imaging, lidar, radar, radio science, seismometer                                                                                                                          |
| The existence and characteristics of Ceres' exosphere                                                                                     | Gas species around Ceres                                                          | Gas chromatography-mass spectrometer (GC-MS), Ion and mass spectrometer, UV spectrometer, dust detector                                                                    |
| <i>The astrobiological implications of Ceres: Was it habitable in the past and is it still today?</i>                                     |                                                                                   |                                                                                                                                                                            |
| Existence of liquid water inside Ceres, its extent, distribution, and depth                                                               | Topography, morphology, gravity                                                   | Camera, radar, lidar, topographic camera, radio science, seismometer, subsurface science package                                                                           |
| Redox condition on Ceres, the existence and forms of oxidants                                                                             | Elemental abundance, forms, and isotopic ratios of O, S, Cl, N                    | Sampling mechanism, NIR, TIR, APXS, GRS                                                                                                                                    |
| Abundance, sources and sinks, and chemical forms of life-forming elements; source of terrestrial water                                    | Elemental abundance, forms, and isotopic ratios of C, H, N, O, S                  | Sampling mechanism, NIR, TIR, APXS, GRS                                                                                                                                    |
| Inventory and composition of organic compounds, their origins and evolutions                                                              | Existence, abundance, and types of organic materials                              | Sample mechanism, NIR, TIR                                                                                                                                                 |
| <i>Mineralogical connection between Ceres and the collections of primitive meteorites: Where are Ceres meteorites?</i>                    |                                                                                   |                                                                                                                                                                            |
| Thermal metamorphism and aqueous alteration history of Ceres                                                                              | Mineralogy, petrological characterization, isotope ratios                         | Sampling mechanism, NIR, TIR                                                                                                                                               |
| Fractionation of elements and the geochemical processes                                                                                   | Mineralogy, elemental abundance, isotope ratios                                   | Sampling mechanism, NIR, TIR                                                                                                                                               |



**Fig. 8** Surface geomorphological features on Ceres and their connections to the interior structure (image credit: NASA/JPL/Caltech/IAPS/MPS/DLR/INAF/ASI).



**Fig. 9** Candidate sites on Ceres for in-situ and/or sample return. The Ceres basemap is a composite color map of Ceres generated by Dawn Framing Camera images with R, G, B = 960, 550, 440 nm, respectively (image credit: NASA/JPL/Caltech/IAPS/MPS/DLR/INAF/ASI).

entific implications serving the scientific goals identified for Project GAUSS for in-situ investigations and sample returns (Fig. 9).

### 3.3 Proposed payload

Only three instruments were carried on Dawn. A more comprehensive payload would be needed to characterize Ceres itself and its atmospheric and space environment. This is especially true with the detection of water plume activity by Herschel (Küppers et al., 2014). Similar to the Cassini measurements at Enceladus, repeated fly-through of the gas plume would allow the identification of the chemical composition and isotopic ratios of the gas molecules, thus providing a probe to the nature of the subsurface lake/ocean. The proposed payload and their heritage are summarized in Table 3. It is assumed that some of the scientific instruments will have been successfully developed for CNSA's Tianwen-1 mission to Mars and the China Asteroid Explorer (CAEX) to small bodies. By the same token, the sampling and reentry technology are also assumed to be derived from China's sampling missions to the Moon, Mars, and asteroids. The instrumentation on the Philae lander on ESA's Rosetta, the Rosalind Franklin Rover (RFR) of ExoMars, the Yutu rovers of Chang'e missions, and the Zhurong rover of the Tianwen-1 mission, will provide strong preparation and heritage for the Ceres rover.

It is important to emphasize the astrobiological significance of the in-situ exploration of geological landmarks of special interest. These include the Ernutet crater and the Occator crater. To achieve high confidence in detecting organic materials on the surface of Ceres and assess their nature, the mid-IR range 2-12  $\mu\text{m}$  turns out to be crucial at the spatial resolution that could be achieved by a spacecraft in orbit around Ceres, not necessarily with imaging capabilities. Covering this sensitivity range would allow:

- Complementing surface mineralogy as derived by VIR in the overall 0.4-5.1  $\mu\text{m}$  spectral range by using thermal emission spectroscopy from 2-12  $\mu\text{m}$ , determining the specific compound responsible for the organics-rich area observed close to crater Ernutet, and shedding light on its origin (endogenous vs. exogenic). This spectral range would ultimately provide the capability to greatly expand our inventory of astrobiologically important compounds, and to remotely sense complex organics.
- Assess any potential ongoing activity and occurrence of organics, particularly on top of Cerealia Facula in crater Occator.
- Perform an in-depth characterization of the 34-km crater Haulani, one of the youngest geologic features on Ceres and home to the most prominent thermal signature observed on the entire surface of the dwarf planet (Tosi et al., 2018, 2019).
- Retrieve surface temperatures, grain size, porosity and surface roughness using thermal emission from 2-12  $\mu\text{m}$ , thus accessing temperature values below 180 K with high accuracy, which were precluded to Dawn/VIR.
- Ultimately characterize the best landing site for a surface element in terms of composition and roughness (regolith depth).

The Fourier Transform Spectrometer (FTS) working principle is based on the Michelson interferometer, which is an infrared spectrometer suitable for


**Table 3** Strawman payload on the Orbiter and the Lander

|         | Payload name                         | Heritage                                |
|---------|--------------------------------------|-----------------------------------------|
| Orbiter | Wide-angle and narrow-angle camera   | Chang'e 1-3, Dawn, Rosetta, BepiColombo |
|         | Infrared imaging spectrometer        | Dawn, Rosetta, Chang'e-4, CAEX          |
|         | Fourier Transform Spectrometer       | Mars Express, ExoMars                   |
|         | Thermal mapper                       | CAEX                                    |
|         | Ultraviolet imaging spectrometer     | Chang'e-3, MMX                          |
|         | GRS                                  | Change'e 1-2                            |
|         | Long-wavelength radar                | CAEX                                    |
|         | Lidar                                | BepiColombo                             |
|         | Dust detector                        | Rosetta, CAEX                           |
|         | Ion and mass spectrometer            | Rosetta, CAEX                           |
|         | Particles and Fields package         | Rosetta, CAEX                           |
|         | Radio Science                        | Rosetta                                 |
| Lander  | Topographic camera system            | Chang'e 3-4; Mars rovers; Philae        |
|         | Microscopic camera                   | Philae, CAEX, CLUPI (ExoMars)           |
|         | APXS                                 | Chang'e 1-3                             |
|         | GRS                                  | Chang'e 1-2                             |
|         | Laser-induced breakdown spectroscopy | Tianwen-1, Curiosity                    |
|         | GC-MS                                | Philae, CAEX                            |
|         | Subsurface science package           | Philae, CAEX                            |
|         | Active seismometer                   | Insight, Chang'e 7                      |

covering a broad spectral range from the near infrared to the mid infrared with constant, high spectral resolution (typically up to  $1\text{-}2\text{ cm}^{-1}$ ). FTSs simultaneously detect light varying the optical path difference (OPD) and encoding the signal at each wavelength with a cosine modulation at a frequency proportional to the wavenumber  $\nu=1/\lambda$ . This interferogram is then transformed into a spectrum, i.e., from the optical path domain to the optical frequency domain. The OPD could be produced by rotating a double pendulum system around its axis, rather than translating the moving mirror along a linear direction like in the classical Michelson interferometer. Compared to the first generation of FTS (e.g. PFS on board the ESA Mars Express spacecraft), in recent years updated versions have been proposed that foresee a significant reduction in mass and size, adopting innovative technical solutions.

### 3.4 Trajectory design

Considering the fact that Ceres is located between the orbits of Mars and Jupiter (the semi-major axis of Ceres is about 2.768 au), in the transfer design we take advantage of a gravity assist of Mars in order to reduce the required fuel consumption. In the preliminary transfer scenario, the sequence is given as follows: (a) launching from the Earth; (b) Mars' gravity assist; (c) rendezvous with Ceres; (d) returning to the Earth. The launch window is set to be after



**Fig. 10** The transfer trajectory for Ceres' sample return mission. In the transfer scenario, a Mars gravity assist is taken into consideration in order to save fuel consumption. The probe launches on August 15 th, 2035. The arc from the Earth to Mars is shown in black line; the arc from Mars to Ceres is given in blue line; the rendezvous segment is presented in cyan; and the return trajectory from Ceres to the Earth is marked in green. The key points of time are represented by red dots.

**Table 4** Parameters of the interplanetary trajectory for the Ceres sample return mission.

| Time points           |                   | Velocity impulse                                            |            |
|-----------------------|-------------------|-------------------------------------------------------------|------------|
| Launch                | August 1, 2035    | Hyperbolic excess velocity ( $V_{\infty}$ )                 | 4.567 km/s |
| Mars gravity assist   | August 26, 2038   | Deep space maneuver at perimartian ( $\Delta V_M$ )         | 0.325 km/s |
| Rendezvous with Ceres | July 13, 2040     | Braking velocity for rendezvous with Ceres ( $\Delta V_f$ ) | 4.283 km/s |
| Departure from Ceres  | July 18, 2042     | Accelerating velocity for departing from Ceres              | 4.813 km/s |
| Earth re-entry        | December 13, 2043 |                                                             |            |

January 1 st, 2035. The planar transfer trajectory is illustrated in Fig. 10 and the associated parameters are provided in Table 4.

### 3.5 Technological requirements

Here we discuss the key technological requirements and main challenges for a Ceres sample return mission.

*Flight dynamics* It is assumed, that by the time of the Voyage 2050 programme, the capacity of launch vehicles poses no challenge for sending a spacecraft with a lander and re-launch system to Ceres. Ceres does not have an atmosphere that exhibits any detectable dynamical effects on the Dawn spacecraft at altitudes down to 35 km. Therefore entry into the atmosphere necessary for Mars landing is not needed to land on Ceres. The descent and landing system for Ceres can be derived from the already mature Chinese Chang'e 3 and 4 descent and landing system. The technology for a re-launch system already exists since the Apollo era, and a similar system is already in development for the various Mars sample return missions in study (see Visions and Voyages, the Planetary Science Decadal Survey 2013-2022 report, National Research Council of the National Academies). The escape velocity from Ceres' surface of about 0.5 km/s is much lower than that of the Moon (2.4 km/s) and Mars (5 km/s). Therefore, the re-launch system from Ceres' surface for sample return can be much smaller and more light-weight than those for the Apollo system (which also had a complicated life-support system) or any Mars sample return system in development. It can be expected that the needed technology will have been developed and matured by the 2030-2040 timeframe, and should be much more efficient (small and light-weight) than the currently available system and hence will require a less powerful launch vehicle than required by the present technology. Earth re-entry is already mature in today's technology and is not expected to pose any challenge. In addition, an ion propulsion system, like the one used in the Dawn mission should be much more efficient and might also be adopted to boost the velocity increment.

*Sampling on Ceres* One or more approaches need to be developed for sampling the surface layer of Ceres. Sampling on Ceres is different from that on the Moon or Mars, which have relatively strong surface gravity and an anchoring mechanism may not be necessary. Ceres sampling would be different from that on small asteroids such as the targets of JAXA's Hayabusa and Hayabusa 2, and NASA's OSIRIS-REx, which have loose regolith and micro-gravity allowing for a touch-and-go approach. With its surface gravitational acceleration of  $\sim 0.27 \text{ m/s}^2$ , a robust anchoring mechanism is required for sampling on Ceres, depending on the sampling system. A drilling system is also needed to drill into the relatively strong rock-ice-salt mixture with a density of  $\sim 1.3 \text{ g/cm}^3$  (Ermakov et al., 2017a). A drilling depth of decimetres, but less than one metre is likely needed in order to reach the water-ice rich layers or subsurface ice table (Prettyman et al., 2017). Technologies currently being developed for landing on and sampling asteroids might provide the necessary technological readiness for project GAUSS, e.g., technologies being developed for the Chinese small body mission (Zhang et al., 2017). The various sampling

mechanisms and collection system for Mars sample return currently in development (Magnani et al., 2011; Zacny, 2013) are good references for the future development of a Ceres sample return system. Note that ESA and NASA signed a statement of intent on Mars sample return.

*Cryogenic sample collection, containment, and curation* The idea of cryogenic sample return from icy Solar System bodies has been around for a long time. The Rosetta mission was originally designed to return samples from a comet at cryogenic temperature (Huber and Schwehm, 1991). This is probably the most challenging part of the entire project. To preserve the volatile and organic compounds in their original status, samples should not be thermally or aqueously altered during the collection, storage and transport, and shielded from contamination by terrestrial volatiles upon return. Therefore, the samples need to be collected and sealed in the containers with cryogenic temperature and overpressure. A temperature of  $<\sim 170$  K is required, and the original temperature of subsurface samples (probably  $\sim 140$  K; Schorghofer, 2008) is desired, throughout the entire collection, return, and curation process in order to prevent water ice melting and aqueously alteration of mineral samples. The sample return capsule should have a number of separated and individually sealed containers that can be overpressured to prevent the loss of volatiles such as  $\text{H}_2\text{O}$ , HCN, S-bearing species, and cyanides (Chuang, 2019; Chiang et al., 2019).

### 3.6 Planetary protection

As revealed by the Dawn observations, the surface of Ceres is organic rich, with certain areas exhibiting higher concentration (e.g. the Ernutet crater; De Sanctis et al., 2017). Furthermore, multiple regions on Ceres show signs of recent or even ongoing geothermal activities that could expose fresh materials from Ceres' interior. Although it is unlikely that the organic matter on Ceres is of biological origin, it is necessary to assess planetary protection strategies for any future lander or sample return mission to Ceres. Based on the current guidelines for planetary protection (Haddaji et al., 2018) and synthesized analyses of the Dawn results, we summarize in Table 5 preliminary considerations on planetary protection for different types of mission to Ceres.

## 4 International context

We note that the ESA roadmap to Ceres needs to be constructed within the international context. NASA has recently chosen in situ exploration and sample return concepts at Ceres to be studied in preparation for the 2023-2032 planetary science decadal survey. This study will cover a range of architectures with a focus on long-range mobility and different approaches to sample acquisition and return to Earth. CNSA has also expressed interest in a mission to

**Table 5** Planetary protection categories for different types of missions to Ceres

| Mission type                                                                                                                                                                                                   |                                                        | Category | Note                                                                                           |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------|----------|------------------------------------------------------------------------------------------------|
| Lander/Rover                                                                                                                                                                                                   | Most part of the surface                               | II*      | Like Dawn, the mission is at least Category III if it includes a Mars gravity assist.          |
|                                                                                                                                                                                                                | Occator faculae                                        | IV       |                                                                                                |
| Sample return                                                                                                                                                                                                  | Different possible sites, including organic-rich areas | V        | Determination on the sub-category of being “restricted” or “unrestricted” depends on location. |
| <i>Note:</i> Further information to be found in Castillo-Rogez, J. C. et al. “Planetary Protection Requirements for Future Exploration of Ceres—State of Understanding after the Dawn Mission” (under review). |                                                        |          |                                                                                                |

Ceres beyond its exploration of the Moon, Mars and small bodies (Zou et al., 2014). It is, therefore, essential for ESA to explore cooperative opportunities through joint assessment studies with international collaborators.

This mission concept came about during a round-table discussion at the 4th Lunar and Deep-Space Exploration International Conference between July 22 and 24, 2019, in Zhuhai, China. One important objective of this proposal is therefore to promote scientific cooperation of the Chinese planetary science community with its European counterpart, taking advantage of the momentum of CNSA and the spirit of the “Voyage 2050 Initiative” of ESA. It might begin with a joint assessment study co-sponsored by both agencies or their representatives. The assessment should identify the scientific objectives, technical requirements, mission architecture, and possible division of responsibilities and authorities.

The extensive orbital measurements of the Dawn mission can be used as a robust basis to the full advantage in planning this large-scale mission. Because of the efforts of several agencies including JAXA, NASA, and ESA over the last two decades, the exploration of asteroids has become a major component of deep space missions. On the one hand, sample return missions and the related technology have been championed by JAXA that has successfully executed the Hayabusa mission to the S-type asteroid Itokawa and the Hayabusa 2 mission to a C-type asteroid Ryugu. This line of approach will be extended to the sample return mission (Mars Moon Explorer or MMX) to the Martian moon Phobos and likely a sample return mission to a Jovian Trojan asteroid (Okada et al., 2019). Following the Dawn mission, NASA has demonstrated its strong interest in asteroid in-situ exploration by running the OSIRIS-REx sample-collection project to the B-type asteroid Bennu, and developing two more asteroid rendezvous missions, one to the large M-type asteroid, Psyche, and the other one (Lucy) to a number of Jovian Trojans (flybys). The Chinese space agency, CNSA, has recently issued an AO calling for instrument proposals and international cooperation for its first asteroid sample return mission to a near-Earth asteroid, followed by a rendezvous mission to a main-belt comet (Zhang

et al., 2019). There is also an assessment study for a sample return mission to an inner-belt E-type asteroid. In parallel to this heightened level of activities by JAXA, NASA and CNSA, ESA has approved the daring “Comet Interceptor” project on the heel of the Rosetta/Philae mission.

If the above set of planned (or proposed) space projects can be successfully carried out, then the first phase of reconnaissance/sample-return missions to major phenotypes of small asteroids and short-period comets—as a global enterprise—would be completed by 2030-2035. It is also expected that the basic knowledge gained and technologies introduced would be applied to the next phase of asteroidal exploration—beginning around 2030 if not earlier—with the goal to address the need of planetary defense against asteroid impact hazard, and asteroidal mining for commercial reasons.

The lunar samples returned by Apollo missions half a century ago continue to spur cutting-edge science. We believe that Ceres should be viewed as the next frontier space laboratory to extend our knowledge of the Solar System much further beyond.

***The GAUSS Team (continued):*** John Carter (Institut d’Astrophysique Spatiale), Grégoire Danger (Aix Marseille Univ, CNRS), Julia de Leon (Instituto de Astrofísica de Canarias), Jörn Helbert (DLR Institute of Planetary Research), Xiyun Hou (Nanjing University), Hauke Hussmann (DLR Institute of Planetary Research), Katherine Joy (University of Manchester), Tomas Kohout (University of Helsinki), Alice Lucchetti (INAF-Astronomical Observatory of Padova), David Mimoun (Université de Toulouse), Olga Muñoz (Instituto de Astrofísica de Andalucía), Jose Luis Ortiz (Instituto de Astrofísica de Andalucía), Maurizio Pajola (INAF-Astronomical Observatory of Padova), Antti Penttilä (University of Helsinki), Frank Preusker (DLR Institute of Planetary Research), Ottaviano Ruesch (University of Münster), Pablo Santos-Sanz (Instituto de Astrofísica de Andalucía), Nico Schmedmann (University of Münster), Nicole Schmitz (DLR Institute of Planetary Research), Katrin Stephan (DLR Institute of Planetary Research), Guneshwar Thangjam (NISER), Josep Maria Trigo-Rodríguez (Institute of Space Sciences, CSIC-IEEC), Cecilia Tubiana (Max Planck Institute for Solar System Research), Vassilissa Vinogradoff (Aix Marseille University, CNRS), Liangliang Yu (Macau University of Science and Technology), Francesca Zambon (INAF-IAPS), Yuhui Zhao (Purple Mountain Observatory, CAS)

**Acknowledgements** Part of this work has been carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D0004). J.-Y.L. acknowledges partial support from the Solar System Exploration Research Virtual Institute 2016 (SSERVI16) Cooperative Agreement (grant NNH16ZDA001N), SSERVI-TREX to the Planetary Science Institute. J.A. acknowledges support from the European Research Council Starting Grant 757390 (CAstRA). A.J.C. and G.H.J. acknowledge support from the STFC consolidated grant to UCL-MSSL ST/S000240/1. J.M.T-R. acknowledges support from the Spanish Ministry of Science and Innovation (project PGC2018-097374-B-I00).

## References

- A'Hearn MF, Feldman PD (1992) Water vaporization on Ceres. *Icarus* 98(1):54–60, DOI 10.1016/0019-1035(92)90206-M
- Altwegg K, Balsiger H, Bar-Nun A, Berthelier JJ, Bieler A, Bochslers P, Briois C, Calmonte U, Combi M, De Keyser J, Eberhardt P, Fiethe B, Fuselier S, Gasc S, Gombosi TI, Hansen KC, Hässig M, Jäckel A, Kopp E, Korth A, LeRoy L, Mall U, Marty B, Mousis O, Neefs E, Owen T, Rème H, Rubin M, Sémon T, Tzou CY, Waite H, Wurz P (2015) 67P/Churyumov-Gerasimenko, a Jupiter family comet with a high D/H ratio. *Science* 347(6220):1261952, DOI 10.1126/science.1261952
- Ammannito E, DeSanctis MC, Ciarniello M, Frigeri A, Carrozzo FG, Combe JP, Ehlmann BL, Marchi S, McSween HY, Raponi A, Toplis MJ, Tosi F, Castillo-Rogez JC, Capaccioni F, Capria MT, Fonte S, Giardino M, Jaumann R, Longobardo A, Joy SP, Magni G, McCord TB, McFadden LA, Palomba E, Pieters CM, Polanskey CA, Rayman MD, Raymond CA, Schenk PM, Zambon F, Russell CT (2016) Distribution of phyllosilicates on the surface of Ceres. *Science* 353(6303):aaf4279, DOI 10.1126/science.aaf4279
- Bell CPM, Naylor T, Mayne NJ, Jeffries RD, Littlefair SP (2013) Pre-main-sequence isochrones - II. Revising star and planet formation time-scales. *MNRAS* 434(1):806–831, DOI 10.1093/mnras/stt1075, 1306.3237
- Bland MT, Raymond CA, Schenk PM, Fu RR, Kneissl T, Pasckert JH, Hiesinger H, Preusker F, Park RS, Marchi S, King SD, Castillo-Rogez JC, Russell CT (2016) Composition and structure of the shallow subsurface of Ceres revealed by crater morphology. *Nature Geoscience* 9(7):538–542, DOI 10.1038/ngeo2743
- Bottke J W F, Vokrouhlický D, Rubincam DP, Broz M (2002) The Effect of Yarkovsky Thermal Forces on the Dynamical Evolution of Asteroids and Meteoroids, pp 395–408
- Bottke WF, Durda DD, Nesvorný D, Jedicke R, Morbidelli A, Vokrouhlický D, Levison HF (2005) Linking the collisional history of the main asteroid belt to its dynamical excitation and depletion. *Icarus* 179(1):63–94, DOI 10.1016/j.icarus.2005.05.017
- Bottke Jr WF, Vokrouhlický D, Rubincam DP, Nesvorný D (2006) The yarkovsky and yorp effects: Implications for asteroid dynamics. *Annu Rev Earth Planet Sci* 34:157–191
- Buczkowski DL, Schmidt BE, Williams DA, Mest SC, Scully JEC, Ermakov AI, Preusker F, Schenk P, Otto KA, Hiesinger H, O'Brien D, Marchi S, Sizemore H, Hughson K, Chilton H, Bland M, Byrne S, Schorghofer N, Platz T, Jaumann R, Roatsch T, Sykes MV, Nathues A, De Sanctis MC, Raymond CA, Russell CT (2016) The geomorphology of Ceres. *Science* 353(6303):aaf4332, DOI 10.1126/science.aaf4332
- Buczkowski DL, Scully JEC, Quick L, Castillo-Rogez J, Schenk PM, Park RS, Preusker F, Jaumann R, Raymond CA, Russell CT (2019) Tectonic analysis of fracturing associated with occator crater. *Icarus* 320:49–59, DOI 10.1016/j.icarus.2018.05.012

- Budde G, Burkhardt C, Brennecka GA, Fischer-Gödde M, Kruijer TS, Kleine T (2016) Molybdenum isotopic evidence for the origin of chondrules and a distinct genetic heritage of carbonaceous and non-carbonaceous meteorites. *Earth and Planetary Science Letters* 454:293–303, DOI 10.1016/j.epsl.2016.09.020
- Carrozzo FG, De Sanctis MC, Raponi A, Ammannito E, Castillo-Rogez J, Ehlmann BL, Marchi S, Stein N, Ciarniello M, Tosi F, Capaccioni F, Capria MT, Fonte S, Formisano M, Frigeri A, Giardino M, Longobardo A, Magni G, Palomba E, Zambon F, Raymond CA, Russell CT (2018) Nature, formation, and distribution of carbonates on Ceres. *Science Advances* 4(3):e1701645, DOI 10.1126/sciadv.1701645
- Castillo-Rogez J (2020) Future exploration of Ceres as an ocean world. *Nature Astronomy* 4:732–734, DOI 10.1038/s41550-020-1181-5
- Castillo-Rogez J, Neveu M, McSween HY, Fu RR, Toplis MJ, Prettyman T (2018) Insights into Ceres’s evolution from surface composition. *Meteoritics and Planetary Science* 53(9):1820–1843, DOI 10.1111/maps.13181
- Castillo-Rogez JC, Neveu M, Scully JE, House CH, Quick LC, Bouquet A, Miller K, Bland M, De Sanctis MC, Ermakov A, et al. (2020) Ceres: astrobiological target and possible ocean world. *Astrobiology* 20(2):269–291
- Chiang CC, Kuan YJ, Chuang YL (2019) ACA observations of Ceres’ molecular exosphere. In: 16th AOGS Meeting, vol PS16-A025
- Chuang YL (2019) Submillimeter spectral observations of molecular exosphere of Ceres icy world. In: 16th AOGS Meeting, vol PS16-A025
- Ciesla FJ, Cuzzi JN (2006) The evolution of the water distribution in a viscous protoplanetary disk. *Icarus* 181(1):178–204, DOI 10.1016/j.icarus.2005.11.009, astro-ph/0511372
- Combe JP, McCord TB, Tosi F, Ammannito E, Carrozzo FG, De Sanctis MC, Raponi A, Byrne S, Landis ME, Hughson KHG, Raymond CA, Russell CT (2016) Detection of local H<sub>2</sub>O exposed at the surface of Ceres. *Science* 353(6303):aaf3010, DOI 10.1126/science.aaf3010
- Combe JP, Raponi A, Tosi F, De Sanctis MC, Carrozzo FG, Zambon F, Ammannito E, Hughson KHG, Nathues A, Hoffmann M, Platz T, Thangjam G, Schorghofer N, Schröder S, Byrne S, Landis ME, Ruesch O, McCord TB, Johnson KE, Singh SM, Raymond CA, Russell CT (2019) Exposed H<sub>2</sub>O-rich areas detected on Ceres with the dawn visible and infrared mapping spectrometer. *Icarus* 318:22–41, DOI 10.1016/j.icarus.2017.12.008
- Connelly JN, Amelin Y, Krot AN, Bizzarro M (2008) Chronology of the Solar System’s Oldest Solids. *ApJ* 675(2):L121, DOI 10.1086/533586
- De Sanctis MC, Ammannito E, Raponi A, Marchi S, McCord TB, McSween HY, Capaccioni F, Capria MT, Carrozzo FG, Ciarniello M, Longobardo A, Tosi F, Fonte S, Formisano M, Frigeri A, Giardino M, Magni G, Palomba E, Turrini D, Zambon F, Combe JP, Feldman W, Jaumann R, McFadden LA, Pieters CM, Prettyman T, Toplis M, Raymond CA, Russell CT (2015) Ammoniated phyllosilicates with a likely outer Solar System origin on (1) Ceres. *Nature* 528(7581):241–244, DOI 10.1038/nature16172

- De Sanctis MC, Raponi A, Ammannito E, Ciarniello M, Toplis MJ, McSween HY, Castillo-Rogez JC, Ehlmann BL, Carrozzo FG, Marchi S, Tosi F, Zambon F, Capaccioni F, Capria MT, Fonte S, Formisano M, Frigeri A, Giardino M, Longobardo A, Magni G, Palomba E, McFadden LA, Pieters CM, Jaumann R, Schenk P, Mugnuolo R, Raymond CA, Russell CT (2016) Bright carbonate deposits as evidence of aqueous alteration on (1) Ceres. *Nature* 536(7614):54–57, DOI 10.1038/nature18290
- De Sanctis MC, Ammannito E, McSween HY, Raponi A, Marchi S, Capaccioni F, Capria MT, Carrozzo FG, Ciarniello M, Fonte S, Formisano M, Frigeri A, Giardino M, Longobardo A, Magni G, McFadden LA, Palomba E, Pieters CM, Tosi F, Zambon F, Raymond CA, Russell CT (2017) Localized aliphatic organic material on the surface of Ceres. *Science* 355(6326):719–722, DOI 10.1126/science.aaj2305
- De Sanctis MC, Ammannito E, Raponi A, Frigeri A, Ferrari M, Carrozzo FG, Ciarniello M, Formisano M, Rousseau B, Tosi F, Zambon F, Raymond CA, Russell CT (2020) Fresh emplacement of hydrated sodium chloride on Ceres from ascending salty fluids. *Nature Astronomy* 4:786–793, DOI 10.1038/s41550-020-1138-8
- DeMeo FE, Carry B (2013) The taxonomic distribution of asteroids from multi-filter all-sky photometric surveys. *Icarus* 226(1):723–741, DOI 10.1016/j.icarus.2013.06.027, 1307.2424
- DeMeo FE, Carry B (2014) Solar System evolution from compositional mapping of the asteroid belt. *Nature* 505(7485):629–634, DOI 10.1038/nature12908, 1408.2787
- Ermakov AI, Fu RR, Castillo-Rogez JC, Raymond CA, Park RS, Preusker F, Russell CT, Smith DE, Zuber MT (2017a) Constraints on Ceres’ Internal Structure and Evolution From Its Shape and Gravity Measured by the Dawn Spacecraft. *Journal of Geophysical Research (Planets)* 122(11):2267–2293, DOI 10.1002/2017JE005302
- Ermakov AI, Mazarico E, Schröder SE, Carsenty U, Schorghofer N, Preusker F, Raymond CA, Russell CT, Zuber MT (2017b) Ceres’s obliquity history and its implications for the permanently shadowed regions. *Geophys. Res. Lett.* 44(6):2652–2661, DOI 10.1002/2016GL072250
- Farinella P, Vokrouhlický D, Hartmann WK (1998) Meteorite Delivery via Yarkovsky Orbital Drift. *Icarus* 132(2):378–387, DOI 10.1006/icar.1997.5872
- Fernandez JA, Ip WH (1984) Some dynamical aspects of the accretion of Uranus and Neptune: The exchange of orbital angular momentum with planetesimals. *Icarus* 58(1):109–120, DOI 10.1016/0019-1035(84)90101-5
- Fu RR, Ermakov AI, Marchi S, Castillo-Rogez JC, Raymond CA, Hager BH, Zuber MT, King SD, Bland MT, Cristina De Sanctis M, Preusker F, Park RS, Russell CT (2017) The interior structure of Ceres as revealed by surface topography. *Earth and Planetary Science Letters* 476:153–164, DOI 10.1016/j.epsl.2017.07.053
- Gladman BJ, Migliorini F, Morbidelli A, Zappala V, Michel P, Cellino A, Froeschle C, Levison HF, Bailey M, Duncan M (1997) Dynamical lifetimes of objects injected into asteroid belt resonances. *Science* 277:197–201, DOI

- 10.1126/science.277.5323.197
- Gomes R, Levison HF, Tsiganis K, Morbidelli A (2005) Origin of the cataclysmic Late Heavy Bombardment period of the terrestrial planets. *Nature* 435(7041):466–469, DOI 10.1038/nature03676
- Gradie J, Tedesco E (1982) Compositional Structure of the Asteroid Belt. *Science* 216(4553):1405–1407, DOI 10.1126/science.216.4553.1405
- Grant R (1852) History of Physical Astronomy, from the earliest ages to the middle of the XIXth Century: Comprehending a detailed account of the establishment of the theory of gravitation by Newton, and its developement by his successors; with an exposition of the progress of research on all the other subjects of celestial physics. Bohn, URL <https://books.google.de/books?id=1iUuAAAACAAJ>
- Haddaji A, Spry JA, Walter N, Brucato JR, Rabbow E, Rettberg P, Kminek G, Margheritis DB, McKenna-Lawlor S, Fellous JL, Yano H, Cabezas P, Sephton M (2018) The International Planetary Protection Handbook (IPPH). In: 42nd COSPAR Scientific Assembly, vol 42, pp PPP.3–14–18
- Hahn JM, Malhotra R (1999) Orbital Evolution of Planets Embedded in a Planetesimal Disk. *AJ* 117(6):3041–3053, DOI 10.1086/300891, *astro-ph/9902370*
- Hargrove KD, Emery JP, Campins H, Kelley MSP (2015) Asteroid (90) Antiope: Another icy member of the Themis family? *Icarus* 254:150–156, DOI 10.1016/j.icarus.2015.03.008
- Hartogh P, Lis DC, Bockelée-Morvan D, de Val-Borro M, Biver N, Küppers M, Emprechtinger M, Bergin EA, Crovisier J, Rengel M, Moreno R, Szuwacz S, Blake GA (2011) Ocean-like water in the Jupiter-family comet 103P/Hartley 2. *Nature* 478(7368):218–220, DOI 10.1038/nature10519
- Hendrix AR, Hurford TA, Barge LM, Bland MT, Bowman JS, Brinckerhoff W, Buratti BJ, Cable ML, Castillo-Rogez J, Collins GC, Diniega S, German CR, Hayes AeG, Hoehler T, Hosseini S, Howett CJA, McEwen AS, Neish CD, Neveu M, Nordheim TA, Patterson GW, Patthoff DA, Phillips C, Rhoden A, Schmidt BE, Singer KN, Soderblom JM, Vance SD (2019) The NASA Roadmap to Ocean Worlds. *Astrobiology* 19(1):1–27, DOI 10.1089/ast.2018.1955
- Hiesinger H, Marchi S, Schmedemann N, Schenk P, Pasckert JH, Neesemann A, O’Brien DP, Kneissl T, Ermakov AI, Fu RR, Bland MT, Nathues A, Platz T, Williams DA, Jaumann R, Castillo-Rogez JC, Ruesch O, Schmidt B, Park RS, Preusker F, Buczkowski DL, Russell CT, Raymond CA (2016) Cratering on Ceres: Implications for its crust and evolution. *Science* 353(6303):aaf4758, DOI 10.1126/science.aaf4759
- Hillenbrand LA (2008) Disk-dispersal and planet-formation timescales. *Physica Scripta Volume T* 130:014024, DOI 10.1088/0031-8949/2008/T130/014024, 0805.0386
- Hiroi T, Zolensky ME, Pieters CM, Lipschutz ME (1996) Thermal metamorphism of the C, G, B, and F asteroids seen from the 0.7 micron, 3 micron and UV absorption strengths in comparison with carbonaceous chondrites. *Meteoritics and Planetary Science* 31(3):321–327, DOI 10.1111/j.1945-

- 5100.1996.tb02068.x
- Hsieh HH, Jewitt D (2006) A Population of Comets in the Main Asteroid Belt. *Science* 312(5773):561–563, DOI 10.1126/science.1125150
- Huber MCE, Schwehm G (1991) Comet Nucleus Sample Return - Plans and Capabilities. *Space Sci. Rev.* 56(1-2):109–115, DOI 10.1007/BF00178398
- Jacobson SA, Morbidelli A, Raymond SN, O’Brien DP, Walsh KJ, Rubie DC (2014) Highly siderophile elements in Earth’s mantle as a clock for the Moon-forming impact. *Nature* 508(7494):84–87, DOI 10.1038/nature13172, 1504.01421
- Jones TD, Lebofsky LA, Lewis JS, Marley MS (1990) The composition and origin of the C, P, and D asteroids: Water as a tracer of thermal evolution in the outer belt. *Icarus* 88(1):172–192, DOI 10.1016/0019-1035(90)90184-B
- Keil K (2000) Thermal alteration of asteroids: evidence from meteorites. *Planet. Space Sci.* 48(10):887–903, DOI 10.1016/S0032-0633(00)00054-4
- King SD, Castillo-Rogez JC, Toplis MJ, Bland MT, Raymond CA, Russell CT (2018) Ceres internal structure from geophysical constraints. *Meteoritics and Planetary Science* 53(9):1999–2007, DOI 10.1111/maps.13063
- Kokubo E, Ida S (1995) Orbital evolution of protoplanets embedded in a swarm of planetesimals. *Icarus* 114(2):247–257, DOI 10.1006/icar.1995.1059
- Krohn K, Jaumann R, Stephan K, Otto KA, Schmedemann N, Wagner RJ, Matz KD, Tosi F, Zambon F, Gathen I, Schulzeck F, Schröder SE, Buczkowski DL, Hiesinger H, McSween HY, Pieters CM, Preusker F, Roatsch T, Raymond CA, Russell CT, Williams DA (2016) Cryogenic flow features on Ceres: Implications for crater-related cryovolcanism. *Geophys. Res. Lett.* 43(23):11,994–12,003, DOI 10.1002/2016GL070370
- Kruijer TS, Burkhardt C, Budde G, Kleine T (2017) Age of Jupiter inferred from the distinct genetics and formation times of meteorites. *Proceedings of the National Academy of Science* 114(26):6712–6716, DOI 10.1073/pnas.1704461114
- Küppers M, O’Rourke L, Bockelée-Morvan D, Zakharov V, Lee S, von Allmen P, Carry B, Teyssier D, Marston A, Müller T, Crovisier J, Barucci MA, Moreno R (2014) Localized sources of water vapour on the dwarf planet (1)Ceres. *Nature* 505(7484):525–527, DOI 10.1038/nature12918
- Landis ME, Byrne S, Schörghofer N, Schmidt BE, Hayne PO, Castillo-Rogez J, Sykes MV, Combe JP, Ermakov AI, Prettyman TH, Raymond CA, Russell CT (2017) Conditions for Sublimating Water Ice to Supply Ceres’ Exosphere. *Journal of Geophysical Research (Planets)* 122(10):1984–1995, DOI 10.1002/2017JE005335
- Landis ME, Byrne S, Combe JP, Marchi S, Castillo-Rogez J, Sizemore HG, Schörghofer N, Prettyman TH, Hayne PO, Raymond CA, Russell CT (2019) Water Vapor Contribution to Ceres’ Exosphere From Observed Surface Ice and Postulated Ice-Exposing Impacts. *Journal of Geophysical Research (Planets)* 124(1):61–75, DOI 10.1029/2018JE005780
- Lecar M, Podolak M, Sasselov D, Chiang E (2006) On the Location of the Snow Line in a Protoplanetary Disk. *ApJ* 640(2):1115–1118, DOI 10.1086/500287, astro-ph/0602217

- Levison HF, Bottke WF, Gounelle M, Morbidelli A, Nesvorný D, Tsiganis K (2009) Contamination of the asteroid belt by primordial trans-Neptunian objects. *Nature*460(7253):364–366, DOI 10.1038/nature08094
- Levison HF, Morbidelli A, Tsiganis K, Nesvorný D, Gomes R (2011) Late Orbital Instabilities in the Outer Planets Induced by Interaction with a Self-gravitating Planetesimal Disk. *AJ*142(5):152, DOI 10.1088/0004-6256/142/5/152
- Magnani P, Re E, Fumagalli A, Senese S, Ori G, Gily A, Baglioni P (2011) Testing of exomars em drill tool in mars analogous materials. Proceedings Advanced Space Technologies for Robotics and Automation (ASTRA) Noordwijk, The Netherlands: European Space Agency
- Marchi S, Barbieri C, Küppers M, Marzari F, Davidsson B, Keller HU, Besse S, Lamy P, Mottola S, Massironi M, Cremonese G (2010) The cratering history of asteroid (2867) Steins. *Planet. Space Sci.*58(9):1116–1123, DOI 10.1016/j.pss.2010.03.017, 1003.5655
- Marchi S, Raponi A, Prettyman TH, De Sanctis MC, Castillo-Rogez J, Raymond CA, Ammannito E, Bowling T, Ciarniello M, Kaplan H, Palomba E, Russell CT, Vinogradoff V, Yamashita N (2019) An aqueously altered carbon-rich Ceres. *Nature Astronomy* 3:140–145, DOI 10.1038/s41550-018-0656-0
- McKinnon WB (2012) Where Did Ceres Accrete - In Situ in the Asteroid Belt, Among the Giant Planets, or in the Primordial Transneptunian Belt? In: AAS/Division for Planetary Sciences Meeting Abstracts #44, AAS/Division for Planetary Sciences Meeting Abstracts, p 111.14
- McSween HY, Emery JP, Rivkin AS, Toplis MJ, Castillo-Rogez JC, Prettyman TH, De Sanctis MC, Pieters CM, Raymond CA, Russell CT (2018) Carbonaceous chondrites as analogs for the composition and alteration of Ceres. *Meteoritics and Planetary Science* 53(9):1793–1804, DOI 10.1111/maps.12947
- Morbidelli A, Levison HF, Tsiganis K, Gomes R (2005) Chaotic capture of Jupiter’s Trojan asteroids in the early Solar System. *Nature*435(7041):462–465, DOI 10.1038/nature03540
- Morbidelli A, Bitsch B, Crida A, Gounelle M, Guillot T, Jacobson S, Johansen A, Lambrechts M, Lega E (2016) Fossilized condensation lines in the Solar System protoplanetary disk. *Icarus*267:368–376, DOI 10.1016/j.icarus.2015.11.027, 1511.06556
- Nanne JAM, Nimmo F, Cuzzi JN, Kleine T (2019) Origin of the non-carbonaceous-carbonaceous meteorite dichotomy. *Earth and Planetary Science Letters* 511:44–54, DOI 10.1016/j.epsl.2019.01.027
- Nathues A, Platz T, Thangjam G, Hoffmann M, Mengel K, Cloutis EA, Le Corre L, Reddy V, Kallisch J, Crown DA (2017) Evolution of Occator Crater on (1) Ceres. *AJ*153(3):112, DOI 10.3847/1538-3881/153/3/112
- Nathues A, Platz T, Thangjam G, Hoffmann M, Scully JEC, Stein N, Ruesch O, Mengel K (2019) Occator crater in color at highest spatial resolution. *Icarus*320:24–38, DOI 10.1016/j.icarus.2017.12.021
- Neesemann A, van Gasselt S, Schmedemann N, Marchi S, Walter SHG, Preusker F, Michael GG, Kneissl T, Hiesinger H, Jaumann R, Roatsch

- T, Raymond CA, Russell CT (2019) The various ages of Occator crater, Ceres: Results of a comprehensive synthesis approach. *Icarus* 320:60–82, DOI 10.1016/j.icarus.2018.09.006
- O'Brien DP, Izidoro A, Jacobson SA, Raymond SN, Rubie DC (2018) The Delivery of Water During Terrestrial Planet Formation. *Space Sci. Rev.* 214(1):47, DOI 10.1007/s11214-018-0475-8, 1801.05456
- Okada T, Iwata T, Matsumoto J, Chujo T, Kebukawa Y, Ito M, Aoki J, Kawai Y, Yokota S, Saito S, Terada K, Toyoda M, Yabuta H, Yurimoto H, Matsuura S, Tsumura K, Yonetoku D, Mihara T, Matsuoka A, Nomura R, Yano H, Hirai T, Nakamura R, Ulamec S, Jaumann R, Bibring JP, Grand N, Szopa C, Palomba E, Helbert J, Herique A, Kumamoto A, Grott M, Auster HU, Klingelhofer G, Yoshida F, Yoshikawa M, Matsushita M, Saiki T, Kato H, Mori O, Kawaguchi J (2019) OKEANOS - A Solar Power Sail Mission to a Jupiter Trojan Asteroid and Its Updated Science Mission Proposal. In: Lunar and Planetary Science Conference, Lunar and Planetary Science Conference, p 1305
- Platz T, Nathues A, Schorghofer N, Preusker F, Mazarico E, Schröder SE, Byrne S, Kneissl T, Schmedemann N, Combe JP, Schäfer M, Thangjam GS, Hoffmann M, Gutierrez-Marques P, Landis ME, Dietrich W, Ripken J, Matz KD, Russell CT (2016) Surface water-ice deposits in the northern shadowed regions of Ceres. *Nature Astronomy* 1:0007, DOI 10.1038/s41550-016-0007
- Poole GM, Rehkämper M, Coles BJ, Goldberg T, Smith CL (2017) Nucleosynthetic molybdenum isotope anomalies in iron meteorites - new evidence for thermal processing of solar nebula material. *Earth and Planetary Science Letters* 473:215–226, DOI 10.1016/j.epsl.2017.05.001
- Prettyman TH, Yamashita N, Toplis MJ, McSween HY, Schörghofer N, Marchi S, Feldman WC, Castillo-Rogez J, Forni O, Lawrence DJ, Ammannito E, Ehlmann BL, Sizemore HG, Joy SP, Polanskey CA, Rayman MD, Raymond CA, Russell CT (2017) Extensive water ice within Ceres' aqueously altered regolith: Evidence from nuclear spectroscopy. *Science* 355(6320):55–59, DOI 10.1126/science.aah6765
- Prettyman TH, Yamashita N, Ammannito E, Castillo-Rogez JC, Ehlmann BL, McSween HY, Marchi S, Pieters CM, Schorghofer N, Toplis MJ, Russell CT, Raymond CA (2018) Carbon on Ceres: Implications for Origins and Interior Evolution. In: Lunar and Planetary Science Conference, Lunar and Planetary Science Conference, p 1151
- Qin L, Nittler LR, Alexander CMOD, Wang J, Stadermann FJ, Carlson RW (2011) Extreme  $^{54}\text{Cr}$ -rich nano-oxides in the CI chondrite Orgueil - Implication for a late supernova injection into the solar system. *Geochim. Cosmochim. Acta* 75(2):629–644, DOI 10.1016/j.gca.2010.10.017, 1101.4949
- Quick LC, Buczkowski DL, Ruesch O, Scully JEC, Castillo-Rogez J, Raymond CA, Schenk PM, Sizemore HG, Sykes MV (2019) A Possible Brine Reservoir Beneath Occator Crater: Thermal and Compositional Evolution and Formation of the Cerealia Dome and Vinalia Faculae. *Icarus* 320:119–135, DOI 10.1016/j.icarus.2018.07.016

- Raponi A, De Sanctis MC, Frigeri Ar, Ammannito E, Ciarniello M, Formisano M, Combe JP, Magni G, Tosi F, Carrozzo FG, Fonte S, Giardino M, Joy SP, Polanskey CA, Rayman MD, Capaccioni F, Capria MT, Longobardo A, Palomba E, Zambon F, Raymond CA, Russell CT (2018) Variations in the amount of water ice on Ceres' surface suggest a seasonal water cycle. *Science Advances* 4(3):eaao3757, DOI 10.1126/sciadv.aao3757
- Raponi A, De Sanctis MC, Carrozzo FG, Ciarniello M, Castillo-Rogez JC, Ammannito E, Frigeri A, Longobardo A, Palomba E, Tosi F, Zambon F, Raymond CA, Russell CT (2019) Mineralogy of Occator crater on Ceres and insight into its evolution from the properties of carbonates, phyllosilicates, and chlorides. *Icarus* 320:83–96, DOI 10.1016/j.icarus.2018.02.001
- Raymond SN, Izidoro A (2017) Origin of water in the inner Solar System: Planetesimals scattered inward during Jupiter and Saturn's rapid gas accretion. *Icarus* 297:134–148, DOI 10.1016/j.icarus.2017.06.030, 1707.01234
- Rivkin AS, Emery JP (2010) Detection of ice and organics on an asteroidal surface. *Nature* 464(7293):1322–1323, DOI 10.1038/nature09028
- Rivkin AS, Howell ES, Vilas F, Lebofsky LA (2002) Hydrated Minerals on Asteroids: The Astronomical Record, pp 235–253
- Rivkin AS, Thomas CA, Trilling DE, Enga Mt, Grier JA (2011) Ordinary chondrite-like colors in small Koronis family members. *Icarus* 211(2):1294–1297, DOI 10.1016/j.icarus.2010.11.033
- Ruesch O, Platz T, Schenk P, McFadden LA, Castillo-Rogez JC, Quick LC, Byrne S, Preusker F, O'Brien DP, Schmedemann N, Williams DA, Li JY, Bland MT, Hiesinger H, Kneissl T, Neesemann A, Schaefer M, Pasckert JH, Schmidt BE, Buzckowski DL, Sykes MV, Nathues A, Roatsch T, Hoffmann M, Raymond CA, Russell CT (2016) Cryovolcanism on Ceres. *Science* 353(6303):aaf4286, DOI 10.1126/science.aaf4286
- Ruesch O, Genova A, Neumann W, Quick LC, Castillo-Rogez JC, Raymond CA, Russell CT, Zuber MT (2019) Slurry extrusion on Ceres from a convective mud-bearing mantle. *Nature Geoscience* 12(7):505–509, DOI 10.1038/s41561-019-0378-7
- Russell CT, Raymond CA, Ammannito E, Buzckowski DL, De Sanctis MC, Hiesinger H, Jaumann R, Konopliv AS, McSween HY, Nathues A, Park RS, Pieters CM, Prettyman TH, McCord TB, McFadden LA, Mottola S, Zuber MT, Joy SP, Polanskey C, Rayman MD, Castillo-Rogez JC, Chi PJ, Combe JP, Ermakov A, Fu RR, Hoffmann M, Jia YD, King SD, Lawrence DJ, Li JY, Marchi S, Preusker F, Roatsch T, Ruesch O, Schenk P, Villarreal MN, Yamashita N (2016) Dawn arrives at Ceres: Exploration of a small, volatile-rich world. *Science* 353(6303):1008–1010, DOI 10.1126/science.aaf4219
- Sasselov DD, Lecar M (2000) On the Snow Line in Dusty Protoplanetary Disks. *ApJ* 528(2):995–998, DOI 10.1086/308209, astro-ph/9911390
- Schmidt BE, Hughson KHG, Chilton HT, Scully JEC, Platz T, Nathues A, Sizemore H, Bland MT, Byrne S, Marchi S, O'Brien DP, Schorghofer N, Hiesinger H, Jaumann R, Pasckert JH, Lawrence JD, Buzckowski D, Castillo-Rogez JC, Sykes MV, Schenk PM, Desanctis MC, Mitri G, Formisano M, Li JY, Reddy V, Lecorre L, Russell CT, Raymond CA (2017)

- Geomorphological evidence for ground ice on dwarf planet Ceres. *Nature Geoscience* 10(5):338–343, DOI 10.1038/ngeo2936
- Schorghofer N (2008) The lifetime of ice on main belt asteroids. *The Astrophysical Journal* 682(1):697
- Schorghofer N (2016) Predictions of depth-to-ice on asteroids based on an asynchronous model of temperature, impact stirring, and ice loss. *Icarus* 276:88–95, DOI 10.1016/j.icarus.2016.04.037
- Schorghofer N, Mazarico E, Platz T, Preusker F, Schröder SE, Raymond CA, Russell CT (2016) The permanently shadowed regions of dwarf planet Ceres. *Geophys. Res. Lett.* 43(13):6783–6789, DOI 10.1002/2016GL069368
- Schorghofer N, Byrne S, Landis ME, Mazarico E, Prettyman TH, Schmidt BE, Villarreal MN, Castillo-Rogez J, Raymond CA, Russell CT (2017) The Putative Cerean Exosphere. *ApJ* 850(1):85, DOI 10.3847/1538-4357/aa932f
- Scott ERD, Taylor GJ, Newsom HE, Herbert FL, Zolensky M, Kerridge JF (1989) Chemical, thermal and impact processing of asteroids. In: Binzel RP, Gehrels T, Matthews MS (eds) *Asteroids II*, pp 701–739
- Scott ERD, Krot AN, Sanders IS (2018) Isotopic Dichotomy Among Meteorites and Implications for the Evolution of the Protoplanetary Disk. In: *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, p 1713
- Scully JE, Bowling T, Bu C, Buczkowski DL, Longobardo A, Nathues A, Neesemann A, Palomba E, Quick LC, Raponi A, et al. (2019a) Synthesis of the special issue: The formation and evolution of Ceres' Occator crater. *Icarus* 320:213–225
- Scully JE, Buczkowski DL, Raymond CA, Bowling T, Williams DA, Neesemann A, Schenk PM, Castillo-Rogez JC, Russell CT (2019b) Ceres' Occator crater and its faculae explored through geologic mapping. *Icarus* 320:7–23
- Sizemore HG, Platz T, Schorghofer N, Prettyman TH, De Sanctis MC, Crown DA, Schmedemann N, Neesemann A, Kneissl T, Marchi S, Schenk PM, Bland MT, Schmidt BE, Hughson KHG, Tosi F, Zambon F, Mest SC, Yingst RA, Williams DA, Russell CT, Raymond CA (2017) Pitted terrains on (1) Ceres and implications for shallow subsurface volatile distribution. *Geophys. Res. Lett.* 44(13):6570–6578, DOI 10.1002/2017GL073970
- Sori MM, Byrne S, Bland MT, Bramson AM, Ermakov AI, Hamilton CW, Otto KA, Ruesch O, Russell CT (2017) The vanishing cryovolcanoes of Ceres. *Geophys. Res. Lett.* 44(3):1243–1250, DOI 10.1002/2016GL072319
- Sori MM, Sizemore HG, Byrne S, Bramson AM, Bland MT, Stein NT, Russell CT (2018) Cryovolcanic rates on Ceres revealed by topography. *Nature Astronomy* 2:946–950, DOI 10.1038/s41550-018-0574-1
- Takir D, Emery JP (2012) Outer Main Belt asteroids: Identification and distribution of four 3- $\mu$ m spectral groups. *Icarus* 219(2):641–654, DOI 10.1016/j.icarus.2012.02.022
- Teets D, Whitehead K (1999) The discovery of Ceres: How Gauss became famous. *Mathematics Magazine* 72(2):83–93
- Thomas EC, Vu TH, Hodyss R, Johnson PV, Choukroun M (2019) Kinetic effect on the freezing of ammonium-sodium-carbonate-chloride brines and

- implications for the origin of Ceres' bright spots. *Icarus* 320:150–158, DOI 10.1016/j.icarus.2017.12.038
- Tosi F, Carrozzo FG, Raponi A, De Sanctis MC, Thangjam G, Zambon F, Ciarniello M, Nathues A, Capria MT, Rognini E, Ammannito E, Hoffmann M, Krohn K, Longobardo A, Palomba E, Pieters CM, Stephan K, Raymond CA, Russell CT (2018) Mineralogy and temperature of crater Haulani on Ceres. *Meteoritics and Planetary Science* 53(9):1902–1924, DOI 10.1111/maps.13078
- Tosi F, Carrozzo FG, Zambon F, Ciarniello M, Frigeri A, Combe JP, De Sanctis MC, Hoffmann M, Longobardo A, Nathues A, Raponi A, Thangjam G, Ammannito E, Krohn K, McFadden LA, Palomba E, Pieters CM, Stephan K, Raymond CA, Russell CT, Dawn Science Team (2019) Mineralogical analysis of the Ac-H-6 Haulani quadrangle of the dwarf planet Ceres. *Icarus* 318:170–187, DOI 10.1016/j.icarus.2017.08.012
- Tsiganis K, Gomes R, Morbidelli A, Levison HF (2005) Origin of the orbital architecture of the giant planets of the Solar System. *Nature* 435(7041):459–461, DOI 10.1038/nature03539
- Villarréal MN, Russell CT, Luhmann JG, Thompson WT, Prettyman TH, A'Hearn MF, Küppers M, O'Rourke L, Raymond CA (2017) The Dependence of the Cerean Exosphere on Solar Energetic Particle Events. *ApJ* 838(1):L8, DOI 10.3847/2041-8213/aa66cd
- Walsh KJ, Morbidelli A, Raymond SN, O'Brien DP, Mandell AM (2011) A low mass for Mars from Jupiter's early gas-driven migration. *Nature* 475(7355):206–209, DOI 10.1038/nature10201, 1201.5177
- Warren PH (2011) Stable-isotopic anomalies and the accretionary assemblage of the Earth and Mars: A subordinate role for carbonaceous chondrites. *Earth and Planetary Science Letters* 311(1):93–100, DOI 10.1016/j.epsl.2011.08.047
- Weissman PR (1999) Diversity of Comets: Formation Zones and Dynamical Paths. *Space Sci. Rev.* 90:301–311, DOI 10.1023/A:1005278905101
- Williams DA, Buczkowski DL, Crown DA, Frigeri A, Hughson K, Kneissl T, Krohn K, Mest SC, Pasckert JH, Platz T, Ruesch O, Schulzeck F, Scully JEC, Sizemore HG, Nass A, Jaumann R, Raymond CA, Russell CT (2018) High-Resolution Global Geologic Map of Ceres from NASA Dawn Mission. In: *Planetary Geologic Mappers Annual Meeting*, vol 2066, p 7001
- Zacny K (2013) Drilling and Caching Architecture for the Mars2020 Mission. In: *AGU Fall Meeting Abstracts*, vol 2013, pp P51G–1797
- Zambon F, Raponi A, Tosi F, De Sanctis MC, McFadden LA, Carrozzo FG, Longobardo A, Ciarniello M, Krohn K, Stephan K, Palomba E, Pieters CM, Ammannito E, Russell CT, Raymond CA (2017) Spectral analysis of Ahuna Mons from Dawn mission's visible-infrared spectrometer. *Geophys. Res. Lett.* 44(1):97–104, DOI 10.1002/2016GL071303
- Zambon F, Carrozzo FG, Tosi F, Ciarniello M, Combe JP, Frigeri A, De Sanctis MC, Thangjam G, Nathues A, Hoffmann M, Longobardo A, Stephan K, Raponi A, Ammannito E, Krohn K, McFadden LA, Palomba E, Raymond CA, Russell CT, Dawn Science Team (2019) Mineralogical analysis of

- quadrangle Ac-H-10 Rongo on the dwarf planet Ceres. *Icarus* 318:212–229, DOI 10.1016/j.icarus.2017.09.021
- Zhang T, Zhang W, Wang K, Gao S, Hou L, Ji J, Ding X (2017) Drilling, sampling, and sample-handling system for China’s asteroid exploration mission. *Acta Astronautica* 137:192–204, DOI 10.1016/j.actaastro.2017.04.017
- Zhang X, Huang J, Wang T, Huo Z (2019) ZhengHe - A Mission to a Near-Earth Asteroid and a Main Belt Comet. In: *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, p 1045
- Zou Y, Li W, Ouyang Z (2014) China’s Deep-space Exploration to 2030. *Chinese Journal of Space Science* 34(5):516–517